



wwPDB EM Validation Summary Report ⓘ

Dec 6, 2020 – 07:16 PM EST

EMDB ID : EMD-5257
Title : Phosphorylated smooth muscle heavy meromyosin shows an open conformation linked to activation
Authors : Baumann, B.A.J.; Taylor, D.; Huang, Z.; Tama, F.; Fagnant, P.M.; Trybus, K.M.; Taylor, K.
Deposited on : 2011-01-21
Resolution : 20.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMMapValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev61
Validation Pipeline (wwPDB-VP) : 2.15.1

1 Experimental information

Property	Value	Source
EM reconstruction method	CRYSTALLOGRAPHY	Depositor
Imposed symmetry	2D CRYSTAL, $a=219.3$ Å, $b=174.8$ Å, $c=94.4$ Å, $\gamma=94.4^\circ$, space group=P 2	Depositor
Number of images used	Not provided	
Resolution determination method	Not provided	
CTF correction method	determined using ICE and corrected with CTFAPPLY	Depositor
Microscope	FEI/PHILIPS CM300FEG/T	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	4.0	Depositor
Maximum defocus (nm)	Not provided	
Magnification	24000.0	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	4.404	Depositor
Minimum map value	-4.670	Depositor
Average map value	-0.034	Depositor
Map value standard deviation	1.010	Depositor
Recommended contour level	0.609	Depositor
Map size (Å)	419.0693, 428.995, 85.9985	wwPDB
Map dimensions	169, 81, 35	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.4797, 2.4514, 2.4571	Depositor


2 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-5257. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

2.1 Orthogonal projections [i](#)


2.1.1 Primary map



The images above show the map projected in three orthogonal directions.

2.2 Central slices [i](#)


2.2.1 Primary map



The images above show central slices of the map in three orthogonal directions.

2.3 Largest variance slices [i](#)


2.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

2.4 Orthogonal surface views [i](#)

2.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.609. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


2.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

3 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

3.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

3.2 Volume estimate [i](#)



The volume at the recommended contour level is 1100 nm³; this corresponds to an approximate mass of 993 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

3.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.

4 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.