



wwPDB EM Validation Summary Report ⓘ

Jul 14, 2024 – 09:46 am BST

PDB ID : 8A2Q
EMDB ID : EMD-15101
Title : Structure of the DNA-bound FANCD2-FANCI complex containing phospho-mimetic FANCI
Authors : Passmore, L.A.; Sijacki, T.; Alcon, P.
Deposited on : 2022-06-06
Resolution : 3.53 Å(reported)
Based on initial model : 6TNF

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1475	
2	B	1344	
3	S	44	
4	T	44	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 27812 atoms, of which 12784 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FANCD2.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	1119	Total	C	H	N	O	S	0	0
			12650	4384	5656	1297	1278	35		

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1437	ALA	-	expression tag	UNP F1NP22
A	1438	GLY	-	expression tag	UNP F1NP22
A	1439	ARG	-	expression tag	UNP F1NP22
A	1440	LEU	-	expression tag	UNP F1NP22
A	1441	GLU	-	expression tag	UNP F1NP22
A	1442	VAL	-	expression tag	UNP F1NP22
A	1443	LEU	-	expression tag	UNP F1NP22
A	1444	PHE	-	expression tag	UNP F1NP22
A	1445	GLN	-	expression tag	UNP F1NP22
A	1446	GLY	-	expression tag	UNP F1NP22
A	1447	PRO	-	expression tag	UNP F1NP22
A	1448	TRP	-	expression tag	UNP F1NP22
A	1449	SER	-	expression tag	UNP F1NP22
A	1450	HIS	-	expression tag	UNP F1NP22
A	1451	PRO	-	expression tag	UNP F1NP22
A	1452	GLN	-	expression tag	UNP F1NP22
A	1453	PHE	-	expression tag	UNP F1NP22
A	1454	GLU	-	expression tag	UNP F1NP22
A	1455	LYS	-	expression tag	UNP F1NP22
A	1456	GLY	-	expression tag	UNP F1NP22
A	1457	SER	-	expression tag	UNP F1NP22
A	1458	ALA	-	expression tag	UNP F1NP22
A	1459	GLY	-	expression tag	UNP F1NP22
A	1460	SER	-	expression tag	UNP F1NP22
A	1461	ALA	-	expression tag	UNP F1NP22
A	1462	ALA	-	expression tag	UNP F1NP22
A	1463	GLY	-	expression tag	UNP F1NP22
A	1464	SER	-	expression tag	UNP F1NP22

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1465	GLY	-	expression tag	UNP F1NP22
A	1466	ALA	-	expression tag	UNP F1NP22
A	1467	GLY	-	expression tag	UNP F1NP22
A	1468	TRP	-	expression tag	UNP F1NP22
A	1469	SER	-	expression tag	UNP F1NP22
A	1470	HIS	-	expression tag	UNP F1NP22
A	1471	PRO	-	expression tag	UNP F1NP22
A	1472	GLN	-	expression tag	UNP F1NP22
A	1473	PHE	-	expression tag	UNP F1NP22
A	1474	GLU	-	expression tag	UNP F1NP22
A	1475	LYS	-	expression tag	UNP F1NP22

- Molecule 2 is a protein called Fanconi anemia complementation group I.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	B	1084	Total	C	H	N	O	S	0	0
			14136	4745	6732	1294	1332	33		

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	558	ASP	SER	engineered mutation	UNP B0I564
B	561	ASP	SER	engineered mutation	UNP B0I564
B	567	ASP	THR	engineered mutation	UNP B0I564
B	1340	HIS	-	expression tag	UNP B0I564
B	1341	HIS	-	expression tag	UNP B0I564
B	1342	HIS	-	expression tag	UNP B0I564
B	1343	HIS	-	expression tag	UNP B0I564
B	1344	HIS	-	expression tag	UNP B0I564
B	1345	HIS	-	expression tag	UNP B0I564

- Molecule 3 is a DNA chain called DNA (29-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
3	S	29	Total	C	H	O	P	0	0
			515	145	199	143	28		

- Molecule 4 is a DNA chain called DNA (29-MER).


Mol	Chain	Residues	Atoms					AltConf	Trace
4	T	29	Total	C	H	O	P	0	0
			511	144	197	142	28		

- Molecule 1: FANCD2

[illegible]

- Molecule 2: Fanconi anemia complementation group I


MET	ALA	GLN	ARG	ILE	LEU	GLN	LEU	ALA	ALA	GLU	GLY	SER	PRO	GLU	ARG	LEU	GLN	GLU	ALA	GLN	LEU	G30	D31	M32	A45	K52	P55	CYS	SER	GLN	GLN	S60	L63	H71	H82	V93	G94	L137	SER	ASN	SER	SER	LYS	GLU	SER	LEU	LEU
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	-----	-----	-----	-----	-----	-----	-----	-----	-----



• Molecule 3: DNA (29-MER)



• Molecule 4: DNA (29-MER)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	197436	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.043	Depositor
Minimum map value	-0.014	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.0056	Depositor
Map size (Å)	258.0, 258.0, 258.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.86, 0.86, 0.86	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/7120	0.50	0/9571
2	B	0.28	0/7514	0.53	0/10182
3	S	1.10	1/344 (0.3%)	1.39	4/514 (0.8%)
4	T	0.80	0/341	0.98	0/507
All	All	0.34	1/15319 (0.0%)	0.57	4/20774 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	S	27	DG	C3'-O3'	5.09	1.50	1.44

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	S	27	DG	P-O3'-C3'	9.22	130.76	119.70
3	S	27	DG	C3'-C2'-C1'	-5.69	95.67	102.50
3	S	19	DC	P-O3'-C3'	5.59	126.40	119.70
3	S	27	DG	C4'-C3'-C2'	-5.06	98.55	103.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6994	5656	5789	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	7404	6732	6770	25	0
3	S	316	199	176	0	0
4	T	314	197	173	0	0
All	All	15028	12784	12908	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:295:LEU:HD11	2:B:315:LEU:HD21	1.84	0.59
2:B:1000:SER:OG	2:B:1002:GLN:OE1	2.20	0.59
1:A:780:GLU:N	1:A:780:GLU:OE1	2.38	0.55
1:A:1058:SER:O	1:A:1062:GLN:HG2	2.07	0.55
2:B:432:PHE:CD1	2:B:468:ILE:HD11	2.43	0.54

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1097/1475 (74%)	1059 (96%)	38 (4%)	0	100	100
2	B	1054/1344 (78%)	1022 (97%)	31 (3%)	1 (0%)	51	84
All	All	2151/2819 (76%)	2081 (97%)	69 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	645	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	544/1311 (42%)	542 (100%)	2 (0%)	91	97
2	B	666/1190 (56%)	665 (100%)	1 (0%)	93	98
All	All	1210/2501 (48%)	1207 (100%)	3 (0%)	93	98

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	375	THR
1	A	786	SER
2	B	1085	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-15101. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 120




Y Index: 135




Z Index: 146


6.3.2 Raw map



X Index: 119



Y Index: 135




Z Index: 146


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0056. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_15101_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 279 nm³; this corresponds to an approximate mass of 252 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.283 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.283 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.53	-	-
Author-provided FSC curve	3.53	4.04	3.58
Unmasked-calculated*	3.79	4.33	3.84

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-15101 and PDB model 8A2Q. Per-residue inclusion information can be found in section [3](#) on page [5](#).


9.1 Map-model overlay [i](#)



X



Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0056 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0056).

9.4 Atom inclusion [i](#)



At the recommended contour level, 96% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0056) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.9510	<div></div> 0.4020
A	<div></div> 0.9810	<div></div> 0.4190
B	<div></div> 0.9620	<div></div> 0.4110
S	<div></div> 0.4560	<div></div> 0.0950
T	<div></div> 0.4550	<div></div> 0.0990

