



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 08:43 AM EDT

PDB ID : 7ACA  
Title : CRYSTAL STRUCTURE OF AN ACTIVE KRAS G12D (GPPCP) DIMER  
IN COMPLEX WITH BI-5747  
Authors : Kessler, D.  
Deposited on : 2020-09-10  
Resolution : 1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	<div> <div>3%</div> <div>95%</div> <div>••</div> </div>
1	B	170	<div> <div>5%</div> <div>92%</div> <div>6% •</div> </div>
1	C	170	<div> <div>12%</div> <div>87%</div> <div>8% 5%</div> </div>
1	D	170	<div> <div>10%</div> <div>93%</div> <div>5% •</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTPase KRas.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	167	Total	C	N	O	S	0	0	0
			1336	836	229	264	7			
1	B	167	Total	C	N	O	S	0	0	0
			1336	836	229	264	7			
1	C	162	Total	C	N	O	S	0	0	0
			1295	811	224	253	7			
1	D	167	Total	C	N	O	S	0	0	0
			1336	836	229	264	7			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P01116
A	12	ASP	GLY	engineered mutation	UNP P01116
B	0	GLY	-	expression tag	UNP P01116
B	12	ASP	GLY	engineered mutation	UNP P01116
C	0	GLY	-	expression tag	UNP P01116
C	12	ASP	GLY	engineered mutation	UNP P01116
D	0	GLY	-	expression tag	UNP P01116
D	12	ASP	GLY	engineered mutation	UNP P01116

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		


- # GCP
- 
- The image displays the chemical structure of Guanosine Cyclic Phosphate (GCP). The structure consists of a guanine base (a purine ring system with an amino group at position 2) linked to a ribose sugar, which is in turn linked to a cyclic phosphate group. The guanine base is shown with atoms labeled N1, N2, N3, N7, N8, C2, C3, C4, C5, C6, and C8. The ribose sugar is shown with atoms labeled C1', C2', C3', C4', and C5'. The cyclic phosphate group is shown with atoms labeled P1, P2, P3, P4, and P5. The phosphate group is linked to the ribose sugar at the 5' position and to the guanine base at the 1' position. The structure is drawn with standard chemical notation, including wedges and dashes to indicate stereochemistry.

- Molecule 4 is (3 {S})-5-oxidanyl-3-[2-[[6-[[3-[(1 {S})-6-oxidanyl-3-oxidanylidene-1,2-dihydroisoindol-1-yl]-1 {H}-indol-2-yl]methylamino]hexylamino]methyl]-1 {H}-indol-3-yl]-2,3-dihydroisoindol-1-one (three-letter code: R6W) (formula: C<sub>40</sub>H<sub>40</sub>N<sub>6</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			50	40	6	4		
4	B	1	Total	C	N	O	0	0
			50	40	6	4		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	148	Total 148	O 148	0	0
6	B	148	Total 148	O 148	0	0
6	C	114	Total 114	O 114	0	0
6	D	105	Total 105	O 105	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: GTPase KRas




- Molecule 1: GTPase KRas



- Molecule 1: GTPase KRas



- Molecule 1: GTPase KRas



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.26Å 67.19Å 71.44Å 90.00° 96.47° 90.00°	Depositor
Resolution (Å)	22.99 – 1.57 22.94 – 1.57	Depositor EDS
% Data completeness (in resolution range)	97.8 (22.99-1.57) 97.9 (22.94-1.57)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 1.57Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, $R_{free}$	0.189 , 0.205 0.194 , 0.211	Depositor DCC
$R_{free}$ test set	1858 reflections (2.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 38.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6054	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: R6W, EDO, GCP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/1357	0.58	0/1831
1	B	0.41	0/1357	0.59	0/1831
1	C	0.41	0/1314	0.62	0/1771
1	D	0.39	0/1357	0.58	0/1831
All	All	0.41	0/5385	0.59	0/7264

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1336	0	1314	1	0
1	B	1336	0	1314	3	0
1	C	1295	0	1282	8	0
1	D	1336	0	1314	4	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	32	0	14	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	32	0	14	0	0
3	C	32	0	14	0	0
3	D	32	0	14	0	0
4	A	50	0	0	0	0
4	B	50	0	0	0	0
5	C	4	0	6	0	0
6	A	148	0	0	0	0
6	B	148	0	0	0	0
6	C	114	0	0	1	0
6	D	105	0	0	0	0
All	All	6054	0	5286	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:118:CYS:HB2	6:C:357:HOH:O	1.78	0.83
1:C:44:VAL:HG21	1:C:157:TYR:OH	2.02	0.60
1:B:24:ILE:HD13	1:B:42:LYS:HB2	1.85	0.57
1:D:44:VAL:HG21	1:D:157:TYR:OH	2.04	0.56
1:B:65:SER:HA	1:B:68:ARG:HD3	1.89	0.55
1:C:118:CYS:SG	1:C:143:GLU:HB3	2.50	0.52
1:A:68:ARG:HG2	1:A:71:TYR:OH	2.10	0.51
1:B:97:ARG:HG2	1:B:137:TYR:CZ	2.46	0.50
1:C:24:ILE:HD13	1:C:42:LYS:HB2	1.93	0.50
1:D:46:ILE:HG22	1:D:164:ARG:HH22	1.76	0.48
1:D:21:ILE:HD13	1:D:29:VAL:HG21	1.95	0.48
1:C:99:GLN:HG3	1:C:102:ARG:HH21	1.79	0.46
1:C:73:ARG:HG3	1:C:103:VAL:CG1	2.48	0.44
1:C:44:VAL:HG21	1:C:157:TYR:CZ	2.52	0.44
1:D:44:VAL:HG21	1:D:157:TYR:CZ	2.53	0.44
1:C:33:ASP:HA	1:C:34:PRO:HD2	1.95	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/170 (97%)	160 (97%)	5 (3%)	0	100	100
1	B	165/170 (97%)	160 (97%)	5 (3%)	0	100	100
1	C	158/170 (93%)	152 (96%)	6 (4%)	0	100	100
1	D	165/170 (97%)	160 (97%)	5 (3%)	0	100	100
All	All	653/680 (96%)	632 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	148/150 (99%)	145 (98%)	3 (2%)	55	29
1	B	148/150 (99%)	143 (97%)	5 (3%)	37	12
1	C	144/150 (96%)	142 (99%)	2 (1%)	67	45
1	D	148/150 (99%)	145 (98%)	3 (2%)	55	29
All	All	588/600 (98%)	575 (98%)	13 (2%)	52	25

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	TYR
1	A	41	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	63	GLU
1	B	32	TYR
1	B	41	ARG
1	B	62	GLU
1	B	71	TYR
1	B	142	ILE
1	C	41	ARG
1	C	67	MET
1	D	61	GLN
1	D	126	ASP
1	D	142	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	150	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GCP	A	202	2	27,34,34	1.26	4 (14%)	35,54,54	1.88	6 (17%)
4	R6W	A	203	-	49,57,57	0.87	2 (4%)	64,82,82	0.88	4 (6%)
3	GCP	D	202	2	27,34,34	1.24	4 (14%)	35,54,54	1.82	7 (20%)
5	EDO	C	203	-	3,3,3	0.50	0	2,2,2	0.42	0
4	R6W	B	203	-	49,57,57	0.90	2 (4%)	64,82,82	0.93	5 (7%)
3	GCP	B	202	2	27,34,34	1.30	3 (11%)	35,54,54	1.94	6 (17%)
3	GCP	C	202	2	27,34,34	1.33	3 (11%)	35,54,54	1.90	7 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GCP	A	202	2	-	0/15/38/38	0/3/3/3
4	R6W	A	203	-	-	0/9/45/45	0/8/8/8
3	GCP	D	202	2	-	0/15/38/38	0/3/3/3
5	EDO	C	203	-	-	0/1/1/1	-
4	R6W	B	203	-	-	0/9/45/45	0/8/8/8
3	GCP	B	202	2	-	0/15/38/38	0/3/3/3
3	GCP	C	202	2	-	0/15/38/38	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	202	GCP	C5-C6	4.17	1.48	1.41
3	D	202	GCP	C5-C6	3.72	1.47	1.41
3	A	202	GCP	C5-C6	3.70	1.47	1.41
3	B	202	GCP	C5-C6	3.69	1.47	1.41
4	B	203	R6W	C7-N8	3.39	1.49	1.46
4	A	203	R6W	C7-N8	3.32	1.49	1.46
3	B	202	GCP	PG-O3G	3.05	1.61	1.55
3	C	202	GCP	PB-O3A	3.04	1.61	1.58
4	B	203	R6W	C35-N36	2.84	1.49	1.46
4	A	203	R6W	C35-N36	2.79	1.49	1.46
3	A	202	GCP	PB-O3A	2.66	1.61	1.58
3	A	202	GCP	PG-O3G	2.34	1.60	1.55
3	D	202	GCP	PG-O3G	2.33	1.60	1.55
3	B	202	GCP	PB-O3A	2.30	1.61	1.58
3	D	202	GCP	PG-O2G	2.23	1.59	1.55
3	A	202	GCP	PG-O2G	2.18	1.59	1.55

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	202	GCP	PB-O3A	2.15	1.60	1.58
3	C	202	GCP	PG-O3G	2.04	1.59	1.55

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	202	GCP	C2-N1-C6	5.71	123.90	115.96
3	A	202	GCP	C2-N1-C6	5.63	123.79	115.96
3	C	202	GCP	C2-N1-C6	5.52	123.64	115.96
3	D	202	GCP	C2-N1-C6	5.08	123.02	115.96
3	C	202	GCP	C5-C6-N1	-4.76	117.06	123.42
3	B	202	GCP	C5-C6-N1	-4.41	117.52	123.42
3	B	202	GCP	C2-N3-C4	4.32	120.14	115.48
3	A	202	GCP	C5-C6-N1	-4.24	117.75	123.42
3	D	202	GCP	C5-C6-N1	-4.23	117.77	123.42
3	D	202	GCP	C2-N3-C4	4.16	119.97	115.48
3	B	202	GCP	N3-C2-N1	-4.15	121.93	127.21
3	A	202	GCP	C2-N3-C4	4.13	119.94	115.48
3	A	202	GCP	N3-C2-N1	-4.11	121.98	127.21
3	D	202	GCP	N3-C2-N1	-3.83	122.35	127.21
3	C	202	GCP	C2-N3-C4	3.78	119.56	115.48
3	C	202	GCP	N3-C2-N1	-3.72	122.48	127.21
3	A	202	GCP	C4-C5-C6	-3.30	116.19	121.23
3	B	202	GCP	C4-C5-C6	-3.05	116.57	121.23
3	D	202	GCP	C4-C5-C6	-2.93	116.75	121.23
3	C	202	GCP	C4-C5-C6	-2.92	116.77	121.23
4	B	203	R6W	C33-C34-C37	2.85	109.40	108.55
4	B	203	R6W	C5-C6-C9	2.74	109.37	108.55
4	A	203	R6W	C33-C34-C37	2.72	109.36	108.55
4	B	203	R6W	C11-C7-N8	-2.58	108.78	113.32
3	C	202	GCP	C1'-N9-C4	-2.53	122.20	126.64
4	A	203	R6W	C5-C6-C9	2.44	109.28	108.55
3	D	202	GCP	C1'-N9-C4	-2.42	122.39	126.64
4	B	203	R6W	C48-C40-N41	-2.39	119.81	124.27
3	B	202	GCP	PB-O3A-PA	-2.18	125.27	132.37
3	A	202	GCP	PB-O3A-PA	-2.13	125.42	132.37
3	D	202	GCP	PB-O3A-PA	-2.12	125.47	132.37
4	B	203	R6W	C12-C20-N21	-2.06	109.44	113.29
4	A	203	R6W	C20-C12-N13	-2.04	120.46	124.27
3	C	202	GCP	C4-C5-N7	-2.03	107.19	109.34
4	A	203	R6W	C48-C40-N41	-2.00	120.53	124.27


There are no chirality outliers.




There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	167/170 (98%)	0.27	5 (2%) 50 51	14, 22, 40, 57	0
1	B	167/170 (98%)	0.29	9 (5%) 25 26	13, 21, 38, 70	0
1	C	162/170 (95%)	0.82	21 (12%) 3 3	14, 27, 52, 74	0
1	D	167/170 (98%)	0.66	17 (10%) 6 6	14, 28, 55, 70	0
All	All	663/680 (97%)	0.51	52 (7%) 13 13	13, 24, 49, 74	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	122	SER	8.2
1	C	1	MET	6.0
1	C	79	LEU	4.2
1	A	108	ASP	4.2
1	C	67	MET	3.8
1	C	105	ASP	3.6
1	D	48	GLY	3.6
1	D	135	ARG	3.4
1	C	121	PRO	3.4
1	D	128	LYS	3.3
1	C	166	HIS	3.2
1	C	95	HIS	3.2
1	D	157	TYR	3.2
1	B	167	LYS	3.1
1	C	165	LYS	3.1
1	A	166	HIS	3.0
1	C	45	VAL	2.9
1	B	32	TYR	2.9
1	C	47	ASP	2.9
1	C	48	GLY	2.8
1	C	50	THR	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	118	CYS	2.8
1	D	142	ILE	2.8
1	D	64	TYR	2.7
1	D	133	LEU	2.7
1	C	9	VAL	2.7
1	C	99	GLN	2.6
1	C	44	VAL	2.6
1	B	1	MET	2.6
1	C	112	VAL	2.5
1	D	108	ASP	2.4
1	C	113	LEU	2.4
1	B	105	ASP	2.4
1	C	163	ILE	2.4
1	D	126	ASP	2.3
1	D	166	HIS	2.3
1	A	137	TYR	2.3
1	A	63	GLU	2.2
1	A	114	VAL	2.2
1	B	81	VAL	2.2
1	D	8	VAL	2.2
1	D	1	MET	2.2
1	D	114	VAL	2.1
1	D	95	HIS	2.1
1	D	91	GLU	2.1
1	C	81	VAL	2.1
1	D	47	ASP	2.1
1	C	120	LEU	2.1
1	B	62	GLU	2.1
1	B	65	SER	2.0
1	B	108	ASP	2.0
1	B	66	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	C	203	4/4	0.76	0.19	36,37,37,38	0
4	R6W	A	203	50/50	0.95	0.07	14,17,21,22	0
4	R6W	B	203	50/50	0.97	0.07	13,15,18,21	0
3	GCP	B	202	32/32	0.98	0.06	14,17,21,23	0
3	GCP	C	202	32/32	0.98	0.07	13,16,19,22	0
3	GCP	D	202	32/32	0.98	0.07	15,18,20,20	0
2	MG	B	201	1/1	0.99	0.04	15,15,15,15	0
2	MG	C	201	1/1	0.99	0.05	15,15,15,15	0
2	MG	D	201	1/1	0.99	0.07	16,16,16,16	0
3	GCP	A	202	32/32	0.99	0.06	14,16,19,20	0
2	MG	A	201	1/1	0.99	0.04	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around R6W B 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around GCP B 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around GCP C 202:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GCP D 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.