



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 02:22 PM EDT

PDB ID : 5AEA
Title : Crystal structure of human NCAM domain 1
Authors : Kvensakul, M.; Griffiths, K.; Foley, M.
Deposited on : 2015-08-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	118	<div> <div>4%</div> <div> <div></div> <div>71%</div> <div>14%</div> <div>15%</div> </div> </div>
1	B	118	<div> <div>7%</div> <div> <div></div> <div>73%</div> <div>9%</div> <div>17%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FLC	B	1100	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3128 atoms, of which 1444 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEURAL CELL ADHESION MOLECULE 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	100	Total	C	H	N	O	S	0	1	0
			1500	480	734	123	161	2			
1	B	98	Total	C	H	N	O	S	0	0	0
			1447	463	705	121	156	2			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	98	ALA	-	expression tag	UNP P13591
A	99	ALA	-	expression tag	UNP P13591
A	100	ALA	-	expression tag	UNP P13591
A	101	ASP	-	expression tag	UNP P13591
A	102	TYR	-	expression tag	UNP P13591
A	103	LYS	-	expression tag	UNP P13591
A	104	ASP	-	expression tag	UNP P13591
A	105	ASP	-	expression tag	UNP P13591
A	106	ASP	-	expression tag	UNP P13591
A	107	ASP	-	expression tag	UNP P13591
A	108	LYS	-	expression tag	UNP P13591
A	109	ALA	-	expression tag	UNP P13591
A	110	ALA	-	expression tag	UNP P13591
A	111	ASP	-	expression tag	UNP P13591
A	112	TYR	-	expression tag	UNP P13591
A	113	LYS	-	expression tag	UNP P13591
A	114	ASP	-	expression tag	UNP P13591
A	115	ASP	-	expression tag	UNP P13591
A	116	ASP	-	expression tag	UNP P13591
A	117	ASP	-	expression tag	UNP P13591
A	118	LYS	-	expression tag	UNP P13591
B	98	ALA	-	expression tag	UNP P13591
B	99	ALA	-	expression tag	UNP P13591
B	100	ALA	-	expression tag	UNP P13591
B	101	ASP	-	expression tag	UNP P13591

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	102	TYR	-	expression tag	UNP P13591
B	103	LYS	-	expression tag	UNP P13591
B	104	ASP	-	expression tag	UNP P13591
B	105	ASP	-	expression tag	UNP P13591
B	106	ASP	-	expression tag	UNP P13591
B	107	ASP	-	expression tag	UNP P13591
B	108	LYS	-	expression tag	UNP P13591
B	109	ALA	-	expression tag	UNP P13591
B	110	ALA	-	expression tag	UNP P13591
B	111	ASP	-	expression tag	UNP P13591
B	112	TYR	-	expression tag	UNP P13591
B	113	LYS	-	expression tag	UNP P13591
B	114	ASP	-	expression tag	UNP P13591
B	115	ASP	-	expression tag	UNP P13591
B	116	ASP	-	expression tag	UNP P13591
B	117	ASP	-	expression tag	UNP P13591
B	118	LYS	-	expression tag	UNP P13591

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	H	O	0	0
			18	6	5	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	86	Total 86	O 86	0	0
3	B	77	Total 77	O 77	0	0

• Molecule 1: NEURAL CELL ADHESION MOLECULE 1



- Chain B:  7% 73% 9% 17%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	23.85Å 107.33Å 41.75Å 90.00° 99.57° 90.00°	Depositor
Resolution (Å)	38.44 – 1.90 38.44 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.44-1.90) 100.0 (38.44-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 1.89Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.172 , 0.199 0.174 , 0.199	Depositor DCC
R_{free} test set	766 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.715	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 53.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3128	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/782	0.53	0/1063
1	B	0.44	0/753	0.54	0/1022
All	All	0.44	0/1535	0.53	0/2085

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	766	734	734	10	0
1	B	742	705	705	7	1
2	B	13	5	5	2	0
3	A	86	0	0	5	0
3	B	77	0	0	4	1
All	All	1684	1444	1444	18	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:ASN:OD1	3:A:2083:HOH:O	1.93	0.85
1:B:36:PHE:O	3:B:2034:HOH:O	1.97	0.81
1:A:82:GLU:O	3:A:2079:HOH:O	2.12	0.67
1:B:30:ASP:OD2	3:B:2029:HOH:O	2.12	0.66
1:A:56:ASP:O	3:A:2062:HOH:O	2.17	0.60
1:A:50:ILE:O	3:A:2048:HOH:O	2.16	0.59
1:B:19:PHE:CE2	2:B:1100:FLC:OB2	2.62	0.53
1:B:85:SER:OG	1:B:86:GLU:N	2.42	0.53
1:B:76:LYS:NZ	3:B:2070:HOH:O	2.42	0.52
1:A:45:PRO:O	3:A:2048:HOH:O	2.20	0.50
1:A:31:LYS:HG3	1:A:81:GLY:HA2	1.96	0.48
1:A:3:VAL:CG2	1:A:22:CYS:SG	3.02	0.46
2:B:1100:FLC:OG2	2:B:1100:FLC:CBC	2.64	0.44
1:A:37:SER:OG	1:A:39:ASN:OD1	2.35	0.42
1:A:49:ARG:HA	1:A:65[A]:TYR:CE2	2.55	0.42
1:B:75:TYR:HA	3:B:2034:HOH:O	2.20	0.41
1:A:37:SER:HB3	1:A:43:LEU:HD21	2.01	0.41
1:B:80:THR:HG22	1:B:86:GLU:HG2	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:GLN:NE2	3:B:2034:HOH:O[1_455]	1.80	0.40

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/118 (84%)	96 (97%)	3 (3%)	0	100	100
1	B	94/118 (80%)	91 (97%)	3 (3%)	0	100	100
All	All	193/236 (82%)	187 (97%)	6 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/101 (85%)	84 (98%)	2 (2%)	50	45
1	B	83/101 (82%)	79 (95%)	4 (5%)	25	16
All	All	169/202 (84%)	163 (96%)	6 (4%)	35	26

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	32	ASP
1	B	2	GLN
1	B	4	ASP
1	B	53	VAL
1	B	86	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FLC	B	1100	-	12,12,12	2.16	4 (33%)	17,17,17	1.50	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	B	1100	-	-	6/16/16/16	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1100	FLC	CA-CB	-3.78	1.49	1.53
2	B	1100	FLC	CG-CB	-3.52	1.49	1.53
2	B	1100	FLC	CB-CBC	-3.41	1.49	1.53
2	B	1100	FLC	OHB-CB	-2.80	1.37	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1100	FLC	OB2-CBC-CB	3.99	119.97	113.05

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1100	FLC	CB-CA-CAC-OA1
2	B	1100	FLC	CB-CA-CAC-OA2
2	B	1100	FLC	CA-CB-CBC-OB2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	1100	FLC	CG-CB-CBC-OB2
2	B	1100	FLC	CB-CG-CGC-OG1
2	B	1100	FLC	CB-CG-CGC-OG2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1100	FLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/118 (84%)	0.02	5 (5%) 28 32	16, 28, 68, 88	0
1	B	98/118 (83%)	0.27	8 (8%) 11 13	20, 34, 76, 122	0
All	All	198/236 (83%)	0.14	13 (6%) 18 20	16, 31, 75, 122	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	28	ALA	4.0
1	B	83	ASP	4.0
1	B	99	ALA	3.7
1	A	82	GLU	3.4
1	A	29	LYS	3.3
1	B	98	ALA	3.1
1	B	1	LEU	2.5
1	B	9	GLN	2.4
1	B	85	SER	2.3
1	A	27	ASP	2.3
1	B	27	ASP	2.2
1	B	29	LYS	2.2
1	A	25	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FLC	B	1100	13/13	0.76	0.44	110,131,161,165	0

6.5 Other polymers [i](#)

There are no such residues in this entry.