



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 25, 2024 – 03:50 PM EST

PDB ID : 2ATZ  
Title : Crystal structure of protein HP0184 from Helicobacter pylori  
Authors : Chang, C.; Xu, X.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2005-08-26  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	180	<div> <div>28%</div> <div>72%</div> <div>21%</div> <div>••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	202	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called H. pylori predicted coding region HP0184.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	176	1461	943	258	255	1	4	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP O24984
A	4	MSE	MET	modified residue	UNP O24984
A	44	MSE	MET	modified residue	UNP O24984
A	100	MSE	MET	modified residue	UNP O24984
A	136	MSE	MET	modified residue	UNP O24984

- Molecule 2 is 2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (three-letter code: DGT) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	100	Total	O	0	0
			100	100		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: *H. pylori* predicted coding region HP0184



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.09Å 77.09Å 118.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-2.00) 99.7 (50.00-2.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.75 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.198 , 0.256 0.200 , 0.185	Depositor DCC
$R_{free}$ test set	732 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 61.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	1596	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, DGT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.00	3/1495 (0.2%)	0.87	1/2004 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	178	LYS	C-O	12.61	1.47	1.23
1	A	33	CYS	CB-SG	-7.46	1.69	1.82
1	A	176	TRP	C-O	6.32	1.35	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	LEU	CA-CB-CG	6.63	130.54	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	22	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1461	0	1461	36	0
2	A	31	0	12	1	0
3	A	4	0	6	4	0
4	A	100	0	0	7	0
All	All	1596	0	1479	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:GLY:HA3	3:A:202:EDO:H21	1.37	1.05
1:A:22:GLY:HA3	1:A:23:LEU:HB2	1.41	1.01
1:A:22:GLY:CA	1:A:23:LEU:HB2	2.02	0.88
1:A:92:LEU:HD23	1:A:95:ARG:NH2	1.93	0.84
2:A:201:DGT:O2A	4:A:277:HOH:O	1.95	0.84
1:A:88:HIS:HA	1:A:91:GLN:OE1	1.78	0.82
1:A:89:LYS:HD3	1:A:142:LEU:HG	1.60	0.82
1:A:92:LEU:HD23	1:A:95:ARG:HH22	1.47	0.80
1:A:125:GLY:HA3	3:A:202:EDO:C2	2.12	0.78
1:A:91:GLN:O	1:A:95:ARG:HG3	1.87	0.75
1:A:106:ASN:HD22	1:A:166:GLU:HB2	1.50	0.74
1:A:125:GLY:CA	3:A:202:EDO:H21	2.17	0.74
1:A:53:HIS:HD2	1:A:60:ILE:H	1.37	0.71
1:A:77:ASP:OD1	4:A:277:HOH:O	2.09	0.70
1:A:177:ALA:O	1:A:178:LYS:HG2	1.96	0.66
1:A:53:HIS:CD2	1:A:60:ILE:H	2.15	0.63
1:A:91:GLN:HG3	4:A:280:HOH:O	1.98	0.62
1:A:44:MSE:HE2	1:A:46:THR:HG22	1.85	0.59
1:A:125:GLY:CA	3:A:202:EDO:C2	2.79	0.58
1:A:79:ASN:ND2	4:A:277:HOH:O	2.08	0.55
1:A:62:HIS:HE1	1:A:116:TYR:OH	1.91	0.54
1:A:92:LEU:CD2	1:A:95:ARG:HH22	2.19	0.54
1:A:101:ASN:HD22	1:A:172:ARG:HA	1.73	0.53

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:HIS:HD2	4:A:216:HOH:O	1.93	0.51
1:A:10:LYS:HD2	1:A:72:GLU:CD	2.31	0.51
1:A:22:GLY:CA	1:A:23:LEU:CB	2.86	0.49
1:A:135:SER:HB2	1:A:148:VAL:HG11	1.96	0.48
1:A:88:HIS:O	1:A:91:GLN:HG2	2.14	0.47
1:A:8:LEU:HD12	4:A:295:HOH:O	2.15	0.46
1:A:170:LYS:HE2	1:A:172:ARG:NH2	2.32	0.45
1:A:47:SER:O	1:A:51:GLN:HG2	2.17	0.45
1:A:72:GLU:HG2	1:A:121:HIS:CE1	2.53	0.44
1:A:8:LEU:HA	4:A:295:HOH:O	2.17	0.43
1:A:10:LYS:HD2	1:A:72:GLU:OE2	2.19	0.43
1:A:147:LYS:HB2	1:A:147:LYS:HE2	1.97	0.41
1:A:53:HIS:CE1	1:A:163:LEU:HB2	2.55	0.41
1:A:86:PHE:HA	1:A:142:LEU:HD11	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	174/180 (97%)	171 (98%)	1 (1%)	2 (1%)	<b>12</b> <b>7</b>

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	GLU
1	A	23	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	155/155 (100%)	143 (92%)	12 (8%)	10 7

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	33	CYS
1	A	47	SER
1	A	56	ARG
1	A	68	ASN
1	A	81	ARG
1	A	93	LEU
1	A	96	GLU
1	A	121	HIS
1	A	154	TRP
1	A	156	LYS
1	A	174	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	A	62	HIS
1	A	73	ASN
1	A	101	ASN
1	A	106	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	EDO	A	202	-	3,3,3	0.59	0	2,2,2	0.35	0
2	DGT	A	201	-	29,33,33	1.60	8 (27%)	37,52,52	1.43	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	202	-	-	0/1/1/1	-
2	DGT	A	201	-	-	5/18/34/34	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	DGT	PA-O3A	3.79	1.63	1.59
2	A	201	DGT	O6-C6	3.11	1.30	1.23
2	A	201	DGT	PB-O3B	2.84	1.62	1.59
2	A	201	DGT	C5-C4	2.48	1.49	1.43
2	A	201	DGT	PG-O2G	-2.38	1.45	1.54
2	A	201	DGT	C6-N1	-2.36	1.34	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	DGT	C2-N3	2.34	1.38	1.33
2	A	201	DGT	O4'-C4'	-2.00	1.40	1.45

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	DGT	O6-C6-N1	3.12	124.33	120.62
2	A	201	DGT	O2G-PG-O3G	2.99	122.47	110.83
2	A	201	DGT	O1A-PA-O2A	2.79	125.41	112.44
2	A	201	DGT	C8-N7-C5	2.73	107.19	102.55
2	A	201	DGT	O6-C6-C5	-2.64	119.09	124.32
2	A	201	DGT	O1A-PA-O3A	-2.44	100.67	107.27
2	A	201	DGT	O1G-PG-O3B	2.38	112.60	104.64

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	DGT	PB-O3B-PG-O1G
2	A	201	DGT	PA-O3A-PB-O2B
2	A	201	DGT	PB-O3B-PG-O3G
2	A	201	DGT	PB-O3B-PG-O2G
2	A	201	DGT	PA-O3A-PB-O1B


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	202	EDO	4	0
2	A	201	DGT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

**Warning:** The R factor obtained from EDS is 0.3299, which does not match the depositor's R factor of 0.19789. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	172/180 (95%)	1.69	51 (29%) <b>1</b> <b>1</b>	33, 42, 52, 81	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	TRP	5.1
1	A	23	LEU	4.4
1	A	178	LYS	4.1
1	A	30	ALA	4.1
1	A	6	LEU	4.0
1	A	64	LEU	3.7
1	A	34	PHE	3.4
1	A	31	GLY	3.4
1	A	32	ARG	3.4
1	A	7	LYS	3.4
1	A	8	LEU	3.2
1	A	9	ILE	3.2
1	A	175	SER	3.0
1	A	33	CYS	2.9
1	A	95	ARG	2.9
1	A	143	PRO	2.9
1	A	61	ALA	2.9
1	A	111	GLY	2.9
1	A	29	TYR	2.8
1	A	35	TYR	2.7
1	A	142	LEU	2.7
1	A	27	VAL	2.7
1	A	141	GLY	2.7
1	A	163	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	167	VAL	2.6
1	A	16	TYR	2.6
1	A	177	ALA	2.6
1	A	165	TYR	2.6
1	A	65	ILE	2.6
1	A	66	LEU	2.6
1	A	174	SER	2.5
1	A	99	PHE	2.5
1	A	92	LEU	2.5
1	A	63	ASN	2.5
1	A	160	ILE	2.4
1	A	22	GLY	2.4
1	A	91	GLN	2.4
1	A	162	ALA	2.3
1	A	43	ALA	2.3
1	A	69	ASP	2.2
1	A	67	ARG	2.2
1	A	60	ILE	2.1
1	A	5	GLU	2.1
1	A	173	GLY	2.1
1	A	71	VAL	2.1
1	A	124	LEU	2.1
1	A	82	ASN	2.1
1	A	70	LYS	2.1
1	A	125	GLY	2.0
1	A	116	TYR	2.0
1	A	41	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	202	4/4	0.67	0.31	118,119,119,120	0
2	DGT	A	201	31/31	0.93	0.10	21,26,28,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.