



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2024 – 10:23 AM EDT

PDB ID : 4AY6  
Title : Human O-GlcNAc transferase (OGT) in complex with UDP-5SGlcNAc and substrate peptide  
Authors : Schimpl, M.; Zheng, X.; Blair, D.E.; Schuettelkopf, A.W.; Navratilova, I.; Aristotelous, T.; Ferenbach, A.T.; Macnaughtan, M.A.; Borodkin, V.S.; van Aalten, D.M.F.  
Deposited on : 2012-06-18  
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	723	
1	B	723	
1	C	723	
1	D	723	
2	E	13	
2	F	13	
2	G	13	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	13	 A horizontal bar chart showing the quality of chain H. The bar is divided into four segments: green (54%), yellow (23%), orange (8%), and grey (15%).

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 22544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-N-ACETYLGLUCOSAMINE--PEPTIDE N-ACETYL GLUCOSAMINYLTRANS FERASE 110 KDA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	698	Total	C	N	O	S	0	0	0
			5514	3499	964	1013	38			
1	B	698	Total	C	N	O	S	0	0	0
			5514	3499	964	1013	38			
1	C	698	Total	C	N	O	S	0	0	0
			5514	3499	964	1013	38			
1	D	698	Total	C	N	O	S	0	0	0
			5514	3499	964	1013	38			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	309	GLY	-	expression tag	UNP O15294
A	310	PRO	-	expression tag	UNP O15294
A	311	GLY	-	expression tag	UNP O15294
A	312	SER	-	expression tag	UNP O15294
B	309	GLY	-	expression tag	UNP O15294
B	310	PRO	-	expression tag	UNP O15294
B	311	GLY	-	expression tag	UNP O15294
B	312	SER	-	expression tag	UNP O15294
C	309	GLY	-	expression tag	UNP O15294
C	310	PRO	-	expression tag	UNP O15294
C	311	GLY	-	expression tag	UNP O15294
C	312	SER	-	expression tag	UNP O15294
D	309	GLY	-	expression tag	UNP O15294
D	310	PRO	-	expression tag	UNP O15294
D	311	GLY	-	expression tag	UNP O15294
D	312	SER	-	expression tag	UNP O15294


- Molecule 2 is a protein called TGF-BETA-ACTIVATED KINASE 1 AND MAP3K7-BINDING PROTEIN 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	11	Total	C	N	O	0	0	0
			78	49	13	16			
2	F	11	Total	C	N	O	0	0	0
			78	49	13	16			
2	G	11	Total	C	N	O	0	0	0
			78	49	13	16			
2	H	11	Total	C	N	O	0	0	0
			78	49	13	16			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	1395	DNP	SER	SEE REMARK 999	UNP Q15750
F	1395	DNP	SER	SEE REMARK 999	UNP Q15750
G	1395	DNP	SER	SEE REMARK 999	UNP Q15750
H	1395	DNP	SER	SEE REMARK 999	UNP Q15750

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is (2S,3R,4R,5S,6R)-3-(acetylamino)-4,5-dihydroxy-6-(hydroxymethyl)tetrahydro-2H-thiopyran-2-yl [(2R,3S,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3,4-dihydroxytetrahydrofuran-2-yl]methyl dihydrogen diphosphate (three-letter code: 12V) (formula: C<sub>17</sub>H<sub>27</sub>N<sub>3</sub>O<sub>16</sub>P<sub>2</sub>S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			39	17	3	16	2	1		
4	B	1	Total	C	N	O	P	S	0	0
			39	17	3	16	2	1		
4	C	1	Total	C	N	O	P	S	0	0
			39	17	3	16	2	1		
4	D	1	Total	C	N	O	P	S	0	0
			39	17	3	16	2	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: UDP-N-ACETYLGLUCOSAMINE--PEPTIDE N-ACETYLGLUCOSAMINYLTR ANS FERASE 110 KDA SUBUNIT



- Molecule 1: UDP-N-ACETYLGLUCOSAMINE--PEPTIDE N-ACETYLGLUCOSAMINYLTR ANS FERASE 110 KDA SUBUNIT



- Molecule 1: UDP-N-ACETYLGLUCOSAMINE--PEPTIDE N-ACETYLGLUCOSAMINYLTR ANS FERASE 110 KDA SUBUNIT








- Molecule 1: UDP-N-ACETYLGLUCOSAMINE--PEPTIDE N-ACETYLGLUCOSAMINYLTR ANS FERASE 110 KDA SUBUNIT

Chain D: 81% 14%



- Molecule 2: TGF-BETA-ACTIVATED KINASE 1 AND MAP3K7-BINDING PROTEIN 1

Chain E: 46% 31% 8% 15%



- Molecule 2: TGF-BETA-ACTIVATED KINASE 1 AND MAP3K7-BINDING PROTEIN 1

Chain F: 46% 31% 8% 15%




- Molecule 2: TGF-BETA-ACTIVATED KINASE 1 AND MAP3K7-BINDING PROTEIN 1

Chain G: 46% 31% 8% 15%



- Molecule 2: TGF-BETA-ACTIVATED KINASE 1 AND MAP3K7-BINDING PROTEIN 1

Chain H:   
54% 23% 8% 15%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	275.44Å 275.44Å 142.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 3.30 25.01 – 3.30	Depositor EDS
% Data completeness (in resolution range)	98.2 (40.00-3.30) 98.5 (25.01-3.30)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.24 (at 3.30Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.227 , 0.272 0.226 , 0.271	Depositor DCC
$R_{free}$ test set	435 reflections (0.48%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.0	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , -2.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	0.237 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	22544	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DNP, 12V, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	3/5641 (0.1%)	0.66	1/7650 (0.0%)
1	B	0.61	2/5641 (0.0%)	0.67	0/7650
1	C	0.60	1/5641 (0.0%)	0.67	0/7650
1	D	0.57	2/5641 (0.0%)	0.63	0/7650
2	E	0.53	0/73	0.76	0/98
2	F	0.61	0/73	0.72	0/98
2	G	0.52	0/73	0.71	0/98
2	H	0.52	0/73	0.71	0/98
All	All	0.60	8/22856 (0.0%)	0.66	1/30992 (0.0%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	988	TRP	CD2-CE2	5.75	1.48	1.41
1	D	988	TRP	CD2-CE2	5.70	1.48	1.41
1	B	469	TRP	CD2-CE2	5.33	1.47	1.41
1	C	469	TRP	CD2-CE2	5.29	1.47	1.41
1	A	469	TRP	CD2-CE2	5.17	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	903	ARG	NE-CZ-NH1	5.10	122.85	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5514	0	5489	71	0
1	B	5514	0	5489	68	0
1	C	5514	0	5489	58	0
1	D	5514	0	5489	60	0
2	E	78	0	76	4	0
2	F	78	0	76	6	0
2	G	78	0	76	5	0
2	H	78	0	76	3	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
3	C	5	0	0	1	0
3	D	5	0	0	0	0
4	A	39	0	25	1	0
4	B	39	0	25	2	0
4	C	39	0	25	2	0
4	D	39	0	25	1	0
All	All	22544	0	22360	267	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 267 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:644:ARG:HG3	1:D:644:ARG:HH11	1.25	1.00
1:B:644:ARG:HH11	1:B:644:ARG:HG3	1.24	0.99
1:B:644:ARG:HH11	1:B:644:ARG:CG	1.77	0.98
1:D:644:ARG:HH11	1:D:644:ARG:CG	1.80	0.94
1:A:644:ARG:HG3	1:A:644:ARG:HH11	1.33	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	692/723 (96%)	648 (94%)	40 (6%)	4 (1%)	25	57
1	B	692/723 (96%)	641 (93%)	46 (7%)	5 (1%)	22	54
1	C	692/723 (96%)	643 (93%)	46 (7%)	3 (0%)	34	66
1	D	692/723 (96%)	643 (93%)	45 (6%)	4 (1%)	25	57
2	E	8/13 (62%)	6 (75%)	1 (12%)	1 (12%)	0	1
2	F	8/13 (62%)	6 (75%)	1 (12%)	1 (12%)	0	1
2	G	8/13 (62%)	6 (75%)	1 (12%)	1 (12%)	0	1
2	H	8/13 (62%)	6 (75%)	1 (12%)	1 (12%)	0	1
All	All	2800/2944 (95%)	2599 (93%)	181 (6%)	20 (1%)	22	54

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	368	GLN
1	B	368	GLN
1	C	368	GLN
1	D	368	GLN
2	E	1390	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	600/618 (97%)	574 (96%)	26 (4%)	29	59

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	600/618 (97%)	575 (96%)	25 (4%)	30	60
1	C	600/618 (97%)	576 (96%)	24 (4%)	31	61
1	D	600/618 (97%)	575 (96%)	25 (4%)	30	60
2	E	9/11 (82%)	9 (100%)	0	100	100
2	F	9/11 (82%)	9 (100%)	0	100	100
2	G	9/11 (82%)	9 (100%)	0	100	100
2	H	9/11 (82%)	9 (100%)	0	100	100
All	All	2436/2516 (97%)	2336 (96%)	100 (4%)	30	61

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	384	PHE
1	C	793	SER
1	D	1026	MET
1	C	434	ASN
1	C	669	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	390	ASN
2	G	1398	GLN
1	D	406	GLN
1	D	784	GLN
1	B	402	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DNP	H	1395	2	4,5,6	0.60	0	1,5,7	0.11	0
2	DNP	G	1395	2	4,5,6	0.58	0	1,5,7	0.19	0
2	DNP	F	1395	2	4,5,6	0.75	0	1,5,7	0.32	0
2	DNP	E	1395	2	4,5,6	0.82	0	1,5,7	0.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DNP	H	1395	2	-	1/2/4/6	-
2	DNP	G	1395	2	-	1/2/4/6	-
2	DNP	F	1395	2	-	1/2/4/6	-
2	DNP	E	1395	2	-	1/2/4/6	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1395	DNP	N-CA-CB-NG
2	F	1395	DNP	N-CA-CB-NG
2	G	1395	DNP	N-CA-CB-NG
2	H	1395	DNP	N-CA-CB-NG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

There are no monosaccharides in this entry.



## 5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	B	1100	-	4,4,4	0.37	0	6,6,6	0.26	0
3	SO4	C	1100	-	4,4,4	0.41	0	6,6,6	0.20	0
4	12V	B	1200	-	38,41,41	1.27	4 (10%)	52,62,62	1.57	11 (21%)
4	12V	C	1200	-	38,41,41	1.41	6 (15%)	52,62,62	1.71	11 (21%)
4	12V	A	1200	-	38,41,41	1.76	5 (13%)	52,62,62	1.73	9 (17%)
4	12V	D	1200	-	38,41,41	1.16	3 (7%)	52,62,62	1.71	10 (19%)
3	SO4	A	1100	-	4,4,4	0.43	0	6,6,6	0.19	0
3	SO4	D	1100	-	4,4,4	0.27	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	12V	C	1200	-	-	5/25/63/63	0/3/3/3
4	12V	B	1200	-	-	5/25/63/63	0/3/3/3
4	12V	A	1200	-	-	5/25/63/63	0/3/3/3
4	12V	D	1200	-	-	5/25/63/63	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1200	12V	PA-O3A	7.76	1.67	1.59
4	C	1200	12V	PA-O3A	4.40	1.64	1.59
4	D	1200	12V	C5'-S5'	-3.85	1.76	1.82
4	A	1200	12V	C5'-S5'	-3.76	1.76	1.82
4	C	1200	12V	C5'-S5'	-3.07	1.77	1.82

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1200	12V	C4-N3-C2	-6.07	119.08	126.61
4	D	1200	12V	C4-N3-C2	-5.95	119.23	126.61
4	A	1200	12V	C4-N3-C2	-5.38	119.94	126.61
4	D	1200	12V	N3-C2-N1	4.83	121.18	114.89
4	A	1200	12V	C5-C4-N3	4.80	121.53	114.80

There are no chirality outliers.

5 of 20 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
4	A	1200	12V	C3B-C4B-C5B-O5B
4	A	1200	12V	O4B-C4B-C5B-O5B
4	A	1200	12V	C4'-C5'-C6'-O6'
4	B	1200	12V	C3B-C4B-C5B-O5B
4	B	1200	12V	O4B-C4B-C5B-O5B


There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1100	SO4	1	0
4	B	1200	12V	2	0
4	C	1200	12V	2	0
4	A	1200	12V	1	0
4	D	1200	12V	1	0
3	A	1100	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.


### 6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands ⓘ




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around 12V B 1200:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 12V C 1200:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.