



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 31, 2025 – 12:13 PM EDT

PDB ID : 9C2Y / pdb_00009c2y
Title : Crystal Structure of JF1cpCasp2 in complex with MUR-65
Authors : Fuller, J.L.; Finzel, B.C.; Shi, K.
Deposited on : 2024-05-31
Resolution : 1.96 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	<div> <div>4%</div> <div>70% 13% 16%</div> </div>
1	B	282	<div> <div>4%</div> <div>79% 11% 11%</div> </div>
1	C	282	<div> <div>2%</div> <div>71% 13% 16%</div> </div>
1	D	282	<div> <div>2%</div> <div>74% 10% 16%</div> </div>
2	F	5	<div> <div>20%</div> <div>20% 40% 40%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	5	
2	H	5	
2	I	5	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	A1AT7	F	4	-	X	-	-
2	ASJ	G	5	-	-	X	-
2	A1AT6	H	1	-	X	-	-
2	A1AT7	H	4	-	X	-	-
2	A1AT7	I	4	-	X	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called JF1cpCasp2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	2	0
			1833	1152	326	337	18			
1	B	252	Total	C	N	O	S	0	2	0
			1952	1230	345	357	20			
1	C	238	Total	C	N	O	S	0	2	0
			1849	1161	327	342	19			
1	D	237	Total	C	N	O	S	0	3	0
			1840	1157	327	337	19			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P42575
A	2	HIS	-	expression tag	UNP P42575
A	3	HIS	-	expression tag	UNP P42575
A	4	HIS	-	expression tag	UNP P42575
A	5	HIS	-	expression tag	UNP P42575
A	6	HIS	-	expression tag	UNP P42575
A	7	HIS	-	expression tag	UNP P42575
A	21	ALA	ASP	conflict	UNP P42575
A	123	GLY	-	linker	UNP P42575
A	124	GLY	-	linker	UNP P42575
A	125	VAL	-	linker	UNP P42575
B	1	MET	-	initiating methionine	UNP P42575
B	2	HIS	-	expression tag	UNP P42575
B	3	HIS	-	expression tag	UNP P42575
B	4	HIS	-	expression tag	UNP P42575
B	5	HIS	-	expression tag	UNP P42575
B	6	HIS	-	expression tag	UNP P42575
B	7	HIS	-	expression tag	UNP P42575
B	21	ALA	ASP	conflict	UNP P42575
B	123	GLY	-	linker	UNP P42575
B	124	GLY	-	linker	UNP P42575

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	125	VAL	-	linker	UNP P42575
C	1	MET	-	initiating methionine	UNP P42575
C	2	HIS	-	expression tag	UNP P42575
C	3	HIS	-	expression tag	UNP P42575
C	4	HIS	-	expression tag	UNP P42575
C	5	HIS	-	expression tag	UNP P42575
C	6	HIS	-	expression tag	UNP P42575
C	7	HIS	-	expression tag	UNP P42575
C	21	ALA	ASP	conflict	UNP P42575
C	123	GLY	-	linker	UNP P42575
C	124	GLY	-	linker	UNP P42575
C	125	VAL	-	linker	UNP P42575
D	1	MET	-	initiating methionine	UNP P42575
D	2	HIS	-	expression tag	UNP P42575
D	3	HIS	-	expression tag	UNP P42575
D	4	HIS	-	expression tag	UNP P42575
D	5	HIS	-	expression tag	UNP P42575
D	6	HIS	-	expression tag	UNP P42575
D	7	HIS	-	expression tag	UNP P42575
D	21	ALA	ASP	conflict	UNP P42575
D	123	GLY	-	linker	UNP P42575
D	124	GLY	-	linker	UNP P42575
D	125	VAL	-	linker	UNP P42575

- Molecule 2 is a protein called MUR-65.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	F	5	Total C N O 49 35 5 9	0	0	0
2	G	5	Total C N O 49 35 5 9	0	0	0
2	H	5	Total C N O 49 35 5 9	0	0	0
2	I	5	Total C N O 49 35 5 9	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	7	Total O 7 7	0	0
3	B	10	Total O 10 10	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	6	Total	O	0	0
			6	6		
3	D	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: JF1cpCasp2



• Molecule 1: JF1cpCasp2



• Molecule 1: JF1cpCasp2






• Molecule 1: JF1cpCasp2




• Molecule 2: MUR-65



• Molecule 2: MUR-65



• Molecule 2: MUR-65



• Molecule 2: MUR-65



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.29Å 130.25Å 78.30Å 90.00° 104.56° 90.00°	Depositor
Resolution (Å)	34.04 – 1.96 34.04 – 1.96	Depositor EDS
% Data completeness (in resolution range)	46.6 (34.04-1.96) 43.3 (34.04-1.96)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.97Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.196 , 0.248 0.197 , 0.248	Depositor DCC
R_{free} test set	4345 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7701	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASJ, A1AT7, UN1, A1AT6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/1868	0.71	0/2525
1	B	0.50	0/1993	0.71	0/2695
1	C	0.48	0/1884	0.70	0/2545
1	D	0.49	0/1874	0.71	0/2532
2	F	0.86	0/6	2.18	0/7
2	G	1.06	0/6	2.31	0/7
2	H	0.90	0/6	2.98	1/7 (14.3%)
2	I	1.12	0/6	1.30	0/7
All	All	0.49	0/7643	0.72	1/10325 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	3	VAL	CB-CA-C	5.03	120.96	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1833	0	1762	30	0
1	B	1952	0	1881	27	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1849	0	1779	23	0
1	D	1840	0	1777	16	0
2	F	49	0	18	3	0
2	G	49	0	19	7	0
2	H	49	0	18	1	0
2	I	49	0	17	0	0
3	A	7	0	0	0	0
3	B	10	0	0	1	0
3	C	6	0	0	0	0
3	D	8	0	0	0	0
All	All	7701	0	7271	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:CYS:SG	2:G:5:ASJ:H	1.34	1.64
1:A:269:CYS:SG	2:F:5:ASJ:C	2.03	1.46
1:B:269:CYS:SG	2:G:5:ASJ:C	2.06	1.41
1:B:269:CYS:HG	2:G:5:ASJ:C	1.64	0.96
1:B:35:ARG:HH22	1:B:243:GLN:HE21	1.14	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/282 (83%)	224 (95%)	11 (5%)	0	100	100
1	B	252/282 (89%)	243 (96%)	9 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	236/282 (84%)	224 (95%)	12 (5%)	0	100	100
1	D	236/282 (84%)	228 (97%)	6 (2%)	2 (1%)	16	8
2	F	1/5 (20%)	1 (100%)	0	0	100	100
2	G	1/5 (20%)	1 (100%)	0	0	100	100
2	H	1/5 (20%)	1 (100%)	0	0	100	100
2	I	1/5 (20%)	1 (100%)	0	0	100	100
All	All	963/1148 (84%)	923 (96%)	38 (4%)	2 (0%)	44	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	164	GLU
1	D	162	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/238 (81%)	188 (98%)	4 (2%)	48	43
1	B	205/238 (86%)	204 (100%)	1 (0%)	86	86
1	C	195/238 (82%)	189 (97%)	6 (3%)	35	25
1	D	193/238 (81%)	186 (96%)	7 (4%)	30	20
2	F	1/1 (100%)	1 (100%)	0	100	100
2	G	1/1 (100%)	1 (100%)	0	100	100
2	H	1/1 (100%)	1 (100%)	0	100	100
2	I	1/1 (100%)	1 (100%)	0	100	100
All	All	789/956 (82%)	771 (98%)	18 (2%)	45	39

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	143	LEU
1	D	198	GLN
1	D	191	VAL
1	C	115	ARG
1	D	73	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	76	HIS
1	B	243	GLN
1	D	158	HIS
1	D	243	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

16 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	UN1	H	2	2	8,9,10	0.84	0	5,10,12	1.47	1 (20%)
2	A1AT7	H	4	2	14,14,15	7.56	9 (64%)	16,19,21	3.94	10 (62%)
2	A1AT6	I	1	2	11,12,13	6.81	6 (54%)	13,16,18	5.13	9 (69%)
2	UN1	F	2	2	8,9,10	0.89	0	5,10,12	1.56	1 (20%)
2	ASJ	G	5	2	7,7,7	1.05	0	5,8,8	1.53	1 (20%)
2	A1AT7	I	4	2	14,14,15	7.44	9 (64%)	16,19,21	4.09	10 (62%)
2	ASJ	F	5	2	7,7,7	1.00	0	5,8,8	1.45	1 (20%)
2	A1AT7	F	4	2	14,14,15	7.33	8 (57%)	16,19,21	4.05	11 (68%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UN1	G	2	2	8,9,10	0.93	0	5,10,12	1.60	0
2	A1AT6	G	1	2	11,12,13	7.00	6 (54%)	13,16,18	5.43	9 (69%)
2	A1AT7	G	4	2	14,14,15	7.61	8 (57%)	16,19,21	4.52	8 (50%)
2	A1AT6	F	1	2	11,12,13	7.03	6 (54%)	13,16,18	5.26	9 (69%)
2	ASJ	H	5	1,2	7,7,7	0.87	0	5,8,8	1.50	1 (20%)
2	A1AT6	H	1	2	11,12,13	7.03	6 (54%)	13,16,18	5.00	10 (76%)
2	ASJ	I	5	1,2	7,7,7	0.99	0	5,8,8	1.70	1 (20%)
2	UN1	I	2	2	8,9,10	0.99	0	5,10,12	1.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UN1	H	2	2	-	2/7/8/10	-
2	A1AT7	H	4	2	-	0/0/12/14	0/2/2/2
2	A1AT6	I	1	2	-	0/0/10/12	0/2/2/2
2	UN1	F	2	2	-	3/7/8/10	-
2	ASJ	G	5	2	-	6/6/6/6	-
2	A1AT7	I	4	2	-	0/0/12/14	0/2/2/2
2	ASJ	F	5	2	-	2/6/6/6	-
2	A1AT7	F	4	2	-	0/0/12/14	0/2/2/2
2	UN1	G	2	2	-	0/7/8/10	-
2	A1AT6	G	1	2	-	0/0/10/12	0/2/2/2
2	A1AT7	G	4	2	-	0/0/12/14	0/2/2/2
2	A1AT6	F	1	2	-	0/0/10/12	0/2/2/2
2	ASJ	H	5	1,2	-	3/6/6/6	-
2	A1AT6	H	1	2	-	0/0/10/12	0/2/2/2
2	ASJ	I	5	1,2	-	4/6/6/6	-
2	UN1	I	2	2	-	5/7/8/10	-

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	4	A1AT7	C27-CB	22.92	1.67	1.39
2	H	4	A1AT7	C27-CB	22.63	1.67	1.39
2	I	4	A1AT7	C27-CB	22.52	1.67	1.39
2	F	4	A1AT7	C27-CB	21.82	1.66	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	1	A1AT6	C20-C19	17.83	1.68	1.39

The worst 5 of 82 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	A1AT6	C23-C18-C19	-14.26	112.48	119.67
2	F	1	A1AT6	C23-C18-C19	-13.49	112.87	119.67
2	I	1	A1AT6	C23-C18-C19	-12.26	113.49	119.67
2	G	4	A1AT7	C20-C21-C22	-11.53	106.61	120.96
2	H	1	A1AT6	C23-C18-C19	-10.77	114.24	119.67

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	2	UN1	C6-C5-CA-C
2	I	2	UN1	C6-C5-CA-N
2	F	5	ASJ	C-CA-CB-CG
2	F	5	ASJ	N-CA-CB-CG
2	G	5	ASJ	O-C-CA-N

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	5	ASJ	6	0
2	F	5	ASJ	2	0
2	G	1	A1AT6	1	0
2	F	1	A1AT6	1	0
2	H	1	A1AT6	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/282 (84%)	0.26	11 (4%) 38 44	10, 32, 52, 68	2 (0%)
1	B	252/282 (89%)	0.07	11 (4%) 39 46	11, 29, 54, 81	2 (0%)
1	C	238/282 (84%)	0.21	7 (2%) 54 60	12, 32, 54, 71	2 (0%)
1	D	237/282 (84%)	0.05	7 (2%) 52 59	10, 29, 48, 77	3 (1%)
2	F	1/5 (20%)	2.04	1 (100%) 0 0	61, 61, 61, 61	0
2	G	1/5 (20%)	2.52	1 (100%) 0 0	55, 55, 55, 55	0
2	H	1/5 (20%)	2.64	1 (100%) 0 0	58, 58, 58, 58	0
2	I	1/5 (20%)	1.66	0 100 100	48, 48, 48, 48	0
All	All	968/1148 (84%)	0.15	39 (4%) 43 49	10, 30, 53, 81	9 (0%)

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	94	TYR	4.5
1	B	176[A]	SER	3.8
1	D	162	GLU	3.6
1	A	97	GLY	3.2
1	A	100	PHE	3.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A1AT6	H	1	11/12	0.62	0.22	67,71,80,83	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	A1AT6	G	1	11/12	0.76	0.17	54,58,65,66	0
2	A1AT6	F	1	11/12	0.78	0.21	56,66,74,74	0
2	A1AT7	F	4	13/14	0.80	0.16	48,56,63,65	0
2	ASJ	I	5	8/8	0.83	0.13	41,48,52,58	0
2	A1AT7	G	4	13/14	0.85	0.13	50,56,63,64	0
2	A1AT7	H	4	13/14	0.86	0.13	49,55,63,65	0
2	ASJ	H	5	8/8	0.87	0.14	51,57,64,67	0
2	A1AT6	I	1	11/12	0.87	0.12	37,43,51,52	0
2	ASJ	F	5	8/8	0.88	0.14	56,58,64,67	0
2	UN1	G	2	10/11	0.88	0.12	50,58,60,60	0
2	A1AT7	I	4	13/14	0.88	0.12	44,48,53,56	0
2	UN1	F	2	10/11	0.91	0.10	44,50,66,73	0
2	ASJ	G	5	8/8	0.91	0.11	46,50,56,59	0
2	UN1	H	2	10/11	0.92	0.12	42,55,57,65	0
2	UN1	I	2	10/11	0.93	0.08	29,37,44,49	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.