



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 18, 2025 – 06:13 PM EDT

PDB ID : 9C5O / pdb_00009c5o
Title : Trypanosoma cruzi R42F mutant beta-3-HBDH structure
Authors : Hashimoto, H.; Debler, E.W.
Deposited on : 2024-06-06
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div></div> </div> </div>
1	B	270	<div> <div>7%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div></div> </div> </div>
1	C	270	<div> <div>6%</div> <div> <div></div> <div>84%</div> <div>7%</div> <div>9%</div> </div> </div>
1	D	270	<div> <div>3%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div></div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16119 atoms, of which 7812 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydroxybutyrate dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	263	Total	C	H	N	O	S	0	2	0
			3936	1234	1982	340	366	14			
1	B	260	Total	C	H	N	O	S	0	4	0
			3927	1227	1986	338	364	12			
1	C	246	Total	C	H	N	O	S	0	2	0
			3689	1157	1864	317	339	12			
1	D	262	Total	C	H	N	O	S	0	1	0
			3902	1224	1962	338	365	13			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP A0A2V2VPF1
A	-2	PRO	-	expression tag	UNP A0A2V2VPF1
A	-1	HIS	-	expression tag	UNP A0A2V2VPF1
A	0	MET	-	expression tag	UNP A0A2V2VPF1
A	42	PHE	ARG	engineered mutation	UNP A0A2V2VPF1
B	-3	GLY	-	expression tag	UNP A0A2V2VPF1
B	-2	PRO	-	expression tag	UNP A0A2V2VPF1
B	-1	HIS	-	expression tag	UNP A0A2V2VPF1
B	0	MET	-	expression tag	UNP A0A2V2VPF1
B	42	PHE	ARG	engineered mutation	UNP A0A2V2VPF1
C	-3	GLY	-	expression tag	UNP A0A2V2VPF1
C	-2	PRO	-	expression tag	UNP A0A2V2VPF1
C	-1	HIS	-	expression tag	UNP A0A2V2VPF1
C	0	MET	-	expression tag	UNP A0A2V2VPF1
C	42	PHE	ARG	engineered mutation	UNP A0A2V2VPF1
D	-3	GLY	-	expression tag	UNP A0A2V2VPF1
D	-2	PRO	-	expression tag	UNP A0A2V2VPF1
D	-1	HIS	-	expression tag	UNP A0A2V2VPF1
D	0	MET	-	expression tag	UNP A0A2V2VPF1
D	42	PHE	ARG	engineered mutation	UNP A0A2V2VPF1

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	183	Total	O	0	0
			183	183		
3	B	161	Total	O	0	0
			161	161		
3	C	166	Total	O	0	0
			166	166		
3	D	125	Total	O	0	0
			125	125		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Hydroxybutyrate dehydrogenase




- Molecule 1: Hydroxybutyrate dehydrogenase



- Molecule 1: Hydroxybutyrate dehydrogenase



- Molecule 1: Hydroxybutyrate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.29Å 80.90Å 121.74Å 90.00° 107.50° 90.00°	Depositor
Resolution (Å)	38.70 – 1.69 38.70 – 1.69	Depositor EDS
% Data completeness (in resolution range)	86.0 (38.70-1.69) 86.0 (38.70-1.69)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 1.69Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.160 , 0.180 0.158 , 0.178	Depositor DCC
R_{free} test set	2000 reflections (1.63%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.356	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	16119	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	1/1990 (0.1%)	0.62	0/2696
1	B	0.42	0/1981	0.57	0/2682
1	C	0.43	0/1857	0.59	0/2517
1	D	0.35	0/1973	0.56	0/2675
All	All	0.44	1/7801 (0.0%)	0.58	0/10570

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	THR	C-O	-12.89	1.12	1.24

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	19	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1954	1982	1986	17	0
1	B	1941	1986	1989	25	0
1	C	1825	1864	1867	18	0
1	D	1940	1962	1964	16	0
2	B	12	18	18	0	0
3	A	183	0	0	5	0
3	B	161	0	0	14	0
3	C	166	0	0	7	0
3	D	125	0	0	5	0
All	All	8307	7812	7824	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:28:MET:HE3	1:C:58:TYR:HE2	1.20	1.02
1:C:5:LYS:N	3:C:301:HOH:O	1.91	1.01
1:B:220:ALA:O	3:B:501:HOH:O	1.85	0.93
1:A:201:GLN:OE1	3:A:301:HOH:O	1.89	0.90
1:C:28:MET:SD	1:C:55:VAL:HG22	2.13	0.88
1:C:28:MET:CE	1:C:58:TYR:HE2	1.87	0.86
1:C:5:LYS:CB	3:C:301:HOH:O	2.23	0.86
1:A:54:LYS:NZ	3:A:303:HOH:O	2.10	0.83
1:C:28:MET:HE3	1:C:58:TYR:CE2	2.12	0.80
1:B:217:ALA:O	3:B:502:HOH:O	2.01	0.79
1:B:136:GLN:NE2	3:B:507:HOH:O	2.23	0.71
1:B:220:ALA:CB	3:B:504:HOH:O	2.39	0.69
1:B:202:ILE:HD11	1:B:231:PHE:CZ	2.27	0.69
1:B:216:ALA:O	3:B:504:HOH:O	2.11	0.69
1:B:220:ALA:HB3	3:B:504:HOH:O	1.93	0.68
1:B:114:GLU:OE1	3:B:505:HOH:O	2.14	0.66
1:A:251:MET:HE1	1:B:228:ALA:HB3	1.77	0.65
1:B:84:LYS:HD2	3:B:622:HOH:O	1.96	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:28:MET:CE	1:C:58:TYR:CE2	2.77	0.63
1:B:54:LYS:NZ	3:B:503:HOH:O	2.03	0.62
1:C:72:ARG:HD2	3:C:420:HOH:O	2.02	0.60
1:A:32:MET:HE2	1:A:58:TYR:CZ	2.36	0.60
1:C:194:TYR:CB	3:C:302:HOH:O	2.50	0.59
1:A:43:ARG:N	3:A:302:HOH:O	2.02	0.58
1:D:229:LYS:NZ	3:D:303:HOH:O	2.37	0.57
1:B:197:LEU:O	1:B:201:GLN:NE2	2.37	0.57
1:C:127:THR:HG23	1:C:142[B]:ILE:HD13	1.86	0.56
1:A:32:MET:HE2	1:A:58:TYR:CE1	2.42	0.55
1:A:137:ARG:HD2	3:A:313:HOH:O	2.07	0.54
1:B:136:GLN:CD	3:B:507:HOH:O	2.48	0.53
1:D:32:MET:HE2	1:D:58:TYR:CZ	2.44	0.53
1:B:219:GLN:HA	1:B:219:GLN:OE1	2.10	0.52
1:A:251:MET:HE1	1:B:228:ALA:CB	2.40	0.52
1:C:16:THR:HG21	1:C:69:MET:HE2	1.91	0.51
1:D:98:ILE:HD11	1:D:118:LEU:HB2	1.93	0.51
1:C:197:LEU:O	1:C:201:GLN:HG3	2.11	0.51
1:D:229:LYS:CE	3:D:303:HOH:O	2.58	0.51
1:A:29:ARG:NH1	1:A:32:MET:HE1	2.26	0.50
1:B:54:LYS:CE	3:B:503:HOH:O	2.53	0.50
1:B:266:GLN:O	3:B:506:HOH:O	2.19	0.50
1:A:229:LYS:NZ	3:A:306:HOH:O	2.44	0.50
1:A:42:PHE:HE1	1:A:68:ASN:HB2	1.77	0.49
1:D:229:LYS:HE2	3:D:303:HOH:O	2.12	0.49
1:C:45:SER:N	1:C:46:PRO:CD	2.76	0.49
1:D:209:LYS:NZ	1:D:209:LYS:HB3	2.27	0.49
1:D:60:THR:OG1	3:D:301:HOH:O	2.20	0.48
1:D:29:ARG:NH1	1:D:32:MET:HE1	2.30	0.47
1:D:195:THR:HB	1:D:196:PRO:HD2	1.96	0.47
1:B:95:ASN:O	1:B:96:ALA:C	2.58	0.46
1:C:194:TYR:N	3:C:302:HOH:O	2.15	0.46
1:C:28:MET:SD	1:C:55:VAL:HA	2.56	0.46
1:D:15:VAL:HG12	1:D:18:SER:HB3	1.99	0.45
1:C:10:GLY:O	3:C:303:HOH:O	2.21	0.45
1:B:220:ALA:HB2	3:B:504:HOH:O	2.06	0.45
1:D:72:ARG:HD2	1:D:125:HIS:ND1	2.32	0.45
1:C:136:GLN:HG2	3:C:308:HOH:O	2.16	0.45
1:D:203:LYS:NZ	1:D:207:GLU:OE2	2.50	0.44
1:B:50:ALA:O	1:B:54:LYS:HG3	2.16	0.44
1:A:219:GLN:OE1	1:A:220:ALA:N	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:LYS:C	1:A:203:LYS:HE2	2.42	0.44
1:C:7:VAL:HG13	1:C:8:LEU:HG	2.00	0.43
1:A:127:THR:HG23	1:A:142:ILE:HD13	1.99	0.43
1:B:70:LYS:HG3	1:B:118[A]:LEU:HD21	2.00	0.43
1:A:7:VAL:HG23	1:A:8:LEU:HG	2.00	0.43
1:B:70:LYS:HE2	1:B:118[A]:LEU:HD11	2.01	0.43
1:D:125:HIS:HD2	3:D:322:HOH:O	2.01	0.43
1:D:134:MET:HB3	1:D:139:TRP:O	2.19	0.42
1:A:203:LYS:HE3	1:A:207:GLU:HG3	2.02	0.42
1:B:28:MET:O	1:B:32:MET:HG3	2.20	0.41
1:B:202:ILE:HD11	1:B:231:PHE:HZ	1.79	0.41
1:B:136:GLN:HG3	3:B:527:HOH:O	2.20	0.41
1:A:96:ALA:O	1:A:97:GLY:C	2.64	0.41
1:D:168:ILE:HG22	1:D:172:LYS:HE3	2.04	0.40
1:D:137:ARG:NH1	1:D:137:ARG:HG2	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/270 (97%)	257 (98%)	6 (2%)	0	100	100
1	B	260/270 (96%)	254 (98%)	6 (2%)	0	100	100
1	C	242/270 (90%)	236 (98%)	6 (2%)	0	100	100
1	D	261/270 (97%)	257 (98%)	4 (2%)	0	100	100
All	All	1026/1080 (95%)	1004 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/206 (98%)	201 (100%)	0	100	100
1	B	202/206 (98%)	200 (99%)	2 (1%)	73	64
1	C	189/206 (92%)	189 (100%)	0	100	100
1	D	199/206 (97%)	199 (100%)	0	100	100
All	All	791/824 (96%)	789 (100%)	2 (0%)	91	88

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	5	LYS
1	B	137	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	ASN
1	B	94	ASN
1	C	94	ASN
1	D	40	ASN
1	D	94	ASN
1	D	100	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	401	-	3,3,3	0.43	0	2,2,2	0.32	0
2	EDO	B	402	-	3,3,3	0.45	0	2,2,2	0.42	0
2	EDO	B	403	-	3,3,3	0.52	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	401	-	-	0/1/1/1	-
2	EDO	B	402	-	-	0/1/1/1	-
2	EDO	B	403	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	403	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/270 (97%)	-0.28	6 (2%) 61 64	19, 34, 63, 99	2 (0%)
1	B	260/270 (96%)	-0.10	18 (6%) 24 25	18, 34, 95, 121	4 (1%)
1	C	246/270 (91%)	-0.09	15 (6%) 28 29	15, 34, 83, 118	2 (0%)
1	D	262/270 (97%)	-0.00	7 (2%) 56 58	23, 42, 82, 124	1 (0%)
All	All	1031/1080 (95%)	-0.12	46 (4%) 39 41	15, 35, 83, 124	9 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	210	TYR	7.9
1	B	202	ILE	5.0
1	C	216	ALA	4.8
1	C	45	SER	4.7
1	B	205	VAL	4.4
1	D	194	TYR	4.3
1	C	197	LEU	4.2
1	C	198	VAL	4.0
1	B	217	ALA	4.0
1	C	202	ILE	3.9
1	A	4	GLU	3.9
1	B	204	ALA	3.8
1	B	197	LEU	3.7
1	A	97	GLY	3.6
1	D	46	PRO	3.4
1	A	96	ALA	3.2
1	C	194	TYR	3.2
1	B	206	ALA	3.1
1	C	217	ALA	2.9
1	B	196	PRO	2.9
1	C	46	PRO	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	216	ALA	2.8
1	B	218	THR	2.8
1	D	197	LEU	2.8
1	C	87	GLY	2.6
1	D	5	LYS	2.5
1	A	5	LYS	2.5
1	C	200	GLU	2.5
1	D	118	LEU	2.4
1	C	28	MET	2.4
1	C	48	ASP	2.3
1	C	41	GLY	2.3
1	B	98	ILE	2.2
1	A	219	GLN	2.2
1	B	198	VAL	2.2
1	B	54	LYS	2.2
1	D	84	LYS	2.2
1	B	3	GLY	2.2
1	B	215	GLU	2.2
1	A	196	PRO	2.1
1	B	203	LYS	2.1
1	D	42	PHE	2.1
1	C	201	GLN	2.1
1	B	208	ALA	2.0
1	B	209	LYS	2.0
1	C	5	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	B	403	4/4	0.78	0.15	52,62,70,73	0
2	EDO	B	401	4/4	0.90	0.12	49,67,73,88	0
2	EDO	B	402	4/4	0.93	0.09	39,52,67,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.