



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 03:18 PM EDT

PDB ID : 5CED
Title : Penicillin G Acylated Bd3459 Predatory Endopeptidase from *Bdellovibrio bacteriovorus* in complex with immunity protein Bd3460
Authors : Lovering, A.L.; Cadby, I.T.; Lambert, C.; Sockett, R.E.
Deposited on : 2015-07-06
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	<div> <div>11%</div> <div> <div></div> <div>80%</div> <div>••</div> <div>15%</div> </div> </div>
2	B	230	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>•</div> <div>16%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PNM	A	501	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4603 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bd3459.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	381	Total	C	N	O	S	0	1	0
			2960	1870	502	570	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	37	MET	ALA	conflict	UNP Q6MHT0
A	38	MET	LYS	engineered mutation	UNP Q6MHT0


- Molecule 2 is a protein called Bd3460.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	194	Total	C	N	O	S	0	0	0
			1442	899	251	288	4			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	25	MET	ALA	conflict	UNP Q6MHS9
B	221	LEU	-	expression tag	UNP Q6MHS9
B	222	GLU	-	expression tag	UNP Q6MHS9
B	223	HIS	-	expression tag	UNP Q6MHS9
B	224	HIS	-	expression tag	UNP Q6MHS9
B	225	HIS	-	expression tag	UNP Q6MHS9
B	226	HIS	-	expression tag	UNP Q6MHS9
B	227	HIS	-	expression tag	UNP Q6MHS9
B	228	HIS	-	expression tag	UNP Q6MHS9
B	229	HIS	-	expression tag	UNP Q6MHS9
B	230	HIS	-	expression tag	UNP Q6MHS9

- Molecule 3 is OPEN FORM - PENICILLIN G (three-letter code: PNM) (formula: C₁₆H₂₀N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			23	16	2	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	68	Total	O	0	0
			68	68		
4	B	110	Total	O	0	0
			110	110		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bd3459



• Molecule 2: Bd3460



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.54Å 59.16Å 192.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.02 – 2.02 32.21 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.8 (96.02-2.02) 99.9 (32.21-2.02)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.172 , 0.207 0.181 , 0.214	Depositor DCC
R_{free} test set	1977 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.385	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4603	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PNM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/3016	0.78	3/4068 (0.1%)
2	B	0.72	0/1450	0.82	4/1945 (0.2%)
All	All	0.68	0/4466	0.79	7/6013 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	389	ASP	CB-CG-OD1	6.51	124.16	118.30
2	B	110	ASP	CB-CG-OD1	6.50	124.15	118.30
2	B	55	ASP	CB-CG-OD2	-5.69	113.18	118.30
1	A	246	ARG	NE-CZ-NH1	5.61	123.10	120.30
2	B	189	ASP	CB-CG-OD1	5.57	123.31	118.30
1	A	118	ARG	NE-CZ-NH1	5.54	123.07	120.30
2	B	110	ASP	CB-CG-OD2	-5.23	113.59	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	70	SER	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2960	0	2975	10	0
2	B	1442	0	1502	1	0
3	A	23	0	17	2	0
4	A	68	0	0	0	0
4	B	110	0	0	0	0
All	All	4603	0	4494	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:501:PNM:N4	3:A:501:PNM:C3	1.78	1.18
1:A:298:ALA:O	1:A:300:PRO:HD3	1.95	0.66
3:A:501:PNM:N4	3:A:501:PNM:C11	2.61	0.60
1:A:189:GLY:O	1:A:193:ILE:HD12	2.07	0.54
1:A:241:LYS:HE2	1:A:326:TYR:OH	2.15	0.46
1:A:296:SER:OG	1:A:299:LYS:N	2.47	0.46
1:A:53:THR:HG23	1:A:55:GLN:H	1.81	0.44
1:A:115:LEU:HD13	1:A:255:ILE:HD11	1.99	0.43
1:A:50:ASN:C	1:A:50:ASN:HD22	2.22	0.42
1:A:77:THR:HG23	1:A:244:LEU:HD13	2.02	0.42
2:B:49:VAL:HG22	2:B:56:LEU:HD21	2.02	0.42
1:A:173:ILE:HD12	1:A:173:ILE:N	2.36	0.41
1:A:296:SER:HG	1:A:299:LYS:H	1.68	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	378/446 (85%)	361 (96%)	16 (4%)	1 (0%)	41	36
2	B	192/230 (84%)	190 (99%)	2 (1%)	0	100	100
All	All	570/676 (84%)	551 (97%)	18 (3%)	1 (0%)	47	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	298	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/380 (86%)	321 (99%)	4 (1%)	71	75
2	B	150/179 (84%)	150 (100%)	0	100	100
All	All	475/559 (85%)	471 (99%)	4 (1%)	81	85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	299	LYS
1	A	388	TYR
1	A	391	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	95	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	119	ASN
1	A	408	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PNM	A	501	1	19,24,24	3.93	6 (31%)	24,34,34	3.09	7 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PNM	A	501	1	1/1/6/9	1/12/33/33	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	PNM	C3-N4	14.63	1.78	1.46
3	A	501	PNM	C6-N14	-5.66	1.38	1.46
3	A	501	PNM	C5-S1	-4.45	1.74	1.84
3	A	501	PNM	C2-S1	-2.77	1.79	1.85
3	A	501	PNM	C15-N14	2.24	1.38	1.34
3	A	501	PNM	O16-C15	-2.01	1.19	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	PNM	C6-N14-C15	9.90	133.72	123.13
3	A	501	PNM	C5-C6-N14	7.61	126.58	109.98
3	A	501	PNM	O8-C7-C6	-5.11	110.59	124.83
3	A	501	PNM	C2-S1-C5	3.05	100.52	93.99
3	A	501	PNM	C17-C15-N14	-2.37	112.06	115.88
3	A	501	PNM	C9-C2-C10	-2.36	106.93	110.78
3	A	501	PNM	C17-C18-C19	-2.28	117.63	120.89

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	501	PNM	C6

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	PNM	C5-C6-N14-C15


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	PNM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	381/446 (85%)	0.67	51 (13%) 3 2	22, 45, 82, 105	0
2	B	194/230 (84%)	-0.05	7 (3%) 42 42	22, 34, 62, 112	0
All	All	575/676 (85%)	0.43	58 (10%) 7 6	22, 41, 82, 112	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	PHE	10.9
1	A	298	ALA	6.9
2	B	53	LYS	6.9
1	A	222	ALA	6.3
1	A	190	TYR	5.7
1	A	171	ALA	5.5
1	A	100	ALA	5.4
1	A	183	ALA	5.2
1	A	173	ILE	5.2
2	B	52	ASN	5.1
1	A	149	TRP	4.5
1	A	172	SER	4.4
1	A	184	THR	4.3
1	A	180	SER	4.2
1	A	101	ASN	4.2
1	A	187	SER	4.0
1	A	176	ALA	3.9
1	A	177	THR	3.8
1	A	170	GLN	3.7
1	A	53	THR	3.7
1	A	208	ARG	3.7
1	A	210	LYS	3.6
1	A	193	ILE	3.5
1	A	225	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	51	LYS	3.4
1	A	367	LEU	3.4
1	A	261	TRP	3.3
2	B	27	GLY	3.3
1	A	186	ILE	3.2
1	A	196	THR	3.2
1	A	299	LYS	3.2
1	A	178	LEU	3.2
1	A	420	LYS	3.1
1	A	179	THR	3.0
1	A	297	VAL	2.9
2	B	28	LYS	2.9
1	A	54	GLN	2.8
1	A	174	VAL	2.8
1	A	51	SER	2.7
1	A	74	VAL	2.7
1	A	229	LYS	2.6
2	B	47	ASN	2.6
1	A	356	ALA	2.5
1	A	228	GLU	2.5
1	A	189	GLY	2.5
1	A	52	LYS	2.5
1	A	425	LEU	2.5
1	A	71	ILE	2.4
1	A	200	ARG	2.3
1	A	296	SER	2.3
1	A	135	LYS	2.2
1	A	358	THR	2.2
1	A	214	ARG	2.2
1	A	357	LYS	2.1
1	A	368	MET	2.0
1	A	181	SER	2.0
2	B	31	LYS	2.0
1	A	102	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PNM	A	501	23/23	0.93	0.16	24,31,50,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.