



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 22, 2025 – 12:05 PM EST

PDB ID : 9CV8
Title : Crystal Structure of Cytochrome P450 NysL bound to nystatin
Authors : Murarka, V.C.; Poulos, T.L.
Deposited on : 2024-07-28
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	
1	B	394	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NysL.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	5	0
			3049	1943	548	549	9			
1	B	389	Total	C	N	O	S	0	11	0
			3075	1962	552	552	9			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is Nystatin (three-letter code: A1AZ7) (formula: $C_{47}H_{75}NO_{17}$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			65	47	1	17		
3	B	1	Total	C	N	O	0	0
			65	47	1	17		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	195	Total	O	0	0
			195	195		
5	B	340	Total	O	0	2
			342	342		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NysL



• Molecule 1: NysL



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	136.29Å 136.29Å 135.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.79 – 2.00 37.79 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.79-2.00) 99.8 (37.79-2.00)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 2.00Å)	Xtriage
Refinement program	PHENIX v1.18-3845-000	Depositor
R, R_{free}	0.189 , 0.215 0.191 , 0.217	Depositor DCC
R_{free} test set	4721 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6922	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, A1AZ7, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3127	0.54	0/4258
1	B	0.42	0/3171	0.61	0/4317
All	All	0.39	0/6298	0.57	0/8575

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3049	0	3125	11	0
1	B	3075	0	3167	19	0
2	A	43	0	30	2	0
2	B	43	0	30	2	0
3	A	65	0	0	3	0
3	B	65	0	0	4	0
4	A	25	0	0	0	0
4	B	20	0	0	0	0
5	A	195	0	0	2	0
5	B	342	0	0	3	0
All	All	6922	0	6352	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:402:A1AZ7:O13	3:A:402:A1AZ7:O12	2.07	0.71
3:A:402:A1AZ7:O06	5:A:501:HOH:O	2.08	0.70
1:B:25[B]:ARG:NH2	1:B:322:GLU:OE2	2.27	0.68
1:B:7:PRO:O	1:B:37[B]:ARG:NH1	2.31	0.64
1:B:263:LYS:HD3	1:B:266:ARG:NH2	2.13	0.64
1:B:183[A]:GLN:NE2	5:B:503:HOH:O	2.26	0.60
1:A:278:ALA:HA	1:A:313:PHE:CE1	2.39	0.57
1:B:55[A]:GLN:OE1	5:B:501:HOH:O	2.18	0.56
1:B:19:ARG:HH11	1:B:19:ARG:HB3	1.71	0.56
1:A:140:HIS:ND1	1:A:242:ASP:OD2	2.39	0.53
3:B:402:A1AZ7:O06	5:B:502:HOH:O	2.19	0.53
1:A:172:LYS:HA	1:A:175:GLU:HG3	1.91	0.52
1:A:162:GLU:HG2	1:A:192:LEU:HD11	1.92	0.52
1:B:31:ALA:O	1:B:48:THR:HG21	2.11	0.50
1:A:149:LEU:HD22	1:A:167:ILE:HG12	1.92	0.50
1:B:278:ALA:HA	1:B:313:PHE:CE1	2.47	0.50
2:A:401:HEM:HMC2	2:A:401:HEM:HBC2	1.92	0.50
1:B:28:GLN:HA	1:B:48:THR:CG2	2.42	0.50
1:B:32:PRO:HA	1:B:48:THR:HG22	1.94	0.49
1:B:263:LYS:HD3	1:B:266:ARG:HH21	1.78	0.49
3:B:402:A1AZ7:O13	3:B:402:A1AZ7:O12	2.32	0.47
3:B:402:A1AZ7:O13	3:B:402:A1AZ7:O14	2.33	0.47
1:B:13:GLU:HG3	1:B:14:VAL:H	1.80	0.46
1:A:170:LEU:HD13	1:A:234:LEU:HD22	1.96	0.46
3:A:402:A1AZ7:O06	3:A:402:A1AZ7:O02	2.34	0.44
1:A:165:LYS:HE2	1:A:169:ALA:HB2	2.00	0.44
1:B:48:THR:CG2	1:B:49:ARG:HG3	2.48	0.44
1:B:262:GLU:CD	1:B:262:GLU:H	2.21	0.43
1:B:15:PRO:HG3	1:B:21:SER:HB3	2.01	0.43
1:B:28:GLN:HA	1:B:48:THR:HG21	2.01	0.43
1:A:66:HIS:O	1:A:87:ASP:HA	2.19	0.42
1:B:48:THR:HG22	1:B:49:ARG:HG3	2.01	0.42
1:A:327[B]:ARG:NH2	5:A:502:HOH:O	2.22	0.42
1:B:19:ARG:HB3	1:B:19:ARG:NH1	2.34	0.42
1:A:162:GLU:HG2	1:A:192:LEU:CD1	2.49	0.42
1:B:345:GLY:HA3	2:B:401:HEM:C3C	2.55	0.41
2:B:401:HEM:HBB2	2:B:401:HEM:HMB2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:402:A1AZ7:O06	3:B:402:A1AZ7:O02	2.39	0.40
1:A:343:CYS:HB2	2:A:401:HEM:NA	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/394 (100%)	387 (98%)	6 (2%)	0	100	100
1	B	398/394 (101%)	391 (98%)	7 (2%)	0	100	100
All	All	791/788 (100%)	778 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/328 (100%)	326 (99%)	3 (1%)	75	81
1	B	334/328 (102%)	330 (99%)	4 (1%)	67	73
All	All	663/656 (101%)	656 (99%)	7 (1%)	70	76

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LYS
1	A	29	SER
1	A	165	LYS
1	B	48	THR
1	B	62	LEU
1	B	167	ILE
1	B	379	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	407	-	4,4,4	0.26	0	6,6,6	0.17	0
4	SO4	A	405	-	4,4,4	0.23	0	6,6,6	0.14	0
3	A1AZ7	B	402	2	67,67,67	1.44	10 (14%)	80,92,92	2.94	16 (20%)
4	SO4	A	404	-	4,4,4	0.25	0	6,6,6	0.14	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	B	403	-	4,4,4	0.22	0	6,6,6	0.30	0
2	HEM	B	401	1,3	42,50,50	1.53	7 (16%)	46,82,82	1.35	9 (19%)
3	A1AZ7	A	402	2	67,67,67	1.46	11 (16%)	80,92,92	2.97	18 (22%)
4	SO4	A	406	-	4,4,4	0.22	0	6,6,6	0.14	0
4	SO4	B	405	-	4,4,4	0.26	0	6,6,6	0.18	0
2	HEM	A	401	1,3	42,50,50	1.42	5 (11%)	46,82,82	1.41	9 (19%)
4	SO4	A	403	-	4,4,4	0.28	0	6,6,6	0.45	0
4	SO4	B	406	-	4,4,4	0.27	0	6,6,6	0.17	0
4	SO4	B	404	-	4,4,4	0.20	0	6,6,6	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	401	1,3	-	0/12/54/54	-
3	A1AZ7	A	402	2	-	20/71/109/109	0/2/3/3
2	HEM	A	401	1,3	-	0/12/54/54	-
3	A1AZ7	B	402	2	-	15/71/109/109	0/2/3/3

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	A1AZ7	O02-C19	6.43	1.49	1.40
2	B	401	HEM	C3C-C2C	-4.50	1.34	1.40
3	A	402	A1AZ7	O02-C19	4.21	1.46	1.40
3	A	402	A1AZ7	C25-C20	3.82	1.59	1.52
3	A	402	A1AZ7	O15-C49	-3.61	1.41	1.46
2	A	401	HEM	C3C-C2C	-3.56	1.35	1.40
3	A	402	A1AZ7	C47-C48	3.32	1.63	1.54
2	B	401	HEM	C3C-CAC	3.18	1.54	1.47
2	A	401	HEM	C3C-CAC	3.11	1.54	1.47
3	B	402	A1AZ7	O15-C49	-3.11	1.41	1.46
3	A	402	A1AZ7	C47-C49	2.96	1.60	1.53
3	B	402	A1AZ7	C25-C27	2.77	1.58	1.53
3	B	402	A1AZ7	C25-C20	2.72	1.57	1.52
3	B	402	A1AZ7	C54-C49	2.71	1.58	1.51
3	A	402	A1AZ7	C25-C27	2.71	1.58	1.53
2	A	401	HEM	CAB-C3B	2.67	1.54	1.47
2	B	401	HEM	CAB-C3B	2.64	1.54	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	A1AZ7	C21-C22	-2.61	1.50	1.54
2	A	401	HEM	CMB-C2B	2.54	1.56	1.50
2	B	401	HEM	C3C-C4C	2.41	1.44	1.41
2	B	401	HEM	CMA-C3A	2.37	1.56	1.51
2	B	401	HEM	CMB-C2B	2.35	1.55	1.50
3	A	402	A1AZ7	C31-C33	2.33	1.56	1.53
3	A	402	A1AZ7	C23-C19	2.32	1.55	1.52
3	B	402	A1AZ7	C43-C44	2.21	1.57	1.52
3	A	402	A1AZ7	O04-C27	2.16	1.48	1.44
2	B	401	HEM	CMD-C2D	2.13	1.55	1.50
3	B	402	A1AZ7	C53-C47	2.09	1.57	1.53
3	A	402	A1AZ7	C21-C28	2.06	1.56	1.51
3	B	402	A1AZ7	C34-C35	2.05	1.57	1.52
2	A	401	HEM	C3C-C4C	2.03	1.44	1.41
3	B	402	A1AZ7	C23-C22	-2.01	1.50	1.53
3	B	402	A1AZ7	C47-C48	2.00	1.59	1.54

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	A1AZ7	O01-C19-C24	13.27	118.62	106.53
3	B	402	A1AZ7	O01-C19-C24	12.88	118.26	106.53
3	B	402	A1AZ7	O01-C19-C23	12.25	123.58	108.92
3	A	402	A1AZ7	O01-C19-C23	11.84	123.08	108.92
3	A	402	A1AZ7	O02-C19-C24	-11.63	88.75	111.28
3	B	402	A1AZ7	O02-C19-C24	-9.99	91.93	111.28
3	A	402	A1AZ7	O02-C19-C23	-8.12	91.18	110.61
3	B	402	A1AZ7	O02-C19-C23	-7.77	92.02	110.61
3	B	402	A1AZ7	O15-C52-C46	7.42	124.73	111.43
3	A	402	A1AZ7	O15-C52-C46	6.92	123.84	111.43
3	B	402	A1AZ7	O15-C52-O17	-4.32	113.61	123.70
3	B	402	A1AZ7	C24-C26-C29	-3.41	105.54	113.73
3	A	402	A1AZ7	C49-O15-C52	3.36	124.35	118.36
3	A	402	A1AZ7	O01-C20-C25	3.26	110.05	105.95
3	B	402	A1AZ7	O15-C49-C54	3.24	114.68	108.20
3	B	402	A1AZ7	O12-C38-C40	-3.22	102.13	109.25
3	B	402	A1AZ7	C42-C40-C38	-3.17	109.26	114.28
3	A	402	A1AZ7	O15-C52-O17	-3.12	116.40	123.70
3	A	402	A1AZ7	C24-C26-C29	-2.98	106.56	113.73
3	B	402	A1AZ7	O04-C30-C31	2.93	115.31	108.09
3	B	402	A1AZ7	C53-C47-C49	-2.86	108.65	112.22
2	B	401	HEM	C2C-C3C-C4C	2.77	108.83	106.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	C4B-CHC-C1C	2.76	126.21	122.56
3	A	402	A1AZ7	C44-C43-C42	-2.68	110.03	114.28
3	A	402	A1AZ7	O12-C38-C40	-2.68	103.33	109.25
3	B	402	A1AZ7	C44-C43-C42	-2.67	110.05	114.28
3	B	402	A1AZ7	C54-C49-C47	-2.56	110.85	114.46
2	A	401	HEM	C4D-ND-C1D	2.55	108.23	105.21
3	A	402	A1AZ7	C54-C49-C47	-2.47	110.98	114.46
3	A	402	A1AZ7	O04-C30-C31	2.47	114.16	108.09
2	B	401	HEM	C4C-CHD-C1D	2.45	125.79	122.56
2	A	401	HEM	C4C-CHD-C1D	2.39	125.71	122.56
3	A	402	A1AZ7	C53-C47-C49	-2.37	109.27	112.22
2	A	401	HEM	CBA-CAA-C2A	-2.32	108.64	112.54
2	A	401	HEM	C1B-NB-C4B	2.31	107.95	105.21
2	B	401	HEM	CBA-CAA-C2A	-2.30	108.67	112.54
3	A	402	A1AZ7	O07-C28-C21	-2.21	117.20	122.84
2	B	401	HEM	C1B-NB-C4B	2.19	107.80	105.21
2	B	401	HEM	O2A-CGA-CBA	2.18	120.88	114.00
3	B	402	A1AZ7	C34-C33-C31	-2.18	106.33	111.06
2	B	401	HEM	CMC-C2C-C3C	2.15	128.97	124.68
2	B	401	HEM	CMA-C3A-C4A	-2.14	125.32	128.46
3	B	402	A1AZ7	O14-C44-C43	-2.13	104.55	109.25
2	A	401	HEM	C3B-C4B-NB	-2.11	107.95	109.47
3	A	402	A1AZ7	O17-C52-C46	-2.10	119.76	124.65
3	A	402	A1AZ7	O13-C42-C43	-2.10	104.61	109.25
2	A	401	HEM	C2D-C1D-ND	-2.09	107.49	109.90
2	B	401	HEM	O1A-CGA-CBA	-2.07	116.52	123.09
3	A	402	A1AZ7	O06-C26-C24	-2.05	104.48	109.52
2	B	401	HEM	CHC-C4B-NB	2.05	126.64	124.44
2	A	401	HEM	CHD-C1D-ND	2.04	126.63	124.44
2	A	401	HEM	CMC-C2C-C3C	2.02	128.72	124.68

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	A1AZ7	C23-C19-C24-C26
3	A	402	A1AZ7	O01-C19-C24-C26
3	A	402	A1AZ7	C20-C25-C27-C36
3	A	402	A1AZ7	C37-C38-C40-C42
3	A	402	A1AZ7	O12-C38-C40-C42
3	A	402	A1AZ7	C62-C63-C64-C65
3	B	402	A1AZ7	C23-C19-C24-C26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	402	A1AZ7	O01-C19-C24-C26
3	B	402	A1AZ7	C43-C44-C46-C52
3	B	402	A1AZ7	O14-C44-C46-C52
3	B	402	A1AZ7	C53-C47-C49-O15
3	A	402	A1AZ7	C42-C43-C44-O14
3	A	402	A1AZ7	C32-C37-C38-O12
3	B	402	A1AZ7	C46-C52-O15-C49
3	A	402	A1AZ7	C42-C43-C44-C46
3	B	402	A1AZ7	O17-C52-O15-C49
3	A	402	A1AZ7	C57-C59-C61-C65
3	A	402	A1AZ7	O14-C44-C46-C52
3	A	402	A1AZ7	C43-C44-C46-C52
3	A	402	A1AZ7	C21-C20-C25-C27
3	B	402	A1AZ7	C21-C20-C25-C27
3	B	402	A1AZ7	C20-C25-C27-C36
3	A	402	A1AZ7	C26-C29-C32-C37
3	A	402	A1AZ7	O08-C29-C32-C37
3	A	402	A1AZ7	C56-C58-C60-C62
3	B	402	A1AZ7	O13-C42-C43-C44
3	A	402	A1AZ7	C20-C21-C28-O09
3	B	402	A1AZ7	C37-C38-C40-C42
3	A	402	A1AZ7	C22-C21-C28-O09
3	A	402	A1AZ7	C20-C21-C28-O07
3	B	402	A1AZ7	O16-C48-C50-C56
3	B	402	A1AZ7	O12-C38-C40-C42
3	B	402	A1AZ7	C53-C47-C49-C54
3	B	402	A1AZ7	C48-C47-C49-O15
3	A	402	A1AZ7	C22-C21-C28-O07


There are no ring outliers.


4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	A1AZ7	4	0
2	B	401	HEM	2	0
3	A	402	A1AZ7	3	0
2	A	401	HEM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is


within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand A1AZ7 A 402





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/394 (98%)	0.21	22 (5%) 31 29	20, 45, 83, 141	5 (1%)
1	B	389/394 (98%)	-0.08	13 (3%) 49 47	17, 34, 59, 141	11 (2%)
All	All	779/788 (98%)	0.06	35 (4%) 39 37	17, 39, 71, 141	16 (2%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	12	ALA	6.0
1	B	295	ASP	4.9
1	A	5	THR	4.7
1	A	11	LYS	4.5
1	A	6	ALA	4.2
1	B	110[A]	ARG	3.7
1	A	13	GLU	3.7
1	B	13	GLU	3.7
1	A	14	VAL	3.6
1	A	16	PRO	3.6
1	B	113	MET	3.5
1	B	14	VAL	3.5
1	B	12	ALA	3.3
1	A	164	GLY	3.2
1	A	294	GLY	3.2
1	B	319	ASP	3.1
1	A	165	LYS	2.8
1	A	15	PRO	2.8
1	A	262	GLU	2.8
1	A	322	GLU	2.7
1	A	110	ARG	2.7
1	A	129	HIS	2.7
1	A	113	MET	2.6
1	B	253	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	15	PRO	2.5
1	A	177	ASP	2.5
1	B	262	GLU	2.4
1	B	161	GLU	2.2
1	B	34	CYS	2.2
1	A	30	ARG	2.1
1	A	37	ARG	2.1
1	B	37[A]	ARG	2.1
1	A	162	GLU	2.1
1	A	114	ASP	2.1
1	A	19	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	B	406	5/5	0.70	0.12	79,87,104,105	0
4	SO4	A	407	5/5	0.74	0.11	87,90,104,105	0
4	SO4	A	404	5/5	0.76	0.13	94,94,113,127	0
4	SO4	A	405	5/5	0.77	0.10	111,119,132,133	0
4	SO4	A	406	5/5	0.83	0.09	96,98,121,125	0
4	SO4	B	405	5/5	0.88	0.09	68,80,90,103	0
3	A1AZ7	A	402	65/65	0.94	0.10	29,49,59,64	0
3	A1AZ7	B	402	65/65	0.95	0.09	22,37,51,54	0
4	SO4	B	403	5/5	0.96	0.11	49,50,57,59	0
4	SO4	B	404	5/5	0.96	0.09	46,53,55,56	0
4	SO4	A	403	5/5	0.98	0.08	46,46,50,54	0
2	HEM	B	401	43/43	0.99	0.04	19,24,31,33	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	HEM	A	401	43/43	0.99	0.05	22,30,36,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around A1AZ7 B 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.