



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 27, 2025 – 04:05 PM EDT

PDB ID : 9E6E / pdb_00009e6e
Title : Yeast Fzf1 Zn-fingers 1-3 bound to YHB1 26 bp DNA
Authors : Moore, S.A.; Ma, C.; Xiao, W.
Deposited on : 2024-10-29
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1044 (2.98-2.94)
Clashscore	180529	1097 (2.98-2.94)
Ramachandran outliers	177936	1049 (2.98-2.94)
Sidechain outliers	177891	1049 (2.98-2.94)
RSRZ outliers	164620	1044 (2.98-2.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	
1	D	299	
2	B	27	
2	E	27	
3	C	27	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	27	 52% 48%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6339 atoms, of which 2638 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zinc finger protein FZF1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	D	90	Total	C	H	N	O	S	0	0	0
			1404	444	684	145	124	7			
1	A	94	Total	C	H	N	O	S	0	1	0
			1488	467	730	156	128	7			

- Molecule 2 is a DNA chain called DNA 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	E	27	Total	C	H	N	O	P	0	0	0
			858	264	304	108	156	26			
2	B	27	Total	C	H	N	O	P	0	0	0
			858	264	304	108	156	26			

- Molecule 3 is a DNA chain called DNA 2.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	F	27	Total	C	H	N	O	P	0	0	0
			855	263	308	94	164	26			
3	C	27	Total	C	H	N	O	P	0	0	0
			855	263	308	94	164	26			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	3	Total	Zn	0	0
			3	3		
4	A	3	Total	Zn	0	0
			3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	4	Total 4	O 4	0	0
5	E	1	Total 1	O 1	0	0
5	B	1	Total 1	O 1	0	0
5	C	2	Total 2	O 2	0	0
5	A	7	Total 7	O 7	0	0



● Molecule 3: DNA 2



● Molecule 3: DNA 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.91Å 66.27Å 174.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.37 – 2.95 36.37 – 2.95	Depositor EDS
% Data completeness (in resolution range)	98.9 (36.37-2.95) 99.7 (36.37-2.95)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.95Å)	Xtriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, R_{free}	0.218 , 0.247 0.217 , 0.248	Depositor DCC
R_{free} test set	781 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	54.8	Xtriage
Anisotropy	0.962	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6339	wwPDB-VP
Average B, all atoms (Å ²)	94.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/781	0.37	0/1052
1	D	0.17	0/740	0.38	0/999
2	B	0.35	0/623	0.65	0/960
2	E	0.31	0/623	0.60	0/960
3	C	0.33	0/611	0.59	0/941
3	F	0.28	0/611	0.57	0/941
All	All	0.28	0/3989	0.54	0/5853

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	758	730	713	6	0
1	D	720	684	667	2	0
2	B	554	304	304	18	0
2	E	554	304	304	16	0
3	C	547	308	308	6	0
3	F	547	308	308	11	0
4	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	3	0	0	0	0
5	A	7	0	0	1	0
5	B	1	0	0	0	0
5	C	2	0	0	2	0
5	D	4	0	0	0	0
5	E	1	0	0	0	0
All	All	3701	2638	2604	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 58 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:21:DC:H2'	2:B:22:DG:C8	2.30	0.67
3:C:14:DT:OP2	5:C:101:HOH:O	2.12	0.66
2:E:21:DC:H2'	2:E:22:DG:C8	2.34	0.63
3:C:3:DG:H2'	3:C:4:DC:C6	2.33	0.63
1:A:76:LEU:HD12	1:A:94:HIS:CD2	2.34	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	93/299 (31%)	85 (91%)	8 (9%)	0	100	100
1	D	88/299 (29%)	82 (93%)	6 (7%)	0	100	100
All	All	181/598 (30%)	167 (92%)	14 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/277 (29%)	80 (99%)	1 (1%)	67	82
1	D	78/277 (28%)	76 (97%)	2 (3%)	41	65
All	All	159/554 (29%)	156 (98%)	3 (2%)	52	73

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	57	CYS
1	D	75	THR
1	A	17	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	39	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	94/299 (31%)	0.34	3 (3%)	50	47	37, 69, 117, 130	1 (1%)
1	D	90/299 (30%)	0.47	6 (6%)	25	24	53, 85, 120, 132	0
2	B	27/27 (100%)	-0.11	0	100	100	49, 96, 172, 191	0
2	E	27/27 (100%)	-0.08	0	100	100	58, 102, 176, 188	0
3	C	27/27 (100%)	0.02	0	100	100	53, 88, 174, 195	0
3	F	27/27 (100%)	0.11	0	100	100	75, 96, 172, 210	0
All	All	292/706 (41%)	0.25	9 (3%)	51	48	37, 81, 166, 210	1 (0%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	18	GLY	3.4
1	A	104	ILE	3.4
1	A	20	GLU	2.9
1	D	38	ASN	2.7
1	D	20	GLU	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ZN	D	303	1/1	0.98	0.04	102,102,102,102	0
4	ZN	D	302	1/1	0.99	0.05	58,58,58,58	0
4	ZN	D	301	1/1	0.99	0.03	66,66,66,66	0
4	ZN	A	302	1/1	0.99	0.06	63,63,63,63	0
4	ZN	A	301	1/1	1.00	0.06	56,56,56,56	0
4	ZN	A	303	1/1	1.00	0.03	61,61,61,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN D 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.