



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 10, 2025 – 12:23 PM EST

PDB ID : 9ECF / pdb\_00009ecf  
Title : Crystal structure of the hERbeta LBD complexed with androstenediol and SRC 2-2 peptide (crystal form 2)  
Authors : Pederick, J.L.; Bruning, J.B.  
Deposited on : 2024-11-14  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	<div> <div>6%</div> <div> <div></div> <div>75%</div> <div>13%</div> <div>11%</div> </div> </div>
1	B	247	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>11%</div> </div> </div>
1	E	247	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>6%</div> </div> </div>
1	F	247	<div> <div>2%</div> <div> <div></div> <div>74%</div> <div>12%</div> <div>13%</div> </div> </div>
1	I	247	<div> <div>6%</div> <div> <div></div> <div>72%</div> <div>15%</div> <div>13%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	J	247	
1	M	247	
1	N	247	
2	C	13	
2	D	13	
2	G	13	
2	H	13	
2	K	13	
2	L	13	
2	O	13	
2	P	13	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14712 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Estrogen receptor beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	231	Total	C	N	O	S	0	1	0
			1773	1140	293	319	21			
1	F	214	Total	C	N	O	S	0	2	0
			1667	1075	277	295	20			
1	A	219	Total	C	N	O	S	0	0	0
			1684	1086	284	296	18			
1	B	219	Total	C	N	O	S	0	0	0
			1671	1075	280	297	19			
1	I	216	Total	C	N	O	S	0	1	0
			1670	1074	278	299	19			
1	J	196	Total	C	N	O	S	0	0	0
			1447	936	244	248	19			
1	M	219	Total	C	N	O	S	0	0	0
			1676	1084	280	294	18			
1	N	218	Total	C	N	O	S	0	0	0
			1681	1080	281	302	18			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	254	MET	-	expression tag	UNP Q92731
E	255	HIS	-	expression tag	UNP Q92731
E	256	HIS	-	expression tag	UNP Q92731
E	257	HIS	-	expression tag	UNP Q92731
E	258	HIS	-	expression tag	UNP Q92731
E	259	HIS	-	expression tag	UNP Q92731
E	260	HIS	-	expression tag	UNP Q92731
F	254	MET	-	expression tag	UNP Q92731
F	255	HIS	-	expression tag	UNP Q92731
F	256	HIS	-	expression tag	UNP Q92731
F	257	HIS	-	expression tag	UNP Q92731
F	258	HIS	-	expression tag	UNP Q92731
F	259	HIS	-	expression tag	UNP Q92731

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	260	HIS	-	expression tag	UNP Q92731
A	254	MET	-	expression tag	UNP Q92731
A	255	HIS	-	expression tag	UNP Q92731
A	256	HIS	-	expression tag	UNP Q92731
A	257	HIS	-	expression tag	UNP Q92731
A	258	HIS	-	expression tag	UNP Q92731
A	259	HIS	-	expression tag	UNP Q92731
A	260	HIS	-	expression tag	UNP Q92731
B	254	MET	-	expression tag	UNP Q92731
B	255	HIS	-	expression tag	UNP Q92731
B	256	HIS	-	expression tag	UNP Q92731
B	257	HIS	-	expression tag	UNP Q92731
B	258	HIS	-	expression tag	UNP Q92731
B	259	HIS	-	expression tag	UNP Q92731
B	260	HIS	-	expression tag	UNP Q92731
I	254	MET	-	expression tag	UNP Q92731
I	255	HIS	-	expression tag	UNP Q92731
I	256	HIS	-	expression tag	UNP Q92731
I	257	HIS	-	expression tag	UNP Q92731
I	258	HIS	-	expression tag	UNP Q92731
I	259	HIS	-	expression tag	UNP Q92731
I	260	HIS	-	expression tag	UNP Q92731
J	254	MET	-	expression tag	UNP Q92731
J	255	HIS	-	expression tag	UNP Q92731
J	256	HIS	-	expression tag	UNP Q92731
J	257	HIS	-	expression tag	UNP Q92731
J	258	HIS	-	expression tag	UNP Q92731
J	259	HIS	-	expression tag	UNP Q92731
J	260	HIS	-	expression tag	UNP Q92731
M	254	MET	-	expression tag	UNP Q92731
M	255	HIS	-	expression tag	UNP Q92731
M	256	HIS	-	expression tag	UNP Q92731
M	257	HIS	-	expression tag	UNP Q92731
M	258	HIS	-	expression tag	UNP Q92731
M	259	HIS	-	expression tag	UNP Q92731
M	260	HIS	-	expression tag	UNP Q92731
N	254	MET	-	expression tag	UNP Q92731
N	255	HIS	-	expression tag	UNP Q92731
N	256	HIS	-	expression tag	UNP Q92731
N	257	HIS	-	expression tag	UNP Q92731
N	258	HIS	-	expression tag	UNP Q92731
N	259	HIS	-	expression tag	UNP Q92731

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
N	260	HIS	-	expression tag	UNP Q92731

- Molecule 2 is a protein called Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	9	Total	C	N	O	0	0	0
			77	50	17	10			
2	H	9	Total	C	N	O	0	0	0
			66	44	12	10			
2	C	10	Total	C	N	O	0	0	0
			81	51	16	14			
2	D	9	Total	C	N	O	0	0	0
			67	45	13	9			
2	K	12	Total	C	N	O	0	0	0
			96	60	20	16			
2	L	9	Total	C	N	O	0	0	0
			70	45	16	9			
2	O	12	Total	C	N	O	0	0	0
			100	63	21	16			
2	P	11	Total	C	N	O	0	0	0
			86	55	18	13			

- Molecule 3 is (3 $\alpha$ ,8 $\alpha$ ,17 $\beta$ )-androst-5-ene-3,17-diol (CCD ID: B81) (formula: C<sub>19</sub>H<sub>30</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total C O 21 19 2	0	0
3	F	1	Total C O 21 19 2	0	0
3	A	1	Total C O 21 19 2	0	0
3	B	1	Total C O 21 19 2	0	0
3	I	1	Total C O 21 19 2	0	0
3	J	1	Total C O 21 19 2	0	0
3	M	1	Total C O 21 19 2	0	0
3	N	1	Total C O 21 19 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	E	86	Total O 86 86	0	0
4	F	87	Total O 87 87	0	0
4	G	2	Total O 2 2	0	0
4	A	74	Total O 74 74	0	0
4	B	66	Total O 66 66	0	0
4	C	1	Total O 1 1	0	0
4	D	1	Total O 1 1	0	0
4	I	98	Total O 98 98	0	0
4	J	38	Total O 38 38	0	0
4	K	4	Total O 4 4	0	0
4	L	2	Total O 2 2	0	0
4	M	97	Total O 97 97	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	N	73	Total 73	O 73	0	1
4	O	2	Total 2	O 2	0	0
4	P	1	Total 1	O 1	0	0




### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Estrogen receptor beta



- Molecule 1: Estrogen receptor beta




- Molecule 1: Estrogen receptor beta




- Molecule 1: Estrogen receptor beta








• Molecule 1: Estrogen receptor beta




• Molecule 1: Estrogen receptor beta



• Molecule 1: Estrogen receptor beta




• Molecule 1: Estrogen receptor beta






• Molecule 2: Nuclear receptor coactivator 2




• Molecule 2: Nuclear receptor coactivator 2




• Molecule 2: Nuclear receptor coactivator 2



• Molecule 2: Nuclear receptor coactivator 2




• Molecule 2: Nuclear receptor coactivator 2




• Molecule 2: Nuclear receptor coactivator 2






• Molecule 2: Nuclear receptor coactivator 2



• Molecule 2: Nuclear receptor coactivator 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.53Å 108.38Å 120.60Å 90.00° 104.23° 90.00°	Depositor
Resolution (Å)	38.97 – 2.00 38.97 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (38.97-2.00) 99.3 (38.97-2.00)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.250 , 0.278 0.251 , 0.278	Depositor DCC
$R_{free}$ test set	6412 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtriage
Anisotropy	0.465	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	14712	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 57.36 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.3936e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B81

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.09	0/1712	0.25	0/2314
1	B	0.09	0/1699	0.26	0/2300
1	E	0.09	0/1804	0.26	0/2447
1	F	0.09	0/1695	0.26	0/2291
1	I	0.12	0/1697	0.29	0/2294
1	J	0.09	0/1470	0.27	0/1992
1	M	0.08	0/1705	0.26	0/2311
1	N	0.09	0/1707	0.25	0/2310
2	C	0.10	0/81	0.34	0/108
2	D	0.09	0/68	0.17	0/92
2	G	0.05	0/78	0.13	0/104
2	H	0.09	0/66	0.23	0/89
2	K	0.06	0/97	0.27	0/130
2	L	0.07	0/71	0.16	0/95
2	O	0.12	0/101	0.26	0/134
2	P	0.09	0/87	0.20	0/117
All	All	0.09	0/14138	0.26	0/19128

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1684	0	1732	16	0
1	B	1671	0	1684	18	0
1	E	1773	0	1803	18	0
1	F	1667	0	1698	22	0
1	I	1670	0	1703	25	0
1	J	1447	0	1410	16	0
1	M	1676	0	1711	15	0
1	N	1681	0	1719	22	0
2	C	81	0	82	1	0
2	D	67	0	63	0	0
2	G	77	0	80	1	0
2	H	66	0	64	2	0
2	K	96	0	94	2	0
2	L	70	0	65	1	0
2	O	100	0	105	2	0
2	P	86	0	80	2	0
3	A	21	0	30	1	0
3	B	21	0	30	1	0
3	E	21	0	30	2	0
3	F	21	0	30	1	0
3	I	21	0	30	2	0
3	J	21	0	30	2	0
3	M	21	0	30	3	0
3	N	21	0	30	2	0
4	A	74	0	0	0	0
4	B	66	0	0	1	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	86	0	0	0	0
4	F	87	0	0	3	0
4	G	2	0	0	0	0
4	I	98	0	0	3	0
4	J	38	0	0	2	0
4	K	4	0	0	0	0
4	L	2	0	0	0	0
4	M	97	0	0	0	0
4	N	73	0	0	1	0
4	O	2	0	0	0	0
4	P	1	0	0	0	0
All	All	14712	0	14333	144	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:376:ILE:HA	1:I:379:MET:HE2	1.63	0.78
1:A:314:LYS:NZ	2:C:11:ASP:O	2.22	0.72
1:M:332:GLU:OE2	2:O:2:HIS:NE2	2.23	0.71
1:I:332:GLU:OE2	2:K:2:HIS:NE2	2.27	0.66
1:F:407:ASN:ND2	4:F:703:HOH:O	2.30	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/247 (86%)	209 (98%)	4 (2%)	0	100	100
1	B	213/247 (86%)	209 (98%)	4 (2%)	0	100	100
1	E	228/247 (92%)	226 (99%)	2 (1%)	0	100	100
1	F	210/247 (85%)	207 (99%)	3 (1%)	0	100	100
1	I	211/247 (85%)	208 (99%)	3 (1%)	0	100	100
1	J	184/247 (74%)	181 (98%)	3 (2%)	0	100	100
1	M	213/247 (86%)	208 (98%)	4 (2%)	1 (0%)	25	21
1	N	212/247 (86%)	208 (98%)	4 (2%)	0	100	100
2	C	8/13 (62%)	8 (100%)	0	0	100	100
2	D	7/13 (54%)	7 (100%)	0	0	100	100
2	G	7/13 (54%)	7 (100%)	0	0	100	100
2	H	7/13 (54%)	7 (100%)	0	0	100	100
2	K	10/13 (77%)	10 (100%)	0	0	100	100
2	L	7/13 (54%)	7 (100%)	0	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	O	10/13 (77%)	8 (80%)	2 (20%)	0	100	100
2	P	9/13 (69%)	8 (89%)	1 (11%)	0	100	100
All	All	1749/2080 (84%)	1718 (98%)	30 (2%)	1 (0%)	48	47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	486	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/222 (84%)	178 (96%)	8 (4%)	25	23
1	B	181/222 (82%)	177 (98%)	4 (2%)	47	51
1	E	197/222 (89%)	194 (98%)	3 (2%)	60	66
1	F	184/222 (83%)	181 (98%)	3 (2%)	58	64
1	I	184/222 (83%)	179 (97%)	5 (3%)	40	42
1	J	144/222 (65%)	141 (98%)	3 (2%)	48	53
1	M	182/222 (82%)	181 (100%)	1 (0%)	86	90
1	N	186/222 (84%)	182 (98%)	4 (2%)	47	51
2	C	9/13 (69%)	8 (89%)	1 (11%)	5	3
2	D	6/13 (46%)	6 (100%)	0	100	100
2	G	8/13 (62%)	7 (88%)	1 (12%)	3	2
2	H	6/13 (46%)	6 (100%)	0	100	100
2	K	10/13 (77%)	9 (90%)	1 (10%)	6	3
2	L	6/13 (46%)	6 (100%)	0	100	100
2	O	11/13 (85%)	11 (100%)	0	100	100
2	P	8/13 (62%)	8 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1508/1880 (80%)	1474 (98%)	34 (2%)	45 49

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	K	12	SER
1	M	484	VAL
1	N	481	CYS
1	A	483	ASN
1	A	466	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	394	HIS
1	J	470	ASN
1	J	467	HIS
2	L	6	HIS
1	A	407	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	B81	M	601	-	24,24,24	1.84	7 (29%)	39,39,39	2.56	15 (38%)
3	B81	E	601	-	24,24,24	1.88	6 (25%)	39,39,39	2.53	15 (38%)
3	B81	A	601	-	24,24,24	1.88	7 (29%)	39,39,39	2.58	16 (41%)
3	B81	J	601	-	24,24,24	1.83	6 (25%)	39,39,39	2.59	16 (41%)
3	B81	I	601	-	24,24,24	1.84	6 (25%)	39,39,39	2.53	14 (35%)
3	B81	N	601	-	24,24,24	1.86	6 (25%)	39,39,39	2.58	16 (41%)
3	B81	F	601	-	24,24,24	1.91	7 (29%)	39,39,39	2.57	16 (41%)
3	B81	B	601	-	24,24,24	1.90	7 (29%)	39,39,39	2.58	15 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	B81	M	601	-	-	-	0/4/4/4
3	B81	E	601	-	-	-	0/4/4/4
3	B81	A	601	-	-	-	0/4/4/4
3	B81	J	601	-	-	-	0/4/4/4
3	B81	I	601	-	-	-	0/4/4/4
3	B81	N	601	-	-	-	0/4/4/4
3	B81	F	601	-	-	-	0/4/4/4
3	B81	B	601	-	-	-	0/4/4/4

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	601	B81	C10-C9	5.26	1.64	1.56
3	E	601	B81	C10-C9	5.23	1.64	1.56
3	B	601	B81	C10-C9	5.22	1.64	1.56
3	A	601	B81	C10-C9	5.14	1.64	1.56
3	N	601	B81	C10-C9	4.98	1.64	1.56

The worst 5 of 123 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	N	601	B81	C14-C13-C17	6.48	105.86	99.25
3	A	601	B81	C14-C13-C17	6.33	105.70	99.25
3	B	601	B81	C14-C13-C17	6.28	105.65	99.25
3	J	601	B81	C14-C13-C17	6.26	105.63	99.25
3	F	601	B81	C14-C13-C17	6.17	105.54	99.25

There are no chirality outliers.


There are no torsion outliers.



There are no ring outliers.


8 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	M	601	B81	3	0
3	E	601	B81	2	0
3	A	601	B81	1	0
3	J	601	B81	2	0
3	I	601	B81	2	0
3	N	601	B81	2	0
3	F	601	B81	1	0
3	B	601	B81	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	219/247 (88%)	0.62	15 (6%) 25 23	22, 35, 53, 60	0
1	B	219/247 (88%)	0.32	7 (3%) 50 48	18, 29, 47, 60	0
1	E	231/247 (93%)	0.40	8 (3%) 47 45	18, 30, 48, 56	1 (0%)
1	F	214/247 (86%)	0.22	6 (2%) 55 53	10, 28, 44, 51	2 (0%)
1	I	216/247 (87%)	0.49	14 (6%) 26 24	17, 31, 56, 65	1 (0%)
1	J	196/247 (79%)	1.36	47 (23%) 2 2	31, 48, 65, 71	0
1	M	219/247 (88%)	0.22	7 (3%) 50 48	17, 28, 47, 56	0
1	N	218/247 (88%)	0.51	9 (4%) 42 40	22, 34, 57, 67	0
2	C	10/13 (76%)	1.25	1 (10%) 14 13	52, 58, 63, 66	0
2	D	9/13 (69%)	1.42	1 (11%) 12 10	48, 51, 58, 60	0
2	G	9/13 (69%)	0.99	1 (11%) 12 10	46, 50, 55, 56	0
2	H	9/13 (69%)	0.98	1 (11%) 12 10	45, 47, 55, 59	0
2	K	12/13 (92%)	0.91	1 (8%) 19 17	36, 44, 52, 53	0
2	L	9/13 (69%)	1.53	3 (33%) 1 1	53, 57, 62, 66	0
2	O	12/13 (92%)	1.11	0 100 100	36, 51, 55, 56	0
2	P	11/13 (84%)	1.25	1 (9%) 16 15	37, 41, 51, 54	0
All	All	1813/2080 (87%)	0.54	122 (6%) 25 23	10, 34, 57, 71	4 (0%)

The worst 5 of 122 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	324	LEU	4.7
1	J	269	VAL	4.7
1	J	442	ALA	4.4
1	I	456	ALA	4.3
1	J	264	SER	4.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	B81	E	601	21/21	0.89	0.09	20,22,25,27	0
3	B81	J	601	21/21	0.89	0.13	29,36,39,41	0
3	B81	A	601	21/21	0.90	0.10	23,27,29,33	0
3	B81	I	601	21/21	0.91	0.08	15,17,20,21	0
3	B81	B	601	21/21	0.91	0.08	17,24,27,29	0
3	B81	N	601	21/21	0.92	0.10	27,33,36,37	0
3	B81	M	601	21/21	0.93	0.08	15,18,20,22	0
3	B81	F	601	21/21	0.93	0.07	15,19,23,24	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around B81 E 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around B81 J 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around B81 A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around B81 I 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around B81 B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around B81 N 601:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around B81 M 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around B81 F 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.