



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2025 – 10:19 AM EDT

PDB ID : 9EJI
Title : Peptide-independent T cell receptor recognition of HLA-DQ2
Authors : Lim, J.J.; Loh, T.J.; Reid, H.H.; Rossjohn, J.
Deposited on : 2024-11-27
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	<div> <div>4%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> </div>
2	B	196	<div> <div>8%</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>8%</div> </div> </div>
3	C	13	<div> <div>8%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
4	D	204	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>6%</div> </div> </div>
5	E	242	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>10%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	2	 100%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 7042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen DQ alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1460	945	234	279	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	181	THR	-	expression tag	UNP Q08AS3

- Molecule 2 is a protein called HLA class II histocompatibility antigen DQ beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	181	Total	C	N	O	S	0	0	0
			1452	920	258	267	7			

- Molecule 3 is a protein called glut-L1 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	0	0	0
			99	62	16	21			

- Molecule 4 is a protein called G9 T cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	199	Total	C	N	O	S	0	0	0
			1543	967	259	307	10			

- Molecule 5 is a protein called G9 T cell receptor beta chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	241	Total	C	N	O	S	0	0	0
			1898	1195	336	361	6			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	C	1	Total	C	O	0	0
			4	2	2		
9	D	1	Total	C	O	0	0
			4	2	2		
9	E	1	Total	C	O	4	0
			4	2	2		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	119	Total 119	O 119	0	0
10	B	97	Total 97	O 97	0	0
10	C	4	Total 4	O 4	0	0
10	D	152	Total 152	O 152	0	0
10	E	152	Total 152	O 152	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen DQ alpha chain

Chain A: 



- Molecule 2: HLA class II histocompatibility antigen DQ beta chain

Chain B: 



- Molecule 3: glut-L1 peptide

Chain C: 





- Molecule 4: G9 T cell receptor alpha chain

Chain D: 



- Molecule 5: G9 T cell receptor beta chain

Chain E: 



- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	101.84Å 81.83Å 143.17Å 90.00° 107.92° 90.00°	Depositor
Resolution (Å)	48.45 – 2.20 48.45 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.45-2.20) 99.9 (48.45-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.30 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.185 , 0.220 0.184 , 0.220	Depositor DCC
R_{free} test set	2957 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	1.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7042	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, GOL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1504	0.45	0/2056
2	B	0.24	0/1484	0.52	0/2018
3	C	0.27	0/102	0.45	0/141
4	D	0.25	0/1577	0.51	0/2140
5	E	0.26	0/1948	0.52	0/2647
All	All	0.25	0/6615	0.50	0/9002

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1460	0	1400	5	0
2	B	1452	0	1397	7	0
3	C	99	0	96	1	0
4	D	1543	0	1455	6	0
5	E	1898	0	1795	13	0
6	F	28	0	25	0	0
7	A	6	0	8	0	0
7	E	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	14	0	13	0	0
9	C	4	0	3	0	0
9	D	4	0	3	0	0
9	E	4	0	3	0	0
10	A	119	0	0	0	0
10	B	97	0	0	0	0
10	C	4	0	0	1	0
10	D	152	0	0	2	0
10	E	152	0	0	2	0
All	All	7042	0	6206	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:LYS:HG3	1:A:59:LEU:HD11	1.65	0.78
4:D:171:LYS:NZ	10:D:401:HOH:O	2.16	0.77
3:C:0:GLN:N	10:C:501:HOH:O	2.37	0.57
4:D:124:PRO:HD2	10:D:401:HOH:O	2.04	0.56
5:E:183:CYS:HB3	5:E:205:ARG:HD3	1.87	0.55
5:E:142:PRO:HD3	5:E:155:LEU:HD13	1.89	0.54
5:E:111:GLU:HA	7:E:302:GOL:H31	1.91	0.52
5:E:183:CYS:O	5:E:205:ARG:HD2	2.10	0.52
4:D:39:PHE:HB2	4:D:56:VAL:HB	1.92	0.52
2:B:70:ARG:HG2	2:B:70:ARG:HH11	1.76	0.50
5:E:230:SER:OG	5:E:233:ASP:OD2	2.30	0.49
5:E:37:ARG:O	10:E:401:HOH:O	2.20	0.49
5:E:139:VAL:HG23	5:E:249:ALA:HB3	1.95	0.49
4:D:96:LEU:HD11	4:D:182:LYS:HD3	1.94	0.48
1:A:158:GLU:C	1:A:159:ILE:HD12	2.36	0.47
4:D:46:CYS:SG	10:E:542:HOH:O	2.61	0.46
2:B:37:ILE:HG13	2:B:38:VAL:HG23	1.96	0.46
5:E:58:SER:O	5:E:58:SER:OG	2.28	0.46
5:E:96:LEU:HD11	5:E:127:GLU:OE1	2.16	0.46
5:E:128:ASP:OD1	5:E:130:ASN:ND2	2.48	0.46
1:A:133:GLU:OE2	1:A:146:LYS:NZ	2.40	0.45
2:B:97:PRO:HB3	2:B:122:PHE:HB3	1.99	0.45
2:B:143:VAL:HG23	2:B:160:MET:HB2	2.00	0.43
5:E:68:LYS:HG2	5:E:69:GLY:N	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:SER:C	1:A:9:GLY:HA2	2.39	0.43
2:B:70:ARG:HG2	2:B:70:ARG:NH1	2.32	0.43
5:E:165:ASP:HB2	5:E:188:PRO:HG2	2.00	0.42
2:B:116:VAL:HG22	2:B:160:MET:HG3	2.01	0.42
5:E:235:TRP:CG	5:E:241:LYS:HG3	2.55	0.42
4:D:79:GLN:HE21	4:D:79:GLN:HB2	1.64	0.42
2:B:71:LYS:HA	2:B:71:LYS:HD2	1.80	0.41
1:A:44:LYS:HA	1:A:44:LYS:HD2	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/184 (98%)	177 (98%)	2 (1%)	1 (1%)	22	23
2	B	175/196 (89%)	169 (97%)	5 (3%)	1 (1%)	22	23
3	C	11/13 (85%)	11 (100%)	0	0	100	100
4	D	197/204 (97%)	193 (98%)	3 (2%)	1 (0%)	25	28
5	E	239/242 (99%)	233 (98%)	6 (2%)	0	100	100
All	All	802/839 (96%)	783 (98%)	16 (2%)	3 (0%)	30	34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	8	PRO
1	A	18	PRO
2	B	165	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/169 (98%)	164 (99%)	2 (1%)	67	80
2	B	157/178 (88%)	154 (98%)	3 (2%)	52	67
3	C	12/12 (100%)	12 (100%)	0	100	100
4	D	174/185 (94%)	169 (97%)	5 (3%)	37	50
5	E	205/211 (97%)	198 (97%)	7 (3%)	32	42
All	All	714/755 (95%)	697 (98%)	17 (2%)	44	57

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	PRO
1	A	164	VAL
2	B	26	LEU
2	B	130	ARG
2	B	182	SER
4	D	58	SER
4	D	79	GLN
4	D	86	TYR
4	D	172	CYS
4	D	214	SER
5	E	18	GLN
5	E	53	LEU
5	E	54	PHE
5	E	58	SER
5	E	83	SER
5	E	205	ARG
5	E	248	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	126	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	79	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	F	1	1,6	14,14,15	0.38	0	17,19,21	0.57	0
6	NAG	F	2	6	14,14,15	0.27	0	17,19,21	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	F	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	F	2	6	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	F	2	NAG	C4-C5-C6-O6
6	F	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	NAG	B	201	2	14,14,15	0.22	0	17,19,21	0.47	0
9	ACT	D	301	-	3,3,3	1.31	0	3,3,3	1.41	0
9	ACT	E	301	-	3,3,3	1.37	0	3,3,3	1.37	0
7	GOL	A	201	-	5,5,5	0.95	0	5,5,5	1.07	0
7	GOL	E	302	-	5,5,5	0.86	0	5,5,5	1.16	1 (20%)
9	ACT	C	101	-	3,3,3	1.31	0	3,3,3	1.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	E	302	-	-	0/4/4/4	-
8	NAG	B	201	2	-	0/6/23/26	0/1/1/1
7	GOL	A	201	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	302	GOL	C3-C2-C1	-2.09	104.14	111.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	E	302	GOL	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/184 (98%)	-0.05	7 (3%) 44 41	22, 32, 53, 75	0
2	B	181/196 (92%)	0.29	15 (8%) 19 16	23, 33, 70, 87	0
3	C	13/13 (100%)	0.38	1 (7%) 21 18	31, 38, 53, 65	0
4	D	199/204 (97%)	-0.01	8 (4%) 43 39	17, 28, 51, 58	0
5	E	241/242 (99%)	0.02	12 (4%) 35 32	19, 29, 56, 74	0
All	All	816/839 (97%)	0.06	43 (5%) 33 30	17, 31, 58, 87	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	9	GLY	4.5
2	B	193	ALA	4.2
4	D	194	ASP	4.1
1	A	17	GLY	4.1
4	D	140	LYS	3.9
2	B	194	GLN	3.9
2	B	164	THR	3.8
5	E	234	GLU	3.5
1	A	18	PRO	3.3
2	B	166	GLN	3.3
5	E	232	ASN	3.3
5	E	196	ASN	3.2
2	B	140	ALA	3.1
2	B	165	PRO	3.1
2	B	167	ARG	2.9
1	A	48	PHE	2.9
5	E	231	GLU	2.9
5	E	195	LEU	2.9
5	E	194	ALA	2.7
3	C	0	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	162	LYS	2.7
5	E	127	GLU	2.7
5	E	92	SER	2.7
2	B	136	GLN	2.6
1	A	181	THR	2.6
1	A	159	ILE	2.5
2	B	104	SER	2.5
4	D	163	ASP	2.5
1	A	157	ASP	2.5
2	B	162	GLU	2.5
2	B	139	THR	2.4
2	B	3	SER	2.4
5	E	49	GLY	2.4
5	E	93	THR	2.3
2	B	135	GLY	2.3
2	B	141	GLY	2.3
4	D	64	GLY	2.3
4	D	141	SER	2.2
1	A	180	GLU	2.1
4	D	202	ASN	2.1
5	E	189	LEU	2.1
2	B	132	PHE	2.0
5	E	197	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	F	2	14/15	0.74	0.14	50,61,71,76	0
6	NAG	F	1	14/15	0.82	0.12	34,44,55,60	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

6.5 Other polymers [i](#)

There are no such residues in this entry.