



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 4, 2025 – 10:10 AM EDT

PDB ID : 9EJS / pdb_00009ejs
Title : Bruton's tyrosine kinase with mutations in the activation loop in complex with compound PTI52
Authors : Lin, D.Y.; Andreotti, A.H.; Tonge, P.J.; Bravo, E.; Li, X.
Deposited on : 2024-11-28
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>.</div> </div> </div>
1	B	271	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>.</div> </div> </div>
1	C	271	<div> <div></div> <div> <div></div> <div>90%</div> <div>7%</div> <div>.</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13154 atoms, of which 6354 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase BTK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	265	Total	C	H	N	O	S	0	1	0
			4256	1373	2104	356	402	21			
1	B	263	Total	C	H	N	O	S	0	1	0
			4190	1356	2069	351	394	20			
1	C	265	Total	C	H	N	O	S	0	0	0
			4215	1360	2085	353	396	21			

There are 45 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	395	MET	-	initiating methionine	UNP P35991
A	430	ARG	LYS	engineered mutation	UNP P35991
A	542	MET	LEU	engineered mutation	UNP P35991
A	543	THR	SER	engineered mutation	UNP P35991
A	555	THR	VAL	engineered mutation	UNP P35991
A	562	LYS	ARG	engineered mutation	UNP P35991
A	564	ALA	SER	engineered mutation	UNP P35991
A	565	SER	PRO	engineered mutation	UNP P35991
A	617	PRO	TYR	engineered mutation	UNP P35991
A	660	HIS	-	expression tag	UNP P35991
A	661	HIS	-	expression tag	UNP P35991
A	662	HIS	-	expression tag	UNP P35991
A	663	HIS	-	expression tag	UNP P35991
A	664	HIS	-	expression tag	UNP P35991
A	665	HIS	-	expression tag	UNP P35991
B	395	MET	-	initiating methionine	UNP P35991
B	430	ARG	LYS	engineered mutation	UNP P35991
B	542	MET	LEU	engineered mutation	UNP P35991
B	543	THR	SER	engineered mutation	UNP P35991
B	555	THR	VAL	engineered mutation	UNP P35991
B	562	LYS	ARG	engineered mutation	UNP P35991
B	564	ALA	SER	engineered mutation	UNP P35991
B	565	SER	PRO	engineered mutation	UNP P35991

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	617	PRO	TYR	engineered mutation	UNP P35991
B	660	HIS	-	expression tag	UNP P35991
B	661	HIS	-	expression tag	UNP P35991
B	662	HIS	-	expression tag	UNP P35991
B	663	HIS	-	expression tag	UNP P35991
B	664	HIS	-	expression tag	UNP P35991
B	665	HIS	-	expression tag	UNP P35991
C	395	MET	-	initiating methionine	UNP P35991
C	430	ARG	LYS	engineered mutation	UNP P35991
C	542	MET	LEU	engineered mutation	UNP P35991
C	543	THR	SER	engineered mutation	UNP P35991
C	555	THR	VAL	engineered mutation	UNP P35991
C	562	LYS	ARG	engineered mutation	UNP P35991
C	564	ALA	SER	engineered mutation	UNP P35991
C	565	SER	PRO	engineered mutation	UNP P35991
C	617	PRO	TYR	engineered mutation	UNP P35991
C	660	HIS	-	expression tag	UNP P35991
C	661	HIS	-	expression tag	UNP P35991
C	662	HIS	-	expression tag	UNP P35991
C	663	HIS	-	expression tag	UNP P35991
C	664	HIS	-	expression tag	UNP P35991
C	665	HIS	-	expression tag	UNP P35991

- Molecule 2 is 4-{4-amino-1-[(3R)-1-(cyclopropanecarbonyl)piperidin-3-yl]-1H-pyrazolo[3,4-d]pyrimidin-3-yl}-N-(4-cyclopropylpyridin-2-yl)benzamide (CCD ID: A1BI0) (formula: C₂₉H₃₀N₈O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			69	29	30	8	2		
2	B	1	Total	C	H	N	O	0	0
			69	29	30	8	2		
2	C	1	Total	C	H	N	O	0	0
			69	29	30	8	2		

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	H	O	S	0	0
			10	2	6	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	98	Total	O	0	0
			98	98		
4	B	69	Total	O	0	0
			69	69		
4	C	109	Total	O	0	0
			109	109		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tyrosine-protein kinase BTK



- Molecule 1: Tyrosine-protein kinase BTK



- Molecule 1: Tyrosine-protein kinase BTK



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	106.63Å 44.13Å 108.00Å 90.00° 117.61° 90.00°	Depositor
Resolution (Å)	19.99 – 2.26 19.99 – 2.26	Depositor EDS
% Data completeness (in resolution range)	65.2 (19.99-2.26) 59.5 (19.99-2.26)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.27Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.191 , 0.246 0.191 , 0.247	Depositor DCC
R_{free} test set	1309 reflections (3.06%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.038 for -h-l,k,h 0.038 for l,k,-h-l 0.033 for h,-k,-h-l 0.027 for -h-l,-k,l 0.028 for l,-k,h	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13154	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1BI0, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/2202	0.29	0/2967
1	B	0.11	0/2170	0.26	0/2925
1	C	0.11	0/2176	0.26	0/2932
All	All	0.11	0/6548	0.27	0/8824

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2152	2104	2105	13	0
1	B	2121	2069	2070	19	0
1	C	2130	2085	2084	13	0
2	A	39	30	0	0	0
2	B	39	30	0	0	0
2	C	39	30	0	0	0
3	C	4	6	6	0	0
4	A	98	0	0	2	2
4	B	69	0	0	9	0
4	C	109	0	0	6	0
All	All	6800	6354	6265	45	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:618:ARG:NH1	4:C:801:HOH:O	2.08	0.87
1:B:655:MET:SD	4:B:866:HOH:O	2.37	0.81
1:B:618:ARG:NH1	1:B:622:ALA:O	2.13	0.81
1:A:437:MET:HE1	1:A:546:VAL:HG22	1.63	0.78
1:B:608:GLU:OE1	4:B:801:HOH:O	2.01	0.76
1:C:635:HIS:ND1	4:C:804:HOH:O	2.23	0.71
1:C:433:ARG:NH2	4:C:807:HOH:O	2.25	0.70
1:C:520:ARG:NH1	4:C:805:HOH:O	2.24	0.69
1:C:412:GLN:OE1	4:C:802:HOH:O	2.11	0.69
1:B:497:GLN:OE1	4:B:802:HOH:O	2.13	0.66
1:C:580:ILE:HD13	1:C:642:PRO:O	2.02	0.60
1:C:396:GLU:OE2	1:C:465:THR:OG1	2.15	0.59
1:A:642:PRO:HG2	1:A:647:LEU:HD13	1.84	0.58
1:C:504:ASP:OD2	4:C:803:HOH:O	2.18	0.57
1:B:492:ARG:O	4:B:803:HOH:O	2.17	0.56
1:A:521:ASP:O	1:A:526:ASN:ND2	2.38	0.55
1:B:435:GLY:O	1:B:547:LEU:HD12	2.07	0.55
1:B:437:MET:HE3	1:B:545:TYR:HB2	1.89	0.55
1:A:437:MET:CE	1:A:546:VAL:HG22	2.36	0.53
1:A:618:ARG:NH2	4:A:809:HOH:O	2.42	0.53
1:B:455:GLU:OE1	4:B:804:HOH:O	2.18	0.51
1:B:575:SER:O	1:B:578:SER:OG	2.28	0.49
1:B:396:GLU:OE1	4:B:805:HOH:O	2.20	0.49
1:A:629:ILE:HD11	1:A:654:VAL:HG21	1.93	0.48
1:B:444:GLU:OE1	4:B:806:HOH:O	2.20	0.48
1:B:642:PRO:HG2	1:B:647:LEU:HD13	1.95	0.48
1:B:515:LYS:NZ	4:B:817:HOH:O	2.47	0.47
1:B:529:VAL:HG22	1:B:535:VAL:HG22	1.97	0.47
1:A:482:LEU:CD1	1:A:501:MET:HE2	2.44	0.46
1:C:586:LEU:O	1:C:586:LEU:HD23	2.17	0.44
1:A:599:GLU:OE1	4:A:801:HOH:O	2.21	0.44
1:A:481:CYS:SG	1:A:525:ARG:HA	2.58	0.44
1:A:577:LYS:HA	1:A:580:ILE:HD12	1.99	0.44
1:B:457:LEU:HD22	1:B:540:PHE:CE2	2.53	0.43
1:C:586:LEU:HD23	1:C:586:LEU:C	2.43	0.43
1:B:494:GLN:N	4:B:802:HOH:O	2.36	0.42
1:C:499:LEU:HD13	1:C:591:TYR:OH	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:525:ARG:NH2	1:B:560:PRO:HB3	2.35	0.42
1:B:608:GLU:O	1:B:612:GLN:HG2	2.19	0.42
1:C:489:MET:HE2	1:C:493:PHE:CE1	2.55	0.42
1:A:483:LEU:HD22	1:A:487:ARG:CZ	2.50	0.41
1:B:609:HIS:HB3	1:B:614:LEU:HD23	2.02	0.41
1:C:519:HIS:O	1:C:520:ARG:HB2	2.21	0.41
1:A:439:GLU:OE1	1:A:468:ARG:NH1	2.53	0.40
1:A:445:GLU:HG3	1:A:449:MET:HE2	2.02	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:887:HOH:O	4:A:893:HOH:O[1_565]	2.06	0.14
4:A:879:HOH:O	4:A:897:HOH:O[1_565]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/271 (97%)	259 (98%)	5 (2%)	0	100	100
1	B	260/271 (96%)	257 (99%)	3 (1%)	0	100	100
1	C	263/271 (97%)	253 (96%)	10 (4%)	0	100	100
All	All	787/813 (97%)	769 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/245 (96%)	231 (98%)	4 (2%)	56	66
1	B	230/245 (94%)	229 (100%)	1 (0%)	89	92
1	C	231/245 (94%)	227 (98%)	4 (2%)	56	66
All	All	696/735 (95%)	687 (99%)	9 (1%)	65	74

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	483	LEU
1	A	492	ARG
1	A	510	GLU
1	A	655	MET
1	B	483	LEU
1	C	483	LEU
1	C	544	ARG
1	C	600	ARG
1	C	655	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	620	HIS
1	C	484	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1BI0	C	701	-	43,45,45	2.27	8 (18%)	54,66,66	1.94	10 (18%)
3	DMS	C	702	-	3,3,3	0.62	0	3,3,3	0.28	0
2	A1BI0	B	701	-	43,45,45	2.25	8 (18%)	54,66,66	1.86	11 (20%)
2	A1BI0	A	701	-	43,45,45	2.25	7 (16%)	54,66,66	1.93	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1BI0	C	701	-	-	6/28/42/42	0/7/7/7
2	A1BI0	B	701	-	-	9/28/42/42	0/7/7/7
2	A1BI0	A	701	-	-	10/28/42/42	0/7/7/7

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	701	A1BI0	C33-N31	9.28	1.48	1.35
2	B	701	A1BI0	C33-N31	9.23	1.48	1.35
2	A	701	A1BI0	C33-N31	9.17	1.48	1.35
2	A	701	A1BI0	N21-N20	-5.29	1.31	1.37
2	C	701	A1BI0	N21-N20	-5.24	1.31	1.37
2	B	701	A1BI0	N21-N20	-5.15	1.31	1.37
2	A	701	A1BI0	C9-N8	4.03	1.47	1.35
2	B	701	A1BI0	C9-N8	4.02	1.47	1.35
2	C	701	A1BI0	C9-N8	4.01	1.47	1.35
2	B	701	A1BI0	C22-N27	3.51	1.46	1.34
2	A	701	A1BI0	C22-N27	3.48	1.46	1.34
2	C	701	A1BI0	C22-N27	3.48	1.46	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	A1BI0	C14-C17	3.45	1.54	1.49
2	A	701	A1BI0	C14-C17	3.35	1.54	1.49
2	C	701	A1BI0	C14-C17	3.29	1.54	1.49
2	A	701	A1BI0	C2-N8	2.99	1.47	1.40
2	C	701	A1BI0	C2-N8	2.96	1.47	1.40
2	B	701	A1BI0	C2-N8	2.95	1.47	1.40
2	C	701	A1BI0	C17-N21	-2.59	1.33	1.35
2	A	701	A1BI0	C17-N21	-2.47	1.33	1.35
2	B	701	A1BI0	C17-N21	-2.25	1.33	1.35
2	C	701	A1BI0	C34-C33	2.11	1.54	1.51
2	B	701	A1BI0	C1-C6	-2.08	1.36	1.39

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	701	A1BI0	C39-C7-C6	7.95	132.14	121.28
2	B	701	A1BI0	C39-C7-C6	7.52	131.56	121.28
2	A	701	A1BI0	C39-C7-C6	7.31	131.27	121.28
2	A	701	A1BI0	C38-C7-C6	6.93	130.74	121.28
2	C	701	A1BI0	C38-C7-C6	6.86	130.65	121.28
2	B	701	A1BI0	C38-C7-C6	6.39	130.02	121.28
2	A	701	A1BI0	C5-C4-N3	-3.47	119.72	123.97
2	B	701	A1BI0	C5-C4-N3	-3.39	119.82	123.97
2	A	701	A1BI0	C34-C33-N31	3.28	121.89	118.80
2	C	701	A1BI0	C5-C4-N3	-3.17	120.09	123.97
2	A	701	A1BI0	C30-N31-C32	2.88	118.62	113.07
2	A	701	A1BI0	C17-N21-N20	2.73	107.35	105.17
2	C	701	A1BI0	C34-C33-N31	2.73	121.37	118.80
2	B	701	A1BI0	C30-N31-C32	2.67	118.22	113.07
2	C	701	A1BI0	C14-C17-N21	2.50	125.06	120.78
2	A	701	A1BI0	C1-C2-N3	-2.37	119.81	122.92
2	B	701	A1BI0	C34-C33-N31	2.33	121.00	118.80
2	C	701	A1BI0	C1-C2-N3	-2.28	119.93	122.92
2	A	701	A1BI0	C4-N3-C2	2.27	120.45	117.21
2	C	701	A1BI0	C30-N31-C32	2.26	117.42	113.07
2	B	701	A1BI0	C1-C2-N3	-2.25	119.97	122.92
2	B	701	A1BI0	C14-C17-N21	2.24	124.62	120.78
2	B	701	A1BI0	C2-N8-C9	-2.22	121.96	128.21
2	C	701	A1BI0	C36-C34-C33	-2.17	114.12	116.91
2	B	701	A1BI0	C17-N21-N20	2.09	106.84	105.17
2	C	701	A1BI0	C15-C14-C13	2.08	121.40	117.68
2	B	701	A1BI0	C4-N3-C2	2.07	120.17	117.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	A1BI0	C28-C26-C32	-2.07	108.32	110.83
2	C	701	A1BI0	C17-N21-N20	2.03	106.79	105.17
2	A	701	A1BI0	O35-C33-N31	-2.02	119.20	121.61

There are no chirality outliers.

All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	A1BI0	C13-C14-C17-N21
2	A	701	A1BI0	C15-C14-C17-N21
2	A	701	A1BI0	C28-C26-N20-C19
2	A	701	A1BI0	C28-C26-N20-N21
2	A	701	A1BI0	N31-C33-C34-C36
2	A	701	A1BI0	N31-C33-C34-C37
2	A	701	A1BI0	O35-C33-C34-C36
2	A	701	A1BI0	O35-C33-C34-C37
2	B	701	A1BI0	C13-C14-C17-N21
2	B	701	A1BI0	C15-C14-C17-N21
2	B	701	A1BI0	C28-C26-N20-C19
2	B	701	A1BI0	C28-C26-N20-N21
2	C	701	A1BI0	C13-C14-C17-N21
2	C	701	A1BI0	C15-C14-C17-N21
2	C	701	A1BI0	C28-C26-N20-C19
2	C	701	A1BI0	C28-C26-N20-N21
2	A	701	A1BI0	C13-C14-C17-C18
2	B	701	A1BI0	O35-C33-C34-C36
2	C	701	A1BI0	O35-C33-C34-C36
2	B	701	A1BI0	N31-C33-C34-C36
2	C	701	A1BI0	N31-C33-C34-C36
2	A	701	A1BI0	C15-C14-C17-C18
2	B	701	A1BI0	C5-C6-C7-C38
2	B	701	A1BI0	C1-C6-C7-C38
2	B	701	A1BI0	C13-C14-C17-C18

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand A1BI0 B 701



Ligand A1BI0 A 701



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/271 (97%)	-0.30	2 (0%) 82 83	20, 34, 62, 96	1 (0%)
1	B	263/271 (97%)	-0.13	3 (1%) 77 79	19, 39, 66, 102	1 (0%)
1	C	265/271 (97%)	-0.09	1 (0%) 89 90	24, 41, 64, 85	0
All	All	793/813 (97%)	-0.18	6 (0%) 82 83	19, 38, 66, 102	2 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	467	GLN	2.8
1	B	600	ARG	2.8
1	B	655	MET	2.5
1	C	395	MET	2.3
1	B	489	MET	2.2
1	A	489	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	DMS	C	702	4/4	0.69	0.12	33,40,51,80	0
2	A1BI0	C	701	39/39	0.92	0.09	23,36,57,57	0
2	A1BI0	A	701	39/39	0.92	0.08	21,33,50,57	0
2	A1BI0	B	701	39/39	0.94	0.09	18,29,52,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around A1BI0 A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1BI0 B 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.