



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2024 – 06:58 AM EDT

PDB ID : 1EK1
Title : CRYSTAL STRUCTURE OF MURINE SOLUBLE EPOXIDE HYDROLASE
COMPLEXED WITH CIU INHIBITOR
Authors : Argiriadi, M.A.; Morisseau, C.; Goodrow, M.H.; Dowdy, D.L.; Hammock,
B.D.; Christianson, D.W.
Deposited on : 2000-03-06
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	554	 17% 46% 23% • 12%
1	B	554	 19% 49% 27% • •

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8255 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EPOXIDE HYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	61	0	0
			3901	2517	651	704	29			
1	B	541	Total	C	N	O	S	71	0	0
			4299	2766	719	783	31			

- Molecule 2 is N-CYCLOHEXYL-N'-(4-IODOPHENYL)UREA (three-letter code: CIU) (formula: C₁₃H₁₇IN₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	I	N	O	0	0
			17	13	1	2	1		
2	B	1	Total	C	I	N	O	0	0
			17	13	1	2	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total 4	O 4	0	0
3	B	17	Total 17	O 17	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: EPOXIDE HYDROLASE



• Molecule 1: EPOXIDE HYDROLASE



P495	P432	P359	A307	P244	D184	T123	Q62
I496	E453	M370	M308	G245	D185	M124	M63
Y497	D434	K371	E309	R246	F186	N125	V64
L498	P435	V372	L310	R247	G187	W126	P65
R499	N436	I373	L311	L248	S188	L127	L66
P500	L437	R374	C312	H249	M189	D128	M67
E501	S438	S375	K313	F250	L190	D129	D68
M502	K439	I376	E314	V251	K191	G130	E69
S503	I440	P377	M315	E252	P192	D131	S70
K504	T441	V378	V316	M253	A193	K132	Y71
M505	T442	F379	T317	G254	R194	K133	R72
M506		N380	F318	S255	D195	D134	K73
E507	E445	Y381	L319	G256	M196	S135	S74
K508	I446	Q382	D320		G197	L136	S75
M509	E447	L383	K321	L259	M198	A137	K76
I510	F448	Y384	L322		P199	Q138	A77
P511	Y449	F385	G323	H263	T200	M139	C78
F512	I450	Q386	I324	G264	I201	M140	G79
L513	Q451	E387	P325	F265	L202	C141	A80
K514	Q452	P388	Q326	P266	V203	E142	M81
M515	F453	G389	A327	E267	H204	L143	L82
O516	K454	V390			N205	Q144	P83
					T206	Q145	E84
C521	T456	L395	I330	F270	A207	H146	N85
G522	G457	E396	G331	S271	S208	F147	F86
H523	F458	K397	H332	W272	A209	D148	S87
M524	R459	N398	D333	R273	L210	F149	I88
T525	Q460	M399	W334	Y274	R211	L150	S89
Q526	P461	S400	A335	Q275	E212	I151	Q90
I527	L462	R401	G336	I276	L213	E152	I91
E528	N463	T402	V337	P277	E214	S153	F92
K529	W464	F403	M338	A278	K215	C154	S93
P530	Y465	K404	V339	L279	V216	Q155	Q94
T531	R466	S405	W340	A280	T217	V156	A95
E532	N467	F406	N341	Q281	G218	G157	M96
V533	T468	F407	M342	A282	T219	M158	
N534	E469	R408	A343	G283	Q220	I159	R99
Q535	R470	A409	L344	F284	F221	K160	S100
I536	N471	S410		R285	P222	P161	I101
L537	W472	D411	P347	V286	E223	E162	M102
I538	K473	E412	E348	L287	A224	P163	R103
K539	W474	T413	R349	A288	P225	Q164	P104
W540	S475	G414	V350	I289	L226	I165	M105
L541	C476	F415	F351	D290	P227	Y166	L106
Q542	K477	I416	A352	N291	V228	M167	Q107
T543	G478	A417	V353	K292	P229	F168	A108
E544	L479	Y418	A354	G293	C230	L169	A109
VAL	G480	H419	S355	Y294	N231	I110	I110
GLN	R481	K420	L356	G295	P232	A111	A111
ASN	K482	A421	N357	D296	N233	L112	L112
PRO	I483	T422	T358	S297	D234	K173	K113
SER	L484	E423	P359	S298	V235	A175	K114
THR		I424	F360	S299	G236	K176	K115
LYS	L488	G425	M361	P300	H237	P177	G116
ILE	M489	G426	P362	P301	G238	N178	F117
	V490	I427	P363	E302	Y239	E179	T118
	T491	L428	D364	I303	V240	V180	T119
	A492	W429		E304	T241	V181	C120
	E493	M430	V367	E305	V242	F182	I121
	K494	T431	S368	Y306	K243	L183	V122

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	151.90Å 143.00Å 60.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.10	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-3.10)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.194 , 0.305	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8255	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CIU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	1/4004 (0.0%)	0.95	11/5430 (0.2%)
1	B	0.77	0/4413	0.96	11/5984 (0.2%)
All	All	0.77	1/8417 (0.0%)	0.96	22/11414 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	230	CYS	CB-SG	5.35	1.91	1.82

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	231	ASN	C-N-CD	-12.52	93.07	120.60
1	B	231	ASN	C-N-CA	7.61	153.97	122.00
1	A	218	GLY	N-CA-C	-6.85	95.98	113.10
1	B	66	LEU	CA-CB-CG	6.57	130.41	115.30
1	A	231	ASN	N-CA-C	5.82	126.72	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3901	0	3890	581	0
1	B	4299	0	4270	676	0
2	A	17	0	17	2	0
2	B	17	0	17	5	0
3	A	4	0	0	0	0
3	B	17	0	0	5	0
All	All	8255	0	8194	1220	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 76.

The worst 5 of 1220 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293:GLY:HA2	1:A:299:SER:HB2	1.33	1.10
1:B:141:CYS:O	1:B:144:SER:HB3	1.52	1.10
1:B:61:SER:O	1:B:64:VAL:HG23	1.51	1.09
1:B:183:LEU:HD23	1:B:201:ILE:HD12	1.31	1.08
1:B:223:GLU:CD	1:B:223:GLU:H	1.53	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	484/554 (87%)	345 (71%)	90 (19%)	49 (10%)	0	3
1	B	539/554 (97%)	384 (71%)	101 (19%)	54 (10%)	0	3
All	All	1023/1108 (92%)	729 (71%)	191 (19%)	103 (10%)	0	3

5 of 103 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	ASP
1	A	18	SER
1	A	63	TRP
1	A	65	PRO
1	A	91	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/480 (89%)	291 (68%)	136 (32%)	0	0
1	B	468/480 (98%)	317 (68%)	151 (32%)	0	0
All	All	895/960 (93%)	608 (68%)	287 (32%)	0	0

5 of 287 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	319	LEU
1	B	539	LYS
1	B	372	VAL
1	B	440	ILE
1	A	408	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	167	ASN
1	B	204	HIS
1	B	189	ASN
1	B	205	ASN
1	A	281	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIU	A	1100	-	18,18,18	2.84	10 (55%)	23,23,23	1.99	4 (17%)
2	CIU	B	1200	-	18,18,18	2.84	10 (55%)	23,23,23	1.99	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIU	A	1100	-	-	0/8/16/16	0/2/2/2
2	CIU	B	1200	-	-	0/8/16/16	0/2/2/2

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1200	CIU	C11-I4	6.54	2.27	2.10
2	A	1100	CIU	C11-I4	6.51	2.27	2.10
2	B	1200	CIU	C4-C3	4.04	1.61	1.52
2	B	1200	CIU	C12-C11	4.03	1.47	1.38
2	A	1100	CIU	C12-C11	4.02	1.47	1.38

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1200	CIU	N2-C7-N1	5.44	121.25	113.77
2	A	1100	CIU	N2-C7-N1	5.43	121.24	113.77
2	B	1200	CIU	C8-N2-C7	4.56	135.92	126.61
2	A	1100	CIU	C8-N2-C7	4.55	135.89	126.61
2	A	1100	CIU	O2-C7-N1	-3.90	114.34	122.67

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1100	CIU	2	0
2	B	1200	CIU	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.