



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 09:53 AM EDT

PDB ID : 6ELL
Title : FAB Fragment. AbVance: Increasing our knowledge of antibody structural space to enable faster and better decision making in antibody drug discovery
Authors : Benz, J.; Weigand, S.; Dengl, S.; Schlothauer, T.; Auer, J.; Ehler, A.; Kettenberger, H.; Lorenz, S.; Hirschheydt, T.; Georges, G.
Deposited on : 2017-09-29
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>85%</div> <div>8%</div> <div>7%</div> </div>
1	H	227	<div> <div>3%</div> <div>85%</div> <div>11%</div> <div>.</div> </div>
2	B	214	<div> <div>2%</div> <div>91%</div> <div>9%</div> </div>
2	L	214	<div> <div>93%</div> <div>6%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called fAB heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1602	1014	267	314	7			
1	H	218	Total	C	N	O	S	0	0	0
			1669	1058	279	325	7			

- Molecule 2 is a protein called fAB light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	213	Total	C	N	O	S	0	0	0
			1645	1030	278	332	5			
2	L	213	Total	C	N	O	S	0	0	0
			1649	1032	278	334	5			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	121	Total	O	0	0
			121	121		
3	B	111	Total	O	0	0
			111	111		
3	H	102	Total	O	0	0
			102	102		
3	L	99	Total	O	0	0
			99	99		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: fAB heavy chain




- Molecule 1: fAB heavy chain



- Molecule 2: fAB light chain



- Molecule 2: fAB light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.98Å 69.17Å 77.67Å 117.11° 89.50° 103.39°	Depositor
Resolution (Å)	38.00 – 1.90 44.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.4 (38.00-1.90) 97.4 (44.40-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 1.89Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.203 , 0.245 0.216 , 0.258	Depositor DCC
R_{free} test set	3598 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6998	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1642	0.66	0/2242
1	H	0.50	0/1714	0.68	0/2341
2	B	0.51	0/1681	0.64	0/2282
2	L	0.51	0/1685	0.66	0/2287
All	All	0.51	0/6722	0.66	0/9152

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1602	0	1556	11	0
1	H	1669	0	1612	12	0
2	B	1645	0	1597	8	0
2	L	1649	0	1601	5	0
3	A	121	0	0	0	0
3	B	111	0	0	0	0
3	H	102	0	0	0	0
3	L	99	0	0	0	0
All	All	6998	0	6366	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:38:ARG:HB3	1:H:48:ILE:HD11	1.86	0.58
2:L:198:HIS:CD2	2:L:200:GLY:H	2.22	0.57
2:L:124:GLN:HE22	2:L:131:SER:H	1.52	0.57
2:B:198:HIS:CD2	2:B:200:GLY:H	2.25	0.55
1:A:38:ARG:HB3	1:A:48:ILE:HD11	1.88	0.55
1:A:130:PRO:HB3	1:A:156:TYR:HB3	1.90	0.54
1:A:30:SER:O	1:A:53:PRO:HB3	2.07	0.54
1:H:166:ASN:ND2	1:H:206:ILE:H	2.07	0.52
1:H:11:LEU:HB2	1:H:158:PRO:HG3	1.90	0.52
1:A:33:TRP:O	1:A:99:PRO:HD2	2.10	0.52
2:B:145:LYS:HB3	2:B:197:THR:HB	1.94	0.48
2:B:124:GLN:HE22	2:B:131:SER:HB2	1.77	0.48
2:L:20:THR:HG23	2:L:72:THR:HG23	1.95	0.48
2:B:20:THR:HG23	2:B:72:THR:HG23	1.96	0.48
1:H:51:VAL:HG23	1:H:58:ILE:HG12	1.96	0.47
2:L:145:LYS:HB3	2:L:197:THR:HB	1.97	0.47
1:H:30:SER:O	1:H:53:PRO:HB3	2.15	0.46
1:H:64:LEU:HB3	1:H:68:PHE:HD2	1.81	0.46
1:H:130:PRO:HB3	1:H:156:TYR:HB3	1.98	0.45
2:L:198:HIS:HD2	2:L:200:GLY:H	1.65	0.45
1:A:18:LEU:HB3	1:A:83:MET:CE	2.46	0.45
1:H:18:LEU:HB3	1:H:83:MET:HE2	1.99	0.44
1:A:83:MET:HB3	1:A:86:LEU:HD21	1.98	0.44
1:H:137:PRO:HG2	1:H:224:PRO:HB3	1.99	0.44
1:A:179:ALA:HA	1:A:189:LEU:HB3	1.99	0.44
2:B:141:PRO:O	2:B:198:HIS:HE1	2.01	0.43
1:A:18:LEU:HB3	1:A:83:MET:HE2	1.99	0.43
2:B:159:SER:HA	2:B:178:THR:O	2.18	0.43
2:B:35:TRP:CE2	2:B:73:PHE:HB2	2.54	0.43
1:A:36:TRP:CE2	1:A:81:LEU:HB2	2.54	0.43
1:A:166:ASN:ND2	1:A:206:ILE:H	2.16	0.43
1:A:155:ASP:HB3	1:A:186:LEU:HD13	2.01	0.42
1:H:155:ASP:HB3	1:H:186:LEU:HD13	2.02	0.42
2:B:83:ILE:HD12	2:B:104:LEU:O	2.21	0.41
1:H:179:ALA:HA	1:H:189:LEU:HB3	2.02	0.40
1:H:34:MET:HB3	1:H:79:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/227 (90%)	203 (99%)	2 (1%)	0	100	100
1	H	214/227 (94%)	213 (100%)	1 (0%)	0	100	100
2	B	211/214 (99%)	204 (97%)	7 (3%)	0	100	100
2	L	211/214 (99%)	205 (97%)	6 (3%)	0	100	100
All	All	841/882 (95%)	825 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/195 (93%)	181 (100%)	0	100	100
1	H	187/195 (96%)	185 (99%)	2 (1%)	73	73
2	B	188/190 (99%)	184 (98%)	4 (2%)	53	48
2	L	189/190 (100%)	184 (97%)	5 (3%)	46	39
All	All	745/770 (97%)	734 (98%)	11 (2%)	65	62

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	10	SER
2	B	11	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	114	SER
2	B	126	LYS
1	H	17	SER
1	H	85	SER
2	L	10	SER
2	L	11	LEU
2	L	12	SER
2	L	48	ILE
2	L	83	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	ASN
2	B	124	GLN
2	B	137	ASN
2	B	198	HIS
2	B	210	ASN
1	H	82	GLN
1	H	84	ASN
1	H	166	ASN
2	L	124	GLN
2	L	138	ASN
2	L	198	HIS
2	L	210	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/227 (92%)	0.11	1 (0%) 91 92	27, 40, 64, 79	0
1	H	218/227 (96%)	0.06	7 (3%) 47 50	27, 39, 60, 83	0
2	B	213/214 (99%)	0.07	4 (1%) 66 69	25, 36, 56, 72	0
2	L	213/214 (99%)	0.04	0 100 100	25, 37, 53, 73	0
All	All	855/882 (96%)	0.07	12 (1%) 75 77	25, 38, 60, 83	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	106	TYR	6.0
1	H	105	ARG	5.7
1	H	138	SER	2.6
1	H	224	PRO	2.5
1	H	204	THR	2.3
2	B	181	LEU	2.2
1	H	103	GLY	2.2
2	B	125	LEU	2.2
2	B	158	ASN	2.2
2	B	156	SER	2.1
1	H	104	SER	2.0
1	A	184	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.