



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 15, 2025 – 07:06 pm BST

PDB ID : 9EWD / pdb\_00009ewd  
Title : DNA Polymerase Lambda I493K E529D, TMP:At Ca<sup>2+</sup> Ground State Ternary Complex  
Authors : Nourisson, A.; Haouz, A.; Missouri, S.; Delarue, M.  
Deposited on : 2024-04-03  
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7689 (2.14-2.10)
Clashscore	180529	8431 (2.14-2.10)
Ramachandran outliers	177936	8366 (2.14-2.10)
Sidechain outliers	177891	8367 (2.14-2.10)
RSRZ outliers	164620	7689 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div> <div>9%</div> <div>82%</div> <div>15%</div> <div>..</div> </div>
2	D	4	<div> <div>50%</div> <div>50%</div> </div>
3	P	6	<div> <div>100%</div> </div>
4	T	11	<div> <div>73%</div> <div>27%</div> </div>

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 3174 atoms, of which 13 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	3	0
			2481	1563	450	457	11			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	241	GLY	-	expression tag	UNP Q9UGP5
A	463	LYS	-	linker	UNP Q9UGP5
A	464	GLY	-	linker	UNP Q9UGP5
A	470	GLU	-	linker	UNP Q9UGP5
A	471	THR	-	linker	UNP Q9UGP5
A	492	LYS	ILE	engineered mutation	UNP Q9UGP5
A	529	ASP	GLU	engineered mutation	UNP Q9UGP5

- Molecule 2 is a DNA chain called DNA primer strand downstream (5'P).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			


- Molecule 3 is a DNA chain called DNA primer strand upstream.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	6	Total	C	N	O	P	0	0	0
			119	58	23	33	5			

- Molecule 4 is a DNA chain called DNA template strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	T	11	Total	C	N	O	P	0	0	0
			224	107	43	64	10			

- Molecule 5 is THYMIDINE-5'-PHOSPHATE (CCD ID: TMP) (formula:  $C_{10}H_{15}N_2O_8P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	P	0	0
			34	10	13	2	8	1		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Na	0	0
			2	2		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	2	Total 2	K 2	0	0

- Molecule 10 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total 1	Ca 1	0	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	150	Total 150	O 150	0	0
11	D	3	Total 3	O 3	0	0
11	P	26	Total 26	O 26	0	0
11	T	31	Total 31	O 31	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase lambda



- Molecule 2: DNA primer strand downstream (5'P)




- Molecule 3: DNA primer strand upstream



There are no outlier residues recorded for this chain.

- Molecule 4: DNA template strand



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.40Å 62.47Å 139.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.89 – 2.12 35.89 – 2.12	Depositor EDS
% Data completeness (in resolution range)	99.8 (35.89-2.12) 99.8 (35.89-2.12)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.57 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.202 , 0.245 0.199 , 0.242	Depositor DCC
$R_{free}$ test set	1472 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.6	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3174	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TMP, K, GOL, CA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/2537	0.46	0/3425
2	D	0.39	0/92	0.50	0/138
3	P	0.53	0/133	0.64	0/203
4	T	0.50	0/251	0.67	0/386
All	All	0.36	0/3013	0.50	0/4152

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2481	0	2464	44	0
2	D	83	0	45	1	0
3	P	119	0	69	0	0
4	T	224	0	125	4	0
5	A	21	13	13	1	0
6	A	2	0	0	0	0
7	A	12	0	18	2	0
8	A	6	0	8	0	0
9	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	A	1	0	0	0	0
11	A	150	0	0	6	0
11	D	3	0	0	0	0
11	P	26	0	0	0	0
11	T	31	0	0	0	0
All	All	3161	13	2742	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ILE:HG21	1:A:313:ILE:HD13	1.34	1.08
1:A:452:ARG:HG3	7:A:604:EDO:H21	1.55	0.87
4:T:8:DA:H2''	4:T:9:DC:H5''	1.67	0.75
1:A:406:PHE:CZ	1:A:446:ARG:HB3	2.25	0.70
5:A:601:TMP:HN3	4:T:5:DA:H61	1.42	0.65
1:A:256:ILE:HG21	1:A:313:ILE:CD1	2.22	0.64
4:T:8:DA:C2'	4:T:9:DC:H5''	2.27	0.64
4:T:8:DA:H2''	4:T:9:DC:C5'	2.29	0.63
1:A:311:GLU:O	1:A:314:ILE:HG12	1.99	0.63
1:A:519:LEU:HD23	1:A:565:LEU:HD11	1.82	0.62
1:A:434:HIS:CD2	1:A:439:SER:HB2	2.35	0.62
1:A:267:TYR:CZ	1:A:275:ARG:HD3	2.36	0.61
1:A:291:LYS:HD3	1:A:298:GLU:OE2	2.02	0.59
1:A:259:LYS:HE2	1:A:320:GLY:O	2.01	0.59
1:A:452:ARG:HG3	7:A:604:EDO:C2	2.32	0.59
1:A:286:LEU:HD11	1:A:313:ILE:HD11	1.86	0.57
1:A:523:LYS:HB2	1:A:525:MET:CE	2.36	0.55
1:A:446:ARG:HD3	11:A:710:HOH:O	2.04	0.55
1:A:556:GLU:CD	11:A:703:HOH:O	2.49	0.54
1:A:350:GLN:HG2	11:A:720:HOH:O	2.09	0.52
1:A:519:LEU:CD2	1:A:565:LEU:HD11	2.39	0.52
1:A:256:ILE:CG2	1:A:313:ILE:HD13	2.24	0.52
1:A:296:TYR:OH	1:A:307:LYS:HG3	2.10	0.51
1:A:406:PHE:CE1	1:A:446:ARG:HB3	2.46	0.51
1:A:264:ALA:HB2	1:A:279:TYR:HB3	1.92	0.49
1:A:544:LYS:HD3	11:A:735:HOH:O	2.14	0.48
1:A:554:PRO:HG2	1:A:558:ASP:OD2	2.15	0.47
1:A:538:ARG:HG2	1:A:544:LYS:HA	1.97	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:HIS:ND1	1:A:435:PRO:HD2	2.31	0.46
1:A:448:LEU:O	1:A:452:ARG:HB2	2.16	0.46
1:A:572:GLU:HA	11:A:729:HOH:O	2.16	0.46
2:D:3:DC:H2''	2:D:4:DG:C8	2.51	0.46
1:A:523:LYS:HB2	1:A:525:MET:HE3	1.98	0.46
1:A:434:HIS:O	1:A:496:TYR:HB2	2.17	0.45
1:A:307:LYS:O	1:A:311:GLU:HG3	2.16	0.45
1:A:526:SER:CB	1:A:536:VAL:HG21	2.47	0.44
1:A:346:THR:O	1:A:350:GLN:HG3	2.17	0.44
1:A:422:LYS:HA	1:A:422:LYS:HD3	1.84	0.44
1:A:435:PRO:HA	1:A:496:TYR:CD1	2.53	0.43
1:A:290:HIS:CE1	1:A:291:LYS:HG2	2.54	0.43
1:A:431:LEU:HD23	1:A:503:LEU:HD13	2.00	0.43
1:A:355:GLN:CD	11:A:702:HOH:O	2.61	0.43
1:A:526:SER:HB2	1:A:536:VAL:HG21	2.01	0.43
1:A:286:LEU:HD21	1:A:305:ILE:CD1	2.49	0.43
1:A:436:ASP:C	1:A:436:ASP:OD1	2.61	0.42
1:A:512:PHE:CE2	1:A:573:ARG:HD3	2.55	0.42
1:A:307:LYS:HE2	1:A:311:GLU:OE2	2.20	0.42
1:A:252:HIS:HB2	1:A:292:PRO:HG3	2.02	0.41
1:A:312:LYS:HB3	1:A:322:LEU:HD11	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/330 (97%)	310 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	258/275 (94%)	255 (99%)	3 (1%)	67 74

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	305	ILE
1	A	313	ILE
1	A	323	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	252	HIS
1	A	255	HIS
1	A	321	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	TMP	A	601	10	22,22,22	0.40	0	33,33,33	0.62	0
7	EDO	A	603	-	3,3,3	0.26	0	2,2,2	0.47	0
8	GOL	A	606	-	5,5,5	0.36	0	5,5,5	0.39	0
7	EDO	A	604	-	3,3,3	0.22	0	2,2,2	0.83	0
7	EDO	A	605	-	3,3,3	0.27	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TMP	A	601	10	-	0/10/22/22	0/2/2/2
7	EDO	A	603	-	-	1/1/1/1	-
8	GOL	A	606	-	-	2/4/4/4	-
7	EDO	A	604	-	-	1/1/1/1	-
7	EDO	A	605	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	A	605	EDO	O1-C1-C2-O2
8	A	606	GOL	O1-C1-C2-O2
7	A	603	EDO	O1-C1-C2-O2
7	A	604	EDO	O1-C1-C2-O2
8	A	606	GOL	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	601	TMP	1	0
7	A	604	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	320/330 (96%)	0.51	29 (9%) 16 18	19, 48, 92, 107	3 (0%)
2	D	4/4 (100%)	-0.10	0 100 100	51, 51, 51, 54	0
3	P	6/6 (100%)	-1.02	0 100 100	29, 32, 35, 41	0
4	T	11/11 (100%)	-0.52	0 100 100	37, 39, 52, 53	0
All	All	341/351 (97%)	0.44	29 (8%) 18 20	19, 47, 90, 107	3 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	540	THR	4.3
1	A	545	VAL	4.0
1	A	537	VAL	3.9
1	A	575	TRP	3.8
1	A	541	HIS	3.7
1	A	289	PHE	3.5
1	A	256	ILE	3.4
1	A	535	ALA	3.4
1	A	293	VAL	3.2
1	A	286	LEU	3.0
1	A	406	PHE	3.0
1	A	290	HIS	2.9
1	A	255	HIS	2.8
1	A	443	ILE	2.8
1	A	511	HIS	2.8
1	A	302	ILE	2.7
1	A	310	ALA	2.6
1	A	296	TYR	2.6
1	A	321	HIS	2.6
1	A	317	LEU	2.6
1	A	299	ALA	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	313	ILE	2.4
1	A	300	CYS	2.3
1	A	316	ILE	2.3
1	A	314	ILE	2.1
1	A	547	PRO	2.1
1	A	322	LEU	2.1
1	A	254	LEU	2.0
1	A	409	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	TMP	A	601	21/21	0.60	0.21	44,70,85,96	0
7	EDO	A	605	4/4	0.62	0.31	45,46,48,53	4
8	GOL	A	606	6/6	0.83	0.17	44,45,45,52	6
10	CA	A	609	1/1	0.84	0.11	76,76,76,76	0
7	EDO	A	604	4/4	0.87	0.10	48,48,51,56	0
6	NA	A	610	1/1	0.92	0.19	62,62,62,62	0
7	EDO	A	603	4/4	0.93	0.18	36,37,38,41	4
9	K	A	608	1/1	0.94	0.22	72,72,72,72	0
9	K	A	607	1/1	0.94	0.07	76,76,76,76	0
6	NA	A	602	1/1	0.98	0.04	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around TMP A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.