



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2025 – 02:50 PM EDT

PDB ID : 4F5M
Title : Wild-Type E. coli Aspartate Aminotransferase: A Template For The Inter-conversion of Substrate Specificity and Activity To Tyrosine Aminotransferase By The JANUS Algorithm.
Authors : Addington, T.A.; Fisher, A.J.; Toney, M.D.
Deposited on : 2012-05-13
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	<div> <div>3%</div> <div>91%</div> <div>7%</div> <div>..</div> </div>
1	B	406	<div> <div>6%</div> <div>85%</div> <div>11%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	501	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate aminotransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	400	Total	C	N	O	P	S	0	0	0
			3105	1957	541	594	1	12			
1	B	396	Total	C	N	O	P	S	0	0	0
			3074	1938	534	589	1	12			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP P00509
A	2	ALA	-	expression tag	UNP P00509
A	3	HIS	-	expression tag	UNP P00509
A	4	HIS	-	expression tag	UNP P00509
A	5	HIS	-	expression tag	UNP P00509
A	6	HIS	-	expression tag	UNP P00509
A	7	HIS	-	expression tag	UNP P00509
A	8	HIS	-	expression tag	UNP P00509
A	9	VAL	-	expression tag	UNP P00509
A	10	GLY	-	expression tag	UNP P00509
A	11	THR	-	expression tag	UNP P00509
B	1	MET	-	expression tag	UNP P00509
B	2	ALA	-	expression tag	UNP P00509
B	3	HIS	-	expression tag	UNP P00509
B	4	HIS	-	expression tag	UNP P00509
B	5	HIS	-	expression tag	UNP P00509
B	6	HIS	-	expression tag	UNP P00509
B	7	HIS	-	expression tag	UNP P00509
B	8	HIS	-	expression tag	UNP P00509
B	9	VAL	-	expression tag	UNP P00509
B	10	GLY	-	expression tag	UNP P00509
B	11	THR	-	expression tag	UNP P00509

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	419	Total	O	0	0
			419	419		
3	B	368	Total	O	0	0
			368	368		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartate aminotransferase



- Molecule 1: Aspartate aminotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.30Å 108.94Å 144.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.09 – 1.65 37.09 – 1.65	Depositor EDS
% Data completeness (in resolution range)	97.3 (37.09-1.65) 97.3 (37.09-1.65)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.176 , 0.210 0.172 , 0.203	Depositor DCC
R_{free} test set	5418 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtriage
Anisotropy	(Not available)	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.5	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6982	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/3143	0.63	0/4261
1	B	0.53	0/3110	0.65	0/4215
All	All	0.51	0/6253	0.64	0/8476

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3105	0	3024	26	0
1	B	3074	0	3004	35	0
2	A	4	0	6	4	0
2	B	12	0	18	3	0
3	A	419	0	0	4	0
3	B	368	0	0	6	0
All	All	6982	0	6052	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:LLP:CA	1:A:256:LLP:CB	1.75	1.63
1:B:83:GLU:OE2	1:B:86:ARG:NH2	1.93	0.99
1:B:295:ASN:HD21	2:B:503:EDO:H11	1.37	0.89
1:B:142:ASN:ND2	3:B:939:HOH:O	2.09	0.83
1:A:256:LLP:CA	1:A:256:LLP:CG	2.57	0.82
1:B:236:GLU:OE2	3:B:965:HOH:O	2.04	0.73
1:B:333:GLN:HA	1:B:352:ILE:HD11	1.69	0.73
1:B:139:SER:HA	1:B:194:PRO:HD3	1.71	0.73
1:B:142:ASN:OD1	1:B:145:SER:OG	2.06	0.73
1:B:344:ALA:O	1:B:345:ASN:HB2	1.89	0.73
1:B:294:SER:OG	2:B:503:EDO:H12	1.90	0.72
1:B:295:ASN:ND2	2:B:503:EDO:H11	2.04	0.72
1:A:212:VAL:HG21	1:A:244:MET:HG2	1.75	0.69
1:B:118:ARG:NH2	3:B:765:HOH:O	2.25	0.67
1:B:370:ARG:NE	1:B:374:GLU:OE1	2.26	0.67
1:B:336:VAL:HG21	1:B:352:ILE:HD13	1.75	0.66
1:B:95:LYS:NZ	3:B:738:HOH:O	2.32	0.62
1:B:396:ALA:HB3	1:B:397:PRO:HD3	1.83	0.59
1:B:344:ALA:HA	1:B:403:VAL:HG22	1.87	0.56
1:A:164:ASN:O	1:A:166:THR:HG23	2.06	0.55
1:B:339:LEU:HD13	1:B:399:CYS:SG	2.47	0.55
1:A:244:MET:HE3	3:A:728:HOH:O	2.06	0.54
1:B:138:PRO:O	1:B:139:SER:HB3	2.08	0.54
1:A:157:TYR:CD1	1:A:172:LEU:HD13	2.43	0.54
1:B:344:ALA:O	1:B:345:ASN:CB	2.56	0.53
1:A:118:ARG:NH2	3:A:981:HOH:O	2.41	0.53
1:B:140:TRP:HB2	1:B:256:LLP:H2'3	1.92	0.52
1:A:172:LEU:C	1:A:172:LEU:HD12	2.31	0.52
1:A:172:LEU:HD12	1:A:172:LEU:O	2.09	0.52
1:B:157:TYR:CD1	1:B:172:LEU:HD13	2.47	0.50
1:A:193:ASN:HD21	2:A:501:EDO:H12	1.75	0.50
1:A:193:ASN:ND2	2:A:501:EDO:H12	2.27	0.49
1:B:46:TYR:CZ	1:B:324:MET:HG2	2.48	0.49
1:B:219:LEU:C	1:B:219:LEU:HD23	2.33	0.49
1:B:164:ASN:O	1:B:166:THR:HG23	2.12	0.48
1:A:83:GLU:HG2	3:A:952:HOH:O	2.14	0.48
1:A:272:ALA:HB3	1:A:278:VAL:HB	1.96	0.47
1:A:9:VAL:HG11	3:A:794:HOH:O	2.15	0.47
1:B:161:ASP:OD1	1:B:163:GLU:HG2	2.14	0.46
1:B:249:ILE:HG12	1:B:270:LEU:HD12	1.97	0.46
1:B:343:GLY:HA2	3:B:920:HOH:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:TRP:HE1	2:A:501:EDO:C1	2.30	0.45
1:A:167:LEU:HB2	1:A:197:ILE:HG21	1.99	0.45
1:B:343:GLY:O	1:B:344:ALA:C	2.54	0.44
1:B:30:PHE:CZ	1:B:40:ASN:HB2	2.52	0.44
1:A:297:PRO:HA	1:B:264:ARG:HG2	2.00	0.43
1:A:194:PRO:HG2	1:A:360:PHE:CE2	2.53	0.43
1:B:380:VAL:HG12	1:B:381:ALA:N	2.34	0.42
1:B:46:TYR:HB2	1:B:357:MET:CE	2.49	0.42
1:A:157:TYR:HD1	1:A:172:LEU:HD13	1.84	0.42
1:A:270:LEU:HD11	1:A:285:MET:HE3	2.01	0.42
1:B:228:ALA:HB3	1:B:234:ASP:OD1	2.20	0.42
1:B:383:GLY:HA3	3:B:732:HOH:O	2.20	0.41
1:A:396:ALA:O	1:A:400:GLU:HG3	2.20	0.41
1:A:173:ILE:HD12	1:A:210:LEU:HD22	2.03	0.41
1:A:256:LLP:CA	1:A:256:LLP:HG2	2.47	0.41
1:B:211:SER:HG	1:B:245:HIS:CE1	2.32	0.41
1:A:140:TRP:HE1	2:A:501:EDO:H11	1.86	0.41
1:A:49:GLU:H	1:A:49:GLU:CD	2.24	0.41
1:A:328:ILE:HG23	1:A:387:VAL:CG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/406 (98%)	389 (98%)	8 (2%)	0	100	100
1	B	393/406 (97%)	381 (97%)	10 (2%)	2 (0%)	25	11
All	All	790/812 (97%)	770 (98%)	18 (2%)	2 (0%)	37	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	345	ASN
1	B	344	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/327 (98%)	314 (98%)	6 (2%)	52	30
1	B	317/327 (97%)	311 (98%)	6 (2%)	52	30
All	All	637/654 (97%)	625 (98%)	12 (2%)	52	30

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LEU
1	A	91	LEU
1	A	167	LEU
1	A	172	LEU
1	A	222	PHE
1	A	313	LEU
1	B	29	LEU
1	B	152	LEU
1	B	172	LEU
1	B	222	PHE
1	B	270	LEU
1	B	339	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	295	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	256	1	23,24,25	3.79	8 (34%)	25,32,34	2.07	3 (12%)
1	LLP	B	256	1	23,24,25	3.48	9 (39%)	25,32,34	1.94	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	256	1	-	7/16/17/19	0/1/1/1
1	LLP	B	256	1	-	8/16/17/19	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	256	LLP	CB-CA	14.56	1.75	1.53
1	B	256	LLP	CB-CA	13.31	1.73	1.53
1	B	256	LLP	C4'-NZ	5.67	1.46	1.27
1	A	256	LLP	C4'-NZ	5.39	1.45	1.27
1	A	256	LLP	CA-N	4.76	1.61	1.48
1	A	256	LLP	C3-C2	4.68	1.45	1.41
1	B	256	LLP	CA-N	4.50	1.60	1.48
1	A	256	LLP	C6-N1	3.45	1.41	1.34
1	B	256	LLP	C3-C2	3.03	1.44	1.41
1	A	256	LLP	C6-C5	2.81	1.43	1.37
1	B	256	LLP	OP4-C5'	-2.74	1.34	1.44
1	B	256	LLP	C6-C5	2.48	1.42	1.37
1	B	256	LLP	C6-N1	2.45	1.39	1.34
1	A	256	LLP	C4-C3	2.40	1.45	1.41
1	B	256	LLP	C4-C4'	2.18	1.51	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	256	LLP	OP4-C5'	-2.10	1.37	1.44
1	B	256	LLP	C4-C3	2.05	1.44	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	256	LLP	OP4-C5'-C5	7.59	123.58	109.36
1	B	256	LLP	OP4-C5'-C5	6.11	120.81	109.36
1	A	256	LLP	CE-NZ-C4'	-4.51	104.26	118.72
1	B	256	LLP	CE-NZ-C4'	-3.91	106.20	118.72
1	A	256	LLP	CG-CD-CE	-3.17	102.34	113.38
1	B	256	LLP	CG-CD-CE	-2.95	103.11	113.38
1	B	256	LLP	C5'-C5-C6	-2.44	115.38	119.36
1	B	256	LLP	OP2-P-OP1	-2.26	102.05	110.83

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	256	LLP	C4-C5-C5'-OP4
1	A	256	LLP	C6-C5-C5'-OP4
1	A	256	LLP	N-CA-CB-CG
1	B	256	LLP	C4-C5-C5'-OP4
1	B	256	LLP	C6-C5-C5'-OP4
1	B	256	LLP	O-C-CA-CB
1	B	256	LLP	CG-CD-CE-NZ
1	A	256	LLP	C4-C4'-NZ-CE
1	A	256	LLP	CG-CD-CE-NZ
1	B	256	LLP	CA-CB-CG-CD
1	A	256	LLP	CA-CB-CG-CD
1	B	256	LLP	C4-C4'-NZ-CE
1	A	256	LLP	C3-C4-C4'-NZ
1	B	256	LLP	C3-C4-C4'-NZ
1	B	256	LLP	N-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	256	LLP	3	0
1	B	256	LLP	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	501	-	3,3,3	0.66	0	2,2,2	0.15	0
2	EDO	B	503	-	3,3,3	0.52	0	2,2,2	0.45	0
2	EDO	B	502	-	3,3,3	0.43	0	2,2,2	0.45	0
2	EDO	A	501	-	3,3,3	0.50	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	501	-	-	0/1/1/1	-
2	EDO	B	503	-	-	1/1/1/1	-
2	EDO	B	502	-	-	0/1/1/1	-
2	EDO	A	501	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	EDO	O1-C1-C2-O2
2	B	503	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	503	EDO	3	0
2	A	501	EDO	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/406 (98%)	0.08	12 (3%) 52 56	16, 30, 53, 85	0
1	B	395/406 (97%)	0.26	23 (5%) 30 32	14, 29, 67, 114	0
All	All	794/812 (97%)	0.17	35 (4%) 39 43	14, 29, 62, 114	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	96	GLY	5.6
1	B	139	SER	4.4
1	B	140	TRP	4.3
1	B	375	PHE	3.5
1	B	141	PRO	3.4
1	A	244	MET	3.3
1	B	370	ARG	3.0
1	A	142	ASN	2.9
1	B	403	VAL	2.8
1	B	369	LEU	2.8
1	B	344	ALA	2.7
1	A	24	LEU	2.7
1	A	211	SER	2.6
1	B	404	ALA	2.6
1	B	405	VAL	2.5
1	B	401	ALA	2.5
1	B	368	VAL	2.5
1	A	23	ILE	2.4
1	A	216	TRP	2.4
1	B	352	ILE	2.4
1	B	37	GLY	2.3
1	B	346	ARG	2.3
1	A	140	TRP	2.3
1	B	349	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	222	PHE	2.3
1	B	366	GLU	2.3
1	B	36	PRO	2.2
1	A	273	ALA	2.2
1	B	24	LEU	2.2
1	B	378	TYR	2.2
1	A	95	LYS	2.1
1	A	272	ALA	2.1
1	A	177	ASN	2.1
1	B	365	LYS	2.0
1	B	32	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	256	24/25	0.97	0.08	17,24,34,40	0
1	LLP	B	256	24/25	0.97	0.08	17,23,36,40	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	501	4/4	0.86	0.12	39,40,44,56	0
2	EDO	B	503	4/4	0.89	0.11	19,25,38,46	0
2	EDO	B	502	4/4	0.91	0.13	24,37,48,49	0
2	EDO	B	501	4/4	0.95	0.10	19,20,23,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.