



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 25, 2024 – 04:08 PM EST

PDB ID : 2F7Y  
Title : Crystal structure of Molybdenum cofactor biosynthesis protein Mog from *Shewanella oneidensis*  
Authors : Chang, C.; Mulligan, R.; Collart, F.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2005-12-01  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	177	<div> <div>14%</div> <div>82%</div> <div>11%</div> <div>• • •</div> </div>
1	B	177	<div> <div>16%</div> <div>69%</div> <div>20%</div> <div>5% • 5%</div> </div>
1	C	177	<div> <div>14%</div> <div>83%</div> <div>10%</div> <div>• •</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4026 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called molybdenum cofactor biosynthesis protein Mog.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	173	Total	C	N	O	S	Se	0	0	0
			1316	832	214	259	6	5			
1	B	168	Total	C	N	O	S	Se	0	0	0
			1286	814	208	253	6	5			
1	C	171	Total	C	N	O	S	Se	0	0	0
			1302	823	211	257	6	5			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	GB 24371665
A	61	MSE	MET	modified residue	GB 24371665
A	95	MSE	MET	modified residue	GB 24371665
A	96	MSE	MET	modified residue	GB 24371665
A	103	MSE	MET	modified residue	GB 24371665
A	157	MSE	MET	modified residue	GB 24371665
B	1	MSE	MET	modified residue	GB 24371665
B	61	MSE	MET	modified residue	GB 24371665
B	95	MSE	MET	modified residue	GB 24371665
B	96	MSE	MET	modified residue	GB 24371665
B	103	MSE	MET	modified residue	GB 24371665
B	157	MSE	MET	modified residue	GB 24371665
C	1	MSE	MET	modified residue	GB 24371665
C	61	MSE	MET	modified residue	GB 24371665
C	95	MSE	MET	modified residue	GB 24371665
C	96	MSE	MET	modified residue	GB 24371665
C	103	MSE	MET	modified residue	GB 24371665
C	157	MSE	MET	modified residue	GB 24371665


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	52	Total 52	O 52	0	0
2	B	21	Total 21	O 21	0	0
2	C	49	Total 49	O 49	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: molybdenum cofactor biosynthesis protein Mog



- Molecule 1: molybdenum cofactor biosynthesis protein Mog



- Molecule 1: molybdenum cofactor biosynthesis protein Mog



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.57Å 215.36Å 40.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 50.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.6 (50.00-2.30) 98.5 (50.00-2.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.74 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.193 , 0.222 0.191 , 0.221	Depositor DCC
$R_{free}$ test set	1717 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.6	Xtriage
Anisotropy	0.331	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 57.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4026	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.97	1/1334 (0.1%)	1.05	10/1805 (0.6%)
1	B	1.50	12/1303 (0.9%)	0.96	4/1762 (0.2%)
1	C	0.96	4/1320 (0.3%)	0.91	3/1787 (0.2%)
All	All	1.17	17/3957 (0.4%)	0.98	17/5354 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	15	ALA	C-O	29.03	1.78	1.23
1	B	21	GLU	CD-OE1	16.85	1.44	1.25
1	B	21	GLU	CD-OE2	16.23	1.43	1.25
1	B	20	TYR	CG-CD2	12.64	1.55	1.39
1	B	24	SER	CB-OG	12.21	1.58	1.42
1	C	89	GLU	CG-CD	9.39	1.66	1.51
1	B	20	TYR	CE1-CZ	9.27	1.50	1.38
1	B	20	TYR	CE2-CZ	7.93	1.48	1.38
1	B	161	TYR	CG-CD1	7.91	1.49	1.39
1	B	161	TYR	CE2-CZ	7.80	1.48	1.38
1	B	18	GLY	N-CA	7.78	1.57	1.46
1	C	161	TYR	CG-CD2	7.08	1.48	1.39
1	A	142	CYS	CB-SG	6.64	1.93	1.82
1	C	161	TYR	CE1-CZ	6.23	1.46	1.38
1	B	161	TYR	CE1-CZ	5.93	1.46	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	161	TYR	CG-CD2	5.80	1.46	1.39
1	C	89	GLU	CD-OE2	5.14	1.31	1.25

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	20	TYR	CB-CG-CD2	-12.77	113.34	121.00
1	A	173	ARG	NE-CZ-NH1	-11.93	114.33	120.30
1	A	118	ARG	NE-CZ-NH2	-9.35	115.63	120.30
1	B	20	TYR	CG-CD2-CE2	-9.09	114.03	121.30
1	A	155	ASP	CB-CG-OD2	-7.73	111.34	118.30
1	A	173	ARG	NE-CZ-NH2	7.39	124.00	120.30
1	C	173	ARG	NE-CZ-NH2	-6.87	116.86	120.30
1	A	155	ASP	CB-CG-OD1	6.79	124.42	118.30
1	A	102	LEU	CB-CG-CD2	6.72	122.42	111.00
1	A	118	ARG	NE-CZ-NH1	6.28	123.44	120.30
1	B	15	ALA	CA-C-O	-5.89	107.73	120.10
1	C	102	LEU	CB-CG-CD2	5.59	120.51	111.00
1	A	19	ILE	N-CA-C	5.44	125.70	111.00
1	A	94	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	B	21	GLU	OE1-CD-OE2	5.10	129.43	123.30
1	C	173	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	A	24	SER	CB-CA-C	5.06	119.71	110.10

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	19	ILE	Peptide
1	B	14	ARG	Peptide
1	B	20	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1316	0	1324	14	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1286	0	1293	31	0
1	C	1302	0	1309	22	0
2	A	52	0	0	1	0
2	B	21	0	0	2	0
2	C	49	0	0	5	0
All	All	4026	0	3926	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:103:MSE:SE	1:C:103:MSE:CE	2.17	1.41
1:B:15:ALA:O	1:B:15:ALA:C	1.78	1.20
1:C:16:SER:HB3	2:C:184:HOH:O	1.42	1.20
1:A:173:ARG:HG2	1:A:173:ARG:HH11	1.13	1.07
1:B:14:ARG:HH11	1:B:14:ARG:HG3	1.21	1.02
1:A:173:ARG:HH11	1:A:173:ARG:CG	1.82	0.88
1:C:67:CYS:SG	2:C:206:HOH:O	2.33	0.87
1:A:155:ASP:OD2	1:A:173:ARG:NH1	2.07	0.86
1:A:173:ARG:HG2	1:A:173:ARG:NH1	1.84	0.86
2:A:182:HOH:O	1:B:94:ARG:HD2	1.81	0.80
1:C:53:VAL:O	1:C:57:THR:CG2	2.30	0.80
1:C:53:VAL:O	1:C:57:THR:HG23	1.83	0.79
1:B:14:ARG:HH11	1:B:14:ARG:CG	2.00	0.71
1:C:5:LYS:HE2	1:C:65:GLN:O	1.93	0.68
1:B:14:ARG:HG3	1:B:14:ARG:NH1	1.99	0.64
1:B:162:LEU:H	1:B:162:LEU:HD12	1.63	0.63
1:B:135:LYS:HD3	1:B:136:PRO:HD2	1.81	0.62
1:A:173:ARG:CG	1:A:173:ARG:NH1	2.50	0.61
1:B:135:LYS:HD3	1:B:136:PRO:N	2.16	0.61
1:C:94:ARG:NH2	1:C:157:MSE:HE1	2.16	0.60
1:B:135:LYS:HD3	1:B:136:PRO:CD	2.32	0.60
1:C:89:GLU:CG	2:C:216:HOH:O	2.52	0.58
1:C:28:ILE:HG21	1:C:71:VAL:HG21	1.85	0.57
1:C:28:ILE:HD13	1:C:71:VAL:HG23	1.86	0.57
1:C:53:VAL:O	1:C:57:THR:HG22	2.05	0.57
1:B:5:LYS:HE2	1:B:65:GLN:O	2.05	0.56
1:B:102:LEU:HD13	1:B:145:ALA:HB1	1.88	0.56
1:B:135:LYS:HD2	1:B:137:LYS:H	1.71	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:96:MSE:HE3	1:B:99:PHE:HE1	1.73	0.54
1:C:28:ILE:HG21	1:C:71:VAL:CG2	2.40	0.52
1:B:71:VAL:HG11	1:B:132:LEU:CD1	2.41	0.51
1:B:34:ASP:O	1:B:168:VAL:HG21	2.10	0.51
1:A:148:PRO:HD3	1:A:169:ILE:HD11	1.93	0.51
1:B:49:ASP:CB	2:B:182:HOH:O	2.59	0.50
1:A:172:PHE:CE2	1:A:174:PRO:HG3	2.47	0.50
1:A:155:ASP:CG	1:A:173:ARG:HH12	2.09	0.50
1:C:135:LYS:HB2	2:C:192:HOH:O	2.11	0.49
1:A:19:ILE:N	1:A:20:TYR:HB2	2.28	0.49
1:B:5:LYS:HG2	1:B:43:ILE:HD11	1.95	0.49
1:C:28:ILE:CG2	1:C:71:VAL:HG21	2.43	0.49
1:A:165:ASN:HD22	1:A:165:ASN:C	2.16	0.48
1:C:94:ARG:HH22	1:C:157:MSE:HE1	1.79	0.48
1:B:49:ASP:HB3	2:B:182:HOH:O	2.12	0.48
1:B:147:PHE:HA	1:B:150:ILE:HD12	1.97	0.47
1:B:135:LYS:HD3	1:B:135:LYS:C	2.34	0.47
1:A:116:LEU:HD22	1:B:96:MSE:HE1	1.97	0.46
1:A:102:LEU:HD22	1:A:106:GLU:HG2	1.97	0.46
1:B:148:PRO:HD3	1:B:169:ILE:HD11	1.97	0.46
1:B:151:PRO:HA	1:B:162:LEU:HD13	1.97	0.46
1:A:45:GLN:HG3	1:A:61:MSE:HE3	1.99	0.45
1:C:89:GLU:HG2	2:C:216:HOH:O	2.15	0.45
1:C:28:ILE:HD13	1:C:71:VAL:CG2	2.48	0.44
1:B:152:TYR:O	1:B:155:ASP:HB2	2.18	0.44
1:B:89:GLU:HG2	1:B:95:MSE:HE2	1.99	0.44
1:B:162:LEU:HD12	1:B:162:LEU:N	2.33	0.43
1:A:108:LEU:HD11	1:A:115:ILE:CG2	2.49	0.43
1:B:80:LYS:HD2	1:C:94:ARG:CZ	2.48	0.43
1:B:76:THR:CG2	1:B:117:SER:OG	2.67	0.42
1:B:156:LEU:HB2	1:B:157:MSE:HE2	2.01	0.42
1:C:147:PHE:N	1:C:148:PRO:CD	2.83	0.41
1:C:157:MSE:H	1:C:157:MSE:HG2	1.75	0.41
1:C:165:ASN:C	1:C:165:ASN:HD22	2.23	0.41
1:B:8:ILE:HD13	1:B:29:ILE:HG13	2.03	0.40
1:C:84:THR:N	1:C:85:PRO:CD	2.84	0.40
1:B:14:ARG:HB3	1:B:15:ALA:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/177 (97%)	167 (98%)	4 (2%)	0	100	100
1	B	164/177 (93%)	161 (98%)	3 (2%)	0	100	100
1	C	169/177 (96%)	163 (96%)	6 (4%)	0	100	100
All	All	504/531 (95%)	491 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/144 (101%)	134 (92%)	12 (8%)	9	12
1	B	144/144 (100%)	131 (91%)	13 (9%)	8	10
1	C	145/144 (101%)	137 (94%)	8 (6%)	18	26
All	All	435/432 (101%)	402 (92%)	33 (8%)	11	14

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ARG
1	A	21	GLU
1	A	26	LYS
1	A	80	LYS
1	A	89	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	94	ARG
1	A	102	LEU
1	A	108	LEU
1	A	137	LYS
1	A	165	ASN
1	A	170	LYS
1	A	173	ARG
1	B	13	ASP
1	B	14	ARG
1	B	19	ILE
1	B	46	VAL
1	B	50	GLU
1	B	76	THR
1	B	94	ARG
1	B	102	LEU
1	B	135	LYS
1	B	137	LYS
1	B	162	LEU
1	B	166	GLU
1	B	168	VAL
1	C	45	GLN
1	C	57	THR
1	C	102	LEU
1	C	135	LYS
1	C	140	ARG
1	C	157	MSE
1	C	165	ASN
1	C	170	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	ASN
1	B	119	GLN
1	B	165	ASN
1	C	165	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

**Warning:** The R factor obtained from EDS is 0.2774, which does not match the depositor's R factor of 0.19282. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	168/177 (94%)	1.06	25 (14%) 7 8	35, 45, 59, 64	0
1	B	163/177 (92%)	1.05	29 (17%) 4 5	31, 41, 58, 68	0
1	C	166/177 (93%)	1.20	24 (14%) 7 8	29, 43, 57, 61	0
All	All	497/531 (93%)	1.10	78 (15%) 6 7	29, 43, 58, 68	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	20	TYR	6.3
1	B	15	ALA	5.8
1	B	20	TYR	5.7
1	B	19	ILE	5.5
1	A	19	ILE	5.0
1	B	174	PRO	4.4
1	C	20	TYR	4.3
1	B	137	LYS	4.2
1	A	152	TYR	4.1
1	B	138	SER	4.0
1	B	139	ILE	3.9
1	B	18	GLY	3.6
1	A	154	ILE	3.6
1	B	172	PHE	3.5
1	C	19	ILE	3.5
1	C	39	GLU	3.4
1	B	24	SER	3.4
1	A	156	LEU	3.3
1	C	174	PRO	3.3
1	B	142	CYS	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	115	ILE	3.2
1	A	14	ARG	3.1
1	A	17	ALA	3.1
1	A	173	ARG	3.1
1	C	108	LEU	3.0
1	B	166	GLU	3.0
1	B	35	TYR	3.0
1	C	172	PHE	2.9
1	B	23	ILE	2.9
1	A	172	PHE	2.8
1	B	167	ALA	2.8
1	B	136	PRO	2.8
1	A	153	CYS	2.8
1	C	161	TYR	2.7
1	C	111	VAL	2.7
1	C	89	GLU	2.7
1	A	175	LYS	2.7
1	C	138	SER	2.6
1	A	161	TYR	2.6
1	B	68	CYS	2.6
1	B	161	TYR	2.5
1	B	143	LEU	2.5
1	C	112	PRO	2.5
1	A	110	PHE	2.5
1	C	164	CYS	2.5
1	A	3	LYS	2.4
1	C	4	ALA	2.4
1	C	160	PRO	2.4
1	B	156	LEU	2.4
1	A	111	VAL	2.4
1	C	167	ALA	2.4
1	A	150	ILE	2.3
1	C	21	GLU	2.3
1	B	14	ARG	2.3
1	A	21	GLU	2.3
1	B	89	GLU	2.3
1	C	105	ALA	2.3
1	A	135	LYS	2.2
1	A	108	LEU	2.2
1	A	37	THR	2.2
1	A	40	TRP	2.2
1	C	63	ASP	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	64	GLU	2.2
1	B	108	LEU	2.2
1	A	15	ALA	2.2
1	B	13	ASP	2.2
1	A	159	GLY	2.2
1	C	14	ARG	2.2
1	A	24	SER	2.2
1	B	171	PRO	2.2
1	B	46	VAL	2.2
1	B	173	ARG	2.1
1	C	60	LYS	2.1
1	C	134	GLY	2.1
1	B	145	ALA	2.1
1	B	110	PHE	2.0
1	A	80	LYS	2.0
1	C	169	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.