



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 25, 2024 – 05:47 AM EDT

PDB ID : 6FCX
Title : Structure of human 5,10-methylenetetrahydrofolate reductase (MTHFR)
Authors : Kopec, J.; Bezerra, G.A.; Oberholzer, A.E.; Rembeza, E.; Sorrell, F.J.; Chalk, R.; Borkowska, O.; Ellis, K.; Kupinska, K.; Krojer, T.; Burgess-Brown, N.; Von Delft, F.; Arrowsmith, C.; Edwards, A.; Bountra, C.; Froese, D.S.; Baumgartner, M.; Yue, W.W.; Structural Genomics Consortium (SGC)
Deposited on : 2017-12-21
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	615	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 79%, yellow 79%, yellow 96%, green 96%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 79% 17% ... </div> </div>
1	B	615	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 8%, orange 8%, orange 73%, yellow 73%, yellow 94%, green 94%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 8% 73% 21% .. </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9180 atoms, of which 10 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylenetetrahydrofolate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	596	Total	C	N	O	S	0	0	0
			4723	3048	786	870	19			
1	B	590	Total	C	N	O	S	0	0	0
			4191	2660	727	791	13			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	37	MET	LEU	initiating methionine	UNP P42898
A	429	ALA	GLU	variant	UNP P42898
A	594	GLN	ARG	variant	UNP P42898
A	645	ALA	-	expression tag	UNP P42898
A	646	GLU	-	expression tag	UNP P42898
A	647	ASN	-	expression tag	UNP P42898
A	648	LEU	-	expression tag	UNP P42898
A	649	TYR	-	expression tag	UNP P42898
A	650	PHE	-	expression tag	UNP P42898
A	651	GLN	-	expression tag	UNP P42898
B	37	MET	LEU	initiating methionine	UNP P42898
B	429	ALA	GLU	variant	UNP P42898
B	594	GLN	ARG	variant	UNP P42898
B	645	ALA	-	expression tag	UNP P42898
B	646	GLU	-	expression tag	UNP P42898
B	647	ASN	-	expression tag	UNP P42898
B	648	LEU	-	expression tag	UNP P42898
B	649	TYR	-	expression tag	UNP P42898
B	650	PHE	-	expression tag	UNP P42898
B	651	GLN	-	expression tag	UNP P42898

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			18	6	5	7		
4	B	1	Total	C	H	O	0	0
			18	6	5	7		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	48	Total	O	0	0
			48	48		
5	B	24	Total	O	0	0
			24	24		

- Molecule 1: Methylenetetrahydrofolate reductase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.34Å 127.94Å 147.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.97 – 2.50 63.97 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (63.97-2.50) 100.0 (63.97-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.210 , 0.247 0.215 , 0.250	Depositor DCC
R_{free} test set	3266 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	69.2	Xtriage
Anisotropy	0.359	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9180	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, CIT, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/4853	0.83	13/6606 (0.2%)
1	B	0.55	0/4295	0.70	5/5885 (0.1%)
All	All	0.61	0/9148	0.77	18/12491 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	2
All	All	0	7

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	TYR	CB-CG-CD2	-9.15	115.51	121.00
1	B	515	PHE	CB-CG-CD2	-8.94	114.55	120.80
1	A	65	PHE	CB-CG-CD2	-8.56	114.81	120.80
1	A	60	PHE	CB-CG-CD2	-7.69	115.42	120.80
1	A	515	PHE	CB-CG-CD2	-7.60	115.48	120.80

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	404	TYR	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	515	PHE	Sidechain
1	A	60	PHE	Sidechain
1	A	65	PHE	Sidechain
1	A	90	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4723	0	4516	120	0
1	B	4191	0	3577	140	0
2	A	53	0	31	3	0
2	B	53	0	30	2	0
3	A	26	0	19	0	0
3	B	26	0	17	0	0
4	A	13	5	5	1	0
4	B	13	5	5	1	0
5	A	48	0	0	0	0
5	B	24	0	0	2	0
All	All	9170	10	8200	251	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 251 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:LEU:HB2	1:B:91:ILE:HD11	1.31	1.11
1:B:62:LEU:HB2	1:B:91:ILE:CD1	1.81	1.09
1:A:529:LEU:HD13	1:A:538:TYR:CD2	1.92	1.05
1:A:439:LEU:O	1:A:468:LYS:NZ	1.93	0.99
1:B:157:ARG:HB3	2:B:701:FAD:H5'1	1.44	0.99

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	590/615 (96%)	570 (97%)	18 (3%)	2 (0%)	41	61
1	B	580/615 (94%)	548 (94%)	32 (6%)	0	100	100
All	All	1170/1230 (95%)	1118 (96%)	50 (4%)	2 (0%)	47	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	205	GLY
1	A	221	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	491/537 (91%)	481 (98%)	10 (2%)	55	79
1	B	358/537 (67%)	348 (97%)	10 (3%)	43	70
All	All	849/1074 (79%)	829 (98%)	20 (2%)	49	74

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	414	LYS
1	B	538	TYR
1	B	620	ASN
1	B	619	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	435	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	96	HIS
1	B	477	GLN
1	B	555	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SAH	A	702	-	24,28,28	1.38	4 (16%)	25,40,40	1.89	7 (28%)
2	FAD	A	701	-	53,58,58	0.51	0	68,89,89	0.67	1 (1%)
3	SAH	B	702	1	24,28,28	1.29	2 (8%)	25,40,40	1.77	6 (24%)
2	FAD	B	701	1	53,58,58	0.46	0	68,89,89	0.58	1 (1%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CIT	A	703	-	12,12,12	1.04	0	17,17,17	1.72	4 (23%)
4	CIT	B	703	-	12,12,12	0.99	0	17,17,17	1.47	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	A	702	-	-	0/11/31/31	0/3/3/3
2	FAD	A	701	-	-	4/30/50/50	0/6/6/6
3	SAH	B	702	1	-	3/11/31/31	0/3/3/3
2	FAD	B	701	1	-	16/30/50/50	0/6/6/6
4	CIT	A	703	-	-	8/16/16/16	-
4	CIT	B	703	-	-	7/16/16/16	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	SAH	C2-N3	4.08	1.38	1.32
3	B	702	SAH	C2-N3	3.93	1.38	1.32
3	B	702	SAH	C2-N1	3.00	1.39	1.33
3	A	702	SAH	C2-N1	2.61	1.38	1.33
3	A	702	SAH	OXT-C	-2.43	1.22	1.30

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	SAH	N3-C2-N1	-6.21	118.97	128.68
3	B	702	SAH	N3-C2-N1	-4.74	121.26	128.68
4	A	703	CIT	O6-C6-C3	4.01	120.01	113.05
4	B	703	CIT	O6-C6-C3	3.68	119.44	113.05
3	A	702	SAH	C5'-SD-CG	-3.00	93.27	102.27

There are no chirality outliers.

5 of 38 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	FAD	C1'-C2'-C3'-O3'
2	A	701	FAD	C1'-C2'-C3'-C4'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	701	FAD	C5B-O5B-PA-O1A
2	B	701	FAD	C1'-C2'-C3'-C4'
2	B	701	FAD	C3'-C4'-C5'-O5'


There are no ring outliers.

4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	FAD	3	0
2	B	701	FAD	2	0
4	A	703	CIT	1	0
4	B	703	CIT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand SAH A 702




Ligand FAD A 701



Ligand SAH B 702



Ligand FAD B 701



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	596/615 (96%)	-0.08	8 (1%) 77 79	48, 74, 131, 160	0
1	B	590/615 (95%)	0.34	48 (8%) 12 12	52, 97, 194, 241	0
All	All	1186/1230 (96%)	0.13	56 (4%) 31 33	48, 83, 181, 241	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	214	LEU	9.5
1	B	250	CYS	8.8
1	B	240	VAL	6.5
1	B	248	ILE	6.3
1	B	236	PHE	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FAD	B	701	53/53	0.82	0.18	102,125,139,142	0
4	CIT	B	703	13/13	0.83	0.18	144,145,175,175	0
4	CIT	A	703	13/13	0.89	0.11	121,124,150,150	0
2	FAD	A	701	53/53	0.95	0.12	80,88,100,101	0
3	SAH	B	702	26/26	0.95	0.15	60,64,69,70	0
3	SAH	A	702	26/26	0.98	0.15	47,53,59,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FAD B 701:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FAD A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAH B 702:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.