



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 27, 2024 – 08:51 AM EDT

PDB ID : 1FYH
Title : 1:1 COMPLEX BETWEEN AN INTERFERON GAMMA SINGLE-CHAIN
VARIANT AND ITS RECEPTOR
Authors : Randal, M.; Kossiakoff, A.A.
Deposited on : 2000-09-29
Resolution : 2.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : NOT EXECUTED
EDS : NOT EXECUTED
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	2229 (2.04-2.04)
Ramachandran outliers	177936	2217 (2.04-2.04)
Sidechain outliers	177891	2217 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	258	 74% 16% • 6%
1	D	258	 82% 10% • 7%
2	B	229	 77% 9% • 12%
2	E	229	 75% 11% • 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7687 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	242	Total	C	N	O	S	80	0	0
			1996	1272	333	384	7			
1	D	240	Total	C	N	O	S	24	0	0
			1984	1267	331	379	7			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P01579
A	111	ASP	HIS	engineered mutation	UNP P01579
A	121	GLY	-	linker	UNP P01579
A	122	ALA	-	linker	UNP P01579
A	123	ASN	-	linker	UNP P01579
A	124	VAL	-	linker	UNP P01579
A	201	SER	-	linker	UNP P01579
A	202	GLY	-	linker	UNP P01579
A	203	GLU	-	linker	UNP P01579
A	204	PHE	-	linker	UNP P01579
D	0	MET	-	initiating methionine	UNP P01579
D	111	ASP	HIS	engineered mutation	UNP P01579
D	121	GLY	-	linker	UNP P01579
D	122	ALA	-	linker	UNP P01579
D	123	ASN	-	linker	UNP P01579
D	124	VAL	-	linker	UNP P01579
D	201	SER	-	linker	UNP P01579
D	202	GLY	-	linker	UNP P01579
D	203	GLU	-	linker	UNP P01579
D	204	PHE	-	linker	UNP P01579

- Molecule 2 is a protein called Interferon gamma receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	201	Total	C	N	O	S	49	0	0
			1612	1025	267	308	12			
2	E	201	Total	C	N	O	S	92	0	0
			1613	1028	266	307	12			

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	105	Total	O	0	0
			105	105		
4	B	139	Total	O	0	0
			139	139		
4	D	143	Total	O	0	0
			143	143		
4	E	94	Total	O	0	0
			94	94		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: Interferon gamma

Chain A: 





- Molecule 1: Interferon gamma

Chain D: 



- Molecule 2: Interferon gamma receptor 1

Chain B: 



- Molecule 2: Interferon gamma receptor 1

Chain E: 



C178	
C183	
Q184	
I187	
P188	
V189	
S194	
V203	
I221	
F222	
ASN	
SER	
SER	
ILF	
LYS	
GLY	
SER	

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.94Å 107.60Å 85.08Å 90.00° 97.94° 90.00°	Depositor
Resolution (Å)	15.00 – 2.04	Depositor
% Data completeness (in resolution range)	95.2 (15.00-2.04)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.194 , 0.246	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7687	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.71	0/2028	0.88	0/2712
1	D	0.75	1/2016 (0.0%)	0.81	0/2694
2	B	0.71	0/1649	0.91	1/2242 (0.0%)
2	E	0.70	0/1651	0.91	1/2245 (0.0%)
All	All	0.72	1/7344 (0.0%)	0.88	2/9893 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	271	GLU	CG-CD	5.54	1.60	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	115	LEU	CA-CB-CG	6.58	130.44	115.30
2	B	115	LEU	CA-CB-CG	5.01	126.82	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1996	0	1988	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1984	0	1989	17	0
2	B	1612	0	1568	11	0
2	E	1613	0	1577	20	0
3	A	1	0	0	0	0
4	A	105	0	0	2	0
4	B	139	0	0	1	0
4	D	143	0	0	4	0
4	E	94	0	0	1	0
All	All	7687	0	7122	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:GLU:HG3	1:A:272:THR:HG21	1.67	0.76
1:A:275:GLU:O	1:A:279:VAL:HG23	1.88	0.74
1:D:289:ARG:O	1:D:293:GLU:HG3	1.89	0.72
2:E:124:ILE:HD12	2:E:189:VAL:HG22	1.75	0.69
2:B:88:ARG:HH11	2:B:88:ARG:HG2	1.59	0.67

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	238/258 (92%)	225 (94%)	10 (4%)	3 (1%)	10 4
1	D	236/258 (92%)	234 (99%)	2 (1%)	0	100 100
2	B	197/229 (86%)	185 (94%)	11 (6%)	1 (0%)	25 17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	197/229 (86%)	182 (92%)	15 (8%)	0	100	100
All	All	868/974 (89%)	826 (95%)	38 (4%)	4 (0%)	25	17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	264	GLN
2	B	90	GLY
1	A	270	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/236 (94%)	206 (92%)	17 (8%)	11	5
1	D	222/236 (94%)	209 (94%)	13 (6%)	16	9
2	B	185/209 (88%)	171 (92%)	14 (8%)	11	5
2	E	186/209 (89%)	174 (94%)	12 (6%)	14	7
All	All	816/890 (92%)	760 (93%)	56 (7%)	13	7

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	165	GLU
2	E	194	SER
1	D	67	GLN
2	E	184	GLN
2	E	161	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	17	ASN
1	D	259	ASN
1	D	306	GLN
1	D	304	ASN
1	A	304	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.