



## wwPDB EM Validation Summary Report ⓘ

Aug 11, 2025 – 01:50 pm BST

PDB ID : 9GE5 / pdb\_00009ge5  
EMDB ID : EMD-51290  
Title : CryoEM structure of the human INO80-Hexasome complex  
Authors : Sharma, M.; Aggarwal, P.; Hopfner, K.P.  
Deposited on : 2024-08-07  
Resolution : 3.35 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev126  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4-5-2 with Phenix2.0rc1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	
1	B	453	
1	C	453	
2	D	439	
2	E	439	
2	F	439	
3	G	733	
4	H	135	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	I	107	
6	J	576	
7	K	113	
8	L	113	
9	M	94	
9	Q	94	
10	N	81	
10	R	81	
11	S	108	
12	T	92	

## 2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 40017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RuvB-like 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	434	Total	C	N	O	S	0	0
			3345	2107	575	647	16		
1	B	447	Total	C	N	O	S	0	0
			3454	2173	592	673	16		
1	C	443	Total	C	N	O	S	0	0
			3413	2148	585	663	17		

- Molecule 2 is a protein called RuvB-like 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	426	Total	C	N	O	S	0	0
			3318	2074	583	646	15		
2	E	439	Total	C	N	O	S	0	0
			3404	2125	597	666	16		
2	F	434	Total	C	N	O	S	0	0
			3377	2110	592	659	16		

- Molecule 3 is a protein called Chromatin-remodeling ATPase INO80.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	682	Total	C	N	O	S	0	0
			5609	3599	1004	976	30		

- Molecule 4 is a protein called INO80 complex subunit B.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	H	101	Total	C	N	O	S	0	0
			786	478	161	138	9		

- Molecule 5 is a protein called INO80 complex subunit C.

Mol	Chain	Residues	Atoms				AltConf	Trace
5	I	107	Total	C	N	O	0	0
			851	545	150	156		

- Molecule 6 is a protein called Actin-related protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	412	Total	C	N	O	S	0	0
			3278	2088	572	597	21		

- Molecule 7 is a DNA chain called Hexasomal DNA Strand 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	K	113	Total	C	N	O	P	0	0
			2334	1104	441	676	113		

- Molecule 8 is a DNA chain called Hexasomal DNA strand 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	L	113	Total	C	N	O	P	0	0
			2299	1093	413	680	113		

- Molecule 9 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	M	91	Total	C	N	O	S	0	0
			739	467	138	130	4		
9	Q	94	Total	C	N	O	S	0	0
			767	482	147	134	4		

- Molecule 10 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	N	81	Total	C	N	O	S	0	0
			646	407	126	112	1		
10	R	78	Total	C	N	O	S	0	0
			619	391	120	107	1		

- Molecule 11 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				AltConf	Trace
11	S	108	Total	C	N	O	0	0
			836	526	165	145		

- | Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 12  | T     | 92       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 721   | 453 | 129 | 137 | 2 |         |       |

- # ATP

Mol	Chain	Residues	Atoms					AltConf
13	A	1	Total 31	C 10	N 5	O 13	P 3	0

- # ADP

Mol	Chain	Residues	Atoms					AltConf
14	B	1	Total	C	N	O	P	0
			27	10	5	10	2	
14	C	1	Total	C	N	O	P	0
			27	10	5	10	2	
14	D	1	Total	C	N	O	P	0
			27	10	5	10	2	
14	E	1	Total	C	N	O	P	0
			27	10	5	10	2	
14	F	1	Total	C	N	O	P	0
			27	10	5	10	2	
14	G	1	Total	C	N	O	P	0
			27	10	5	10	2	
14	J	1	Total	C	N	O	P	0
			27	10	5	10	2	


- Molecule 15 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
15	H	1	Total	Zn	0
			1	1	


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RuvB-like 1




- Molecule 1: RuvB-like 1



- Molecule 1: RuvB-like 1



- Molecule 2: RuvB-like 2




- Molecule 2: RuvB-like 2








- Molecule 2: RuvB-like 2





- Molecule 3: Chromatin-remodeling ATPase INO80





- Molecule 4: INO80 complex subunit B




- Molecule 5: INO80 complex subunit C





• Molecule 6: Actin-related protein 5




• Molecule 7: Hexasomal DNA Strand 1




• Molecule 8: Hexasomal DNA strand 2




• Molecule 9: Histone H3.1









• Molecule 9: Histone H3.1




• Molecule 10: Histone H4




• Molecule 10: Histone H4



• Molecule 11: Histone H2A type 1-B/E



• Molecule 12: Histone H2B type 1-B



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	34444	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.0	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.156	Depositor
Minimum map value	-0.676	Depositor
Average map value	0.005	Depositor
Map value standard deviation	0.053	Depositor
Recommended contour level	0.2	Depositor
Map size (Å)	290.8, 290.8, 290.8	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.727, 0.727, 0.727	Depositor

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ZN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.68	0/3388	1.24	6/4564 (0.1%)
1	B	0.69	0/3497	1.24	9/4710 (0.2%)
1	C	0.69	0/3459	1.22	8/4662 (0.2%)
2	D	0.70	0/3354	1.25	13/4510 (0.3%)
2	E	0.71	0/3444	1.25	4/4636 (0.1%)
2	F	0.70	0/3416	1.26	11/4596 (0.2%)
3	G	0.78	0/5745	1.31	16/7759 (0.2%)
4	H	0.85	0/803	1.45	3/1080 (0.3%)
5	I	3.15	3/874 (0.3%)	1.54	10/1183 (0.8%)
6	J	0.79	0/3363	1.33	6/4558 (0.1%)
7	K	1.27	5/2622 (0.2%)	1.33	7/4049 (0.2%)
8	L	0.67	0/2574	1.34	5/3966 (0.1%)
9	M	0.80	0/748	1.37	4/1003 (0.4%)
9	Q	0.86	0/776	1.37	2/1039 (0.2%)
10	N	0.87	0/653	1.36	1/873 (0.1%)
10	R	0.86	0/626	1.40	1/837 (0.1%)
11	S	0.82	0/846	1.39	1/1139 (0.1%)
12	T	0.82	0/732	1.31	0/983
All	All	0.90	8/40920 (0.0%)	1.30	107/56147 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2
2	D	0	1
2	E	0	4
2	F	0	4
3	G	0	3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	#Chirality outliers	#Planarity outliers
6	J	0	2
7	K	0	44
8	L	0	26
9	Q	0	1
11	S	0	1
All	All	0	88

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	136	LYS	CA-CB	89.44	3.03	1.53
7	K	3	DG	O4'-C1'	26.23	1.94	1.41
7	K	3	DG	C2'-C1'	24.93	2.02	1.52
7	K	3	DG	C4'-C3'	24.76	2.02	1.52
7	K	3	DG	C3'-C2'	23.67	2.00	1.52

The worst 5 of 107 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	136	LYS	N-CA-C	-14.11	95.39	112.89
5	I	136	LYS	N-CA-CB	10.31	127.20	110.40
5	I	136	LYS	CA-CB-CG	10.19	134.49	114.10
7	K	3	DG	O4'-C1'-N9	9.96	123.34	108.40
8	L	-26	DC	C4'-C3'-O3'	7.01	120.52	110.00

There are no chirality outliers.

5 of 88 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	192	TYR	Sidechain
1	B	202	ARG	Sidechain
2	D	172	TYR	Sidechain
2	E	18	ARG	Sidechain
2	E	71	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3345	0	3451	2	0
1	B	3454	0	3563	1	0
1	C	3413	0	3508	3	0
2	D	3318	0	3395	0	0
2	E	3404	0	3472	2	0
2	F	3377	0	3446	1	0
3	G	5609	0	5651	4	0
4	H	786	0	798	1	0
5	I	851	0	852	30	0
6	J	3278	0	3215	2	0
7	K	2334	0	1268	36	0
8	L	2299	0	1270	0	0
9	M	739	0	777	0	0
9	Q	767	0	807	0	0
10	N	646	0	687	1	0
10	R	619	0	659	0	0
11	S	836	0	897	1	0
12	T	721	0	740	0	0
13	A	31	0	12	0	0
14	B	27	0	12	0	0
14	C	27	0	12	0	0
14	D	27	0	12	0	0
14	E	27	0	12	0	0
14	F	27	0	12	0	0
14	G	27	0	12	0	0
14	J	27	0	12	0	0
15	H	1	0	0	0	0
All	All	40017	0	38552	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:K:3:DG:C3'	7:K:3:DG:C2'	2.00	1.39
7:K:3:DG:C2'	7:K:3:DG:C1'	2.02	1.37
7:K:3:DG:C3'	7:K:3:DG:C4'	2.02	1.35
5:I:136:LYS:HA	7:K:3:DG:C2'	1.63	1.28
5:I:136:LYS:HA	7:K:3:DG:C1'	1.73	1.19

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/453 (95%)	413 (96%)	15 (4%)	2 (0%)	25	53
1	B	443/453 (98%)	430 (97%)	11 (2%)	2 (0%)	25	53
1	C	441/453 (97%)	421 (96%)	19 (4%)	1 (0%)	44	71
2	D	420/439 (96%)	408 (97%)	10 (2%)	2 (0%)	25	53
2	E	437/439 (100%)	427 (98%)	10 (2%)	0	100	100
2	F	430/439 (98%)	414 (96%)	15 (4%)	1 (0%)	44	71
3	G	674/733 (92%)	634 (94%)	32 (5%)	8 (1%)	11	34
4	H	97/135 (72%)	93 (96%)	3 (3%)	1 (1%)	13	39
5	I	105/107 (98%)	97 (92%)	7 (7%)	1 (1%)	13	39
6	J	406/576 (70%)	373 (92%)	30 (7%)	3 (1%)	19	47
9	M	89/94 (95%)	86 (97%)	3 (3%)	0	100	100
9	Q	92/94 (98%)	87 (95%)	4 (4%)	1 (1%)	12	37
10	N	79/81 (98%)	73 (92%)	4 (5%)	2 (2%)	4	21
10	R	76/81 (94%)	73 (96%)	2 (3%)	1 (1%)	10	32
11	S	106/108 (98%)	104 (98%)	2 (2%)	0	100	100
12	T	90/92 (98%)	86 (96%)	4 (4%)	0	100	100
All	All	4415/4777 (92%)	4219 (96%)	171 (4%)	25 (1%)	24	50

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	257	LEU
10	N	26	ILE
1	B	59	LYS
1	B	342	GLU
2	F	452	PHE



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/384 (96%)	364 (99%)	3 (1%)	79	88
1	B	381/384 (99%)	376 (99%)	5 (1%)	65	79
1	C	374/384 (97%)	370 (99%)	4 (1%)	70	82
2	D	361/369 (98%)	355 (98%)	6 (2%)	56	74
2	E	369/369 (100%)	367 (100%)	2 (0%)	86	92
2	F	367/369 (100%)	357 (97%)	10 (3%)	40	64
3	G	613/660 (93%)	609 (99%)	4 (1%)	81	89
4	H	86/105 (82%)	84 (98%)	2 (2%)	45	68
5	I	92/92 (100%)	92 (100%)	0	100	100
6	J	351/504 (70%)	348 (99%)	3 (1%)	75	87
9	M	79/81 (98%)	78 (99%)	1 (1%)	65	79
9	Q	81/81 (100%)	79 (98%)	2 (2%)	42	66
10	N	66/66 (100%)	66 (100%)	0	100	100
10	R	63/66 (96%)	63 (100%)	0	100	100
11	S	85/85 (100%)	85 (100%)	0	100	100
12	T	79/79 (100%)	77 (98%)	2 (2%)	42	66
All	All	3814/4078 (94%)	3770 (99%)	44 (1%)	66	80

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	328	THR
4	H	296	VAL
2	F	421	VAL
3	G	1148	ASP
6	J	258	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	580	ASN
3	G	855	HIS
11	S	24	GLN
3	G	581	ASN
3	G	811	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
14	ADP	B	501	-	24,29,29	1.19	2 (8%)	29,45,45	1.39	3 (10%)
14	ADP	C	501	-	24,29,29	1.11	2 (8%)	29,45,45	1.39	3 (10%)
14	ADP	D	501	-	24,29,29	1.13	2 (8%)	29,45,45	1.42	5 (17%)
14	ADP	G	1601	-	24,29,29	1.04	1 (4%)	29,45,45	1.60	7 (24%)
14	ADP	J	701	-	24,29,29	0.99	1 (4%)	29,45,45	1.46	6 (20%)
14	ADP	F	501	-	24,29,29	1.13	2 (8%)	29,45,45	1.43	3 (10%)
14	ADP	E	501	-	24,29,29	1.08	2 (8%)	29,45,45	1.39	6 (20%)
13	ATP	A	501	-	26,33,33	1.05	1 (3%)	31,52,52	1.67	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	ADP	B	501	-	-	1/12/32/32	0/3/3/3
14	ADP	C	501	-	-	0/12/32/32	0/3/3/3
14	ADP	D	501	-	-	3/12/32/32	0/3/3/3
14	ADP	G	1601	-	-	4/12/32/32	0/3/3/3
14	ADP	J	701	-	-	4/12/32/32	0/3/3/3
14	ADP	F	501	-	-	0/12/32/32	0/3/3/3
14	ADP	E	501	-	-	2/12/32/32	0/3/3/3
13	ATP	A	501	-	-	1/18/38/38	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	C	501	ADP	O4'-C1'	2.56	1.44	1.41
14	B	501	ADP	O4'-C1'	2.53	1.44	1.41
14	F	501	ADP	O4'-C1'	2.42	1.44	1.41
14	E	501	ADP	O4'-C1'	2.37	1.44	1.41
14	D	501	ADP	C5-C4	-2.32	1.34	1.40

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	C	501	ADP	C4-C5-N7	4.66	114.25	109.40
13	A	501	ATP	C4-C5-N7	4.52	114.11	109.40
14	G	1601	ADP	C4-C5-N7	4.38	113.96	109.40
14	B	501	ADP	C4-C5-N7	4.28	113.86	109.40
14	E	501	ADP	C4-C5-N7	4.20	113.77	109.40

There are no chirality outliers.


5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
14	G	1601	ADP	C5'-O5'-PA-O1A
14	J	701	ADP	O4'-C4'-C5'-O5'
14	J	701	ADP	C3'-C4'-C5'-O5'
14	J	701	ADP	C4'-C5'-O5'-PA
14	D	501	ADP	O4'-C4'-C5'-O5'



There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## Ligand ADP J 701




## Ligand ADP F 501









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-51290. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


### 6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 206




Y Index: 215




Z Index: 150


### 6.3.2 Raw map



X Index: 206



Y Index: 215




Z Index: 157

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)


### 6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


### 6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)




The volume at the recommended contour level is 297  $\text{nm}^3$ ; this corresponds to an approximate mass of 268 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.299  $\text{\AA}^{-1}$

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.299 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.35	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.44	8.24	4.69

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.44 differs from the reported value 3.35 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-51290 and PDB model 9GE5. Per-residue inclusion information can be found in section [3](#) on page [8](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).







































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7400	 0.3850
A	 0.8550	 0.4740
B	 0.8280	 0.4630
C	 0.8250	 0.4580
D	 0.8350	 0.4660
E	 0.8280	 0.4590
F	 0.8350	 0.4670
G	 0.6800	 0.3660
H	 0.7220	 0.3600
I	 0.6660	 0.3710
J	 0.7020	 0.3610
K	 0.7110	 0.2440
L	 0.6820	 0.2360
M	 0.5240	 0.2440
N	 0.4680	 0.2530
Q	 0.3750	 0.1990
R	 0.4320	 0.2090
S	 0.4970	 0.2110
T	 0.4770	 0.2020

