



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 11, 2024 – 04:52 PM EST

PDB ID : 2GV5  
Title : crystal structure of Sfi1p/Cdc31p complex  
Authors : Li, S.; Sandercock, A.M.; Conduit, P.T.; Robinson, C.V.; Williams, R.L.; Kilmartin, J.V.  
Deposited on : 2006-05-02  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2511 (3.00-3.00)
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)
RSRZ outliers	164620	2523 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	 4% 76% 16% 7%
1	B	161	 4% 73% 17% 8%
1	D	161	 2% 80% 12% 9%
1	E	161	 2% 73% 17% 8%
2	C	73	 5% 70% 27% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	73	<div><div></div><div>8%</div><div>66%</div><div>33%</div><div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6175 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell division control protein 31.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	149	Total	C	N	O	S	Se	0	0	0
			1227	775	196	250	1	5			
1	B	148	Total	C	N	O	S	Se	0	0	0
			1219	769	195	249	1	5			
1	D	147	Total	C	N	O	S	Se	0	0	0
			1211	765	193	247	1	5			
1	E	148	Total	C	N	O	S	Se	0	0	0
			1219	769	195	249	1	5			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP P06704
A	34	MSE	MET	modified residue	UNP P06704
A	49	MSE	MET	modified residue	UNP P06704
A	76	MSE	MET	modified residue	UNP P06704
A	85	MSE	MET	modified residue	UNP P06704
A	137	MSE	MET	modified residue	UNP P06704
B	1	MSE	MET	modified residue	UNP P06704
B	34	MSE	MET	modified residue	UNP P06704
B	49	MSE	MET	modified residue	UNP P06704
B	76	MSE	MET	modified residue	UNP P06704
B	85	MSE	MET	modified residue	UNP P06704
B	137	MSE	MET	modified residue	UNP P06704
D	1	MSE	MET	modified residue	UNP P06704
D	34	MSE	MET	modified residue	UNP P06704
D	49	MSE	MET	modified residue	UNP P06704
D	76	MSE	MET	modified residue	UNP P06704
D	85	MSE	MET	modified residue	UNP P06704
D	137	MSE	MET	modified residue	UNP P06704
E	1	MSE	MET	modified residue	UNP P06704
E	34	MSE	MET	modified residue	UNP P06704
E	49	MSE	MET	modified residue	UNP P06704

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	76	MSE	MET	modified residue	UNP P06704
E	85	MSE	MET	modified residue	UNP P06704
E	137	MSE	MET	modified residue	UNP P06704

- Molecule 2 is a protein called Sfi1p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	73	Total	C	N	O	S	Se	0	0	0
			623	396	111	113	2	1			
2	F	73	Total	C	N	O	S	Se	0	0	0
			623	396	111	113	2	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	638	GLY	-	cloning artifact	UNP Q12369
C	639	PRO	-	cloning artifact	UNP Q12369
C	640	LEU	-	cloning artifact	UNP Q12369
C	641	GLY	-	cloning artifact	UNP Q12369
C	642	SER	-	cloning artifact	UNP Q12369
C	693	MSE	MET	modified residue	UNP Q12369
F	638	GLY	-	cloning artifact	UNP Q12369
F	639	PRO	-	cloning artifact	UNP Q12369
F	640	LEU	-	cloning artifact	UNP Q12369
F	641	GLY	-	cloning artifact	UNP Q12369
F	642	SER	-	cloning artifact	UNP Q12369
F	693	MSE	MET	modified residue	UNP Q12369

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total	O	0	0
			13	13		
3	B	6	Total	O	0	0
			6	6		
3	C	6	Total	O	0	0
			6	6		
3	D	10	Total	O	0	0
			10	10		
3	E	13	Total	O	0	0
			13	13		

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	5	Total	O	0	0
			5	5		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cell division control protein 31





- Molecule 1: Cell division control protein 31



- Molecule 1: Cell division control protein 31




- Molecule 1: Cell division control protein 31



- Molecule 2: Sfi1p



● Molecule 2: Sfi1p





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.78Å 104.73Å 184.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.45 – 3.00 92.45 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (92.45-3.00) 100.0 (92.45-3.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.48 (at 2.82Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.253 , 0.297 0.238 , 0.279	Depositor DCC
$R_{free}$ test set	1501 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.5	Xtriage
Anisotropy	0.204	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 89.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6175	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.9550e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/1240	0.57	0/1654
1	B	0.36	0/1232	0.56	0/1643
1	D	0.40	0/1224	0.56	0/1632
1	E	0.37	0/1232	0.56	0/1643
2	C	0.47	0/632	0.57	0/841
2	F	0.46	0/632	0.53	0/841
All	All	0.41	0/6192	0.56	0/8254

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1227	0	1182	16	0
1	B	1219	0	1171	15	0
1	D	1211	0	1165	14	0
1	E	1219	0	1171	16	0
2	C	623	0	631	12	0
2	F	623	0	631	13	0
3	A	13	0	0	2	0
3	B	6	0	0	1	0
3	C	6	0	0	0	0
3	D	10	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	13	0	0	1	0
3	F	5	0	0	1	0
All	All	6175	0	5951	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:34:MSE:CE	3:D:165:HOH:O	2.21	0.87
1:D:34:MSE:HA	1:D:34:MSE:HE2	1.65	0.79
1:D:34:MSE:HE1	3:D:165:HOH:O	1.85	0.71
1:A:34:MSE:HA	1:A:34:MSE:HE2	1.72	0.69
1:E:160:ASP:HB2	3:E:166:HOH:O	1.95	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/161 (91%)	139 (95%)	7 (5%)	1 (1%)	19	54
1	B	146/161 (91%)	141 (97%)	5 (3%)	0	100	100
1	D	145/161 (90%)	138 (95%)	6 (4%)	1 (1%)	19	54
1	E	146/161 (91%)	139 (95%)	7 (5%)	0	100	100
2	C	71/73 (97%)	70 (99%)	1 (1%)	0	100	100
2	F	71/73 (97%)	70 (99%)	1 (1%)	0	100	100
All	All	726/790 (92%)	697 (96%)	27 (4%)	2 (0%)	37	70

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	GLY
1	D	72	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/139 (96%)	128 (96%)	6 (4%)	23	57
1	B	133/139 (96%)	119 (90%)	14 (10%)	5	23
1	D	132/139 (95%)	127 (96%)	5 (4%)	28	62
1	E	133/139 (96%)	119 (90%)	14 (10%)	5	23
2	C	69/68 (102%)	64 (93%)	5 (7%)	12	39
2	F	69/68 (102%)	64 (93%)	5 (7%)	12	39
All	All	670/692 (97%)	621 (93%)	49 (7%)	11	39

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	71	GLU
1	E	35	ASN
1	D	73	ARG
1	E	20	GLU
1	E	45	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	665	ASN
2	F	689	GLN
2	C	665	ASN
2	C	689	GLN
1	D	152	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	144/161 (89%)	0.28	6 (4%)	41	24	64, 70, 76, 84	0
1	B	143/161 (88%)	0.66	7 (4%)	36	20	67, 70, 74, 80	0
1	D	142/161 (88%)	0.28	4 (2%)	55	33	64, 70, 75, 79	0
1	E	143/161 (88%)	0.54	4 (2%)	55	33	67, 70, 74, 80	0
2	C	72/73 (98%)	0.60	4 (5%)	31	18	63, 71, 81, 88	0
2	F	72/73 (98%)	0.74	6 (8%)	19	10	63, 70, 81, 91	0
All	All	716/790 (90%)	0.49	31 (4%)	40	23	63, 70, 76, 91	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	13	LEU	6.5
2	F	639	PRO	6.3
2	F	640	LEU	5.5
1	A	14	ASN	5.4
1	B	161	SER	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.