



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 23, 2024 – 04:06 AM EDT

PDB ID : 6GVB
Title : Crystal structure of Cutibacterium acnes exo-beta-1,4-mannosidase
Authors : Reichenbach, T.; Divne, C.
Deposited on : 2018-06-20
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	<div> <div>89%</div> <div>6%</div> <div>5%</div> </div>
1	B	416	<div> <div>89%</div> <div>6%</div> <div>6%</div> </div>
1	C	416	<div> <div>88%</div> <div>7%</div> <div>5%</div> </div>
1	D	416	<div> <div>90%</div> <div>5%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called exo-beta-1,4-mannosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	3	0
			3154	2008	546	589	11			
1	B	392	Total	C	N	O	S	0	3	0
			3133	1996	542	584	11			
1	C	394	Total	C	N	O	S	0	2	0
			3141	2002	543	585	11			
1	D	394	Total	C	N	O	S	0	5	0
			3164	2014	549	590	11			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	409	SER	-	expression tag	UNP A0A2B7I7A6
A	410	ALA	-	expression tag	UNP A0A2B7I7A6
A	411	HIS	-	expression tag	UNP A0A2B7I7A6
A	412	HIS	-	expression tag	UNP A0A2B7I7A6
A	413	HIS	-	expression tag	UNP A0A2B7I7A6
A	414	HIS	-	expression tag	UNP A0A2B7I7A6
A	415	HIS	-	expression tag	UNP A0A2B7I7A6
A	416	HIS	-	expression tag	UNP A0A2B7I7A6
B	409	SER	-	expression tag	UNP A0A2B7I7A6
B	410	ALA	-	expression tag	UNP A0A2B7I7A6
B	411	HIS	-	expression tag	UNP A0A2B7I7A6
B	412	HIS	-	expression tag	UNP A0A2B7I7A6
B	413	HIS	-	expression tag	UNP A0A2B7I7A6
B	414	HIS	-	expression tag	UNP A0A2B7I7A6
B	415	HIS	-	expression tag	UNP A0A2B7I7A6
B	416	HIS	-	expression tag	UNP A0A2B7I7A6
C	409	SER	-	expression tag	UNP A0A2B7I7A6
C	410	ALA	-	expression tag	UNP A0A2B7I7A6
C	411	HIS	-	expression tag	UNP A0A2B7I7A6
C	412	HIS	-	expression tag	UNP A0A2B7I7A6
C	413	HIS	-	expression tag	UNP A0A2B7I7A6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	414	HIS	-	expression tag	UNP A0A2B7I7A6
C	415	HIS	-	expression tag	UNP A0A2B7I7A6
C	416	HIS	-	expression tag	UNP A0A2B7I7A6
D	409	SER	-	expression tag	UNP A0A2B7I7A6
D	410	ALA	-	expression tag	UNP A0A2B7I7A6
D	411	HIS	-	expression tag	UNP A0A2B7I7A6
D	412	HIS	-	expression tag	UNP A0A2B7I7A6
D	413	HIS	-	expression tag	UNP A0A2B7I7A6
D	414	HIS	-	expression tag	UNP A0A2B7I7A6
D	415	HIS	-	expression tag	UNP A0A2B7I7A6
D	416	HIS	-	expression tag	UNP A0A2B7I7A6

- Molecule 2 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (three-letter code: MPO) (formula: C₇H₁₅NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
2	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
2	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total 13	C 7	N 1	O 4	S 1	0	0
2	C	1	Total 13	C 7	N 1	O 4	S 1	0	0
2	C	1	Total 13	C 7	N 1	O 4	S 1	0	0
2	C	1	Total 13	C 7	N 1	O 4	S 1	0	0
2	D	1	Total 13	C 7	N 1	O 4	S 1	0	0
2	D	1	Total 13	C 7	N 1	O 4	S 1	0	0
2	D	1	Total 13	C 7	N 1	O 4	S 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	207	Total 207	O 207	0	0
3	B	217	Total 217	O 217	0	0
3	C	182	Total 182	O 182	0	0
3	D	226	Total 226	O 226	0	0

- Molecule 1: exo-beta-1,4-mannosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.18Å 104.37Å 112.75Å 90.00° 90.27° 90.00°	Depositor
Resolution (Å)	49.60 – 1.80 49.60 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.5 (49.60-1.80) 90.5 (49.60-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.91 (at 1.79Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.195 , 0.227 0.196 , 0.227	Depositor DCC
R_{free} test set	2001 reflections (1.14%)	wwPDB-VP
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.541	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.056 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13580	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.10 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6015e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/3261	0.52	0/4472
1	B	0.39	0/3242	0.55	0/4445
1	C	0.38	0/3251	0.54	0/4459
1	D	0.39	0/3274	0.54	0/4490
All	All	0.38	0/13028	0.54	0/17866

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3154	0	2972	14	0
1	B	3133	0	2956	16	0
1	C	3141	0	2966	18	0
1	D	3164	0	2987	9	0
2	A	39	0	45	4	0
2	B	39	0	45	5	0
2	C	39	0	45	8	0
2	D	39	0	45	4	0
3	A	207	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	217	0	0	1	0
3	C	182	0	0	3	0
3	D	226	0	0	2	0
All	All	13580	0	12061	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:86:HIS:NE2	2:D:502:MPO:H11	1.99	0.78
1:A:370:ASP:O	1:A:374:GLN:HG3	1.84	0.77
1:C:86:HIS:NE2	2:C:502:MPO:H11	2.01	0.75
1:B:147:LEU:HD13	1:B:197:ASP:OD1	1.85	0.75
1:B:89:SER:HA	2:B:502:MPO:H31	1.72	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/416 (95%)	387 (98%)	9 (2%)	0	100	100
1	B	393/416 (94%)	383 (98%)	10 (2%)	0	100	100
1	C	394/416 (95%)	386 (98%)	8 (2%)	0	100	100
1	D	397/416 (95%)	387 (98%)	10 (2%)	0	100	100
All	All	1580/1664 (95%)	1543 (98%)	37 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/350 (96%)	333 (99%)	4 (1%)	71	65
1	B	335/350 (96%)	331 (99%)	4 (1%)	71	65
1	C	336/350 (96%)	333 (99%)	3 (1%)	78	75
1	D	339/350 (97%)	331 (98%)	8 (2%)	49	36
All	All	1347/1400 (96%)	1328 (99%)	19 (1%)	71	59

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	147	LEU
1	D	331[B]	SER
1	D	344	HIS
1	D	331[A]	SER
1	C	81	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	374	GLN
1	C	344	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MPO	B	503	-	13,13,13	1.36	3 (23%)	17,17,17	2.00	6 (35%)
2	MPO	D	502	-	13,13,13	1.54	3 (23%)	17,17,17	2.27	5 (29%)
2	MPO	B	501	-	13,13,13	1.47	3 (23%)	17,17,17	2.39	6 (35%)
2	MPO	C	503	-	13,13,13	1.44	3 (23%)	17,17,17	2.32	9 (52%)
2	MPO	D	503	-	13,13,13	1.41	3 (23%)	17,17,17	2.42	9 (52%)
2	MPO	A	502	-	13,13,13	1.40	3 (23%)	17,17,17	2.60	6 (35%)
2	MPO	A	501	-	13,13,13	1.51	3 (23%)	17,17,17	2.14	7 (41%)
2	MPO	A	503	-	13,13,13	1.43	3 (23%)	17,17,17	2.37	9 (52%)
2	MPO	C	502	-	13,13,13	1.57	3 (23%)	17,17,17	2.29	6 (35%)
2	MPO	D	501	-	13,13,13	1.50	3 (23%)	17,17,17	2.27	7 (41%)
2	MPO	B	502	-	13,13,13	1.57	3 (23%)	17,17,17	2.11	6 (35%)
2	MPO	C	501	-	13,13,13	1.53	3 (23%)	17,17,17	2.33	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPO	B	503	-	-	3/7/15/15	0/1/1/1
2	MPO	D	502	-	-	4/7/15/15	0/1/1/1
2	MPO	B	501	-	-	3/7/15/15	0/1/1/1
2	MPO	C	503	-	-	1/7/15/15	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPO	D	503	-	-	1/7/15/15	0/1/1/1
2	MPO	A	502	-	-	4/7/15/15	0/1/1/1
2	MPO	A	501	-	-	5/7/15/15	0/1/1/1
2	MPO	A	503	-	-	2/7/15/15	0/1/1/1
2	MPO	C	502	-	-	5/7/15/15	0/1/1/1
2	MPO	D	501	-	-	4/7/15/15	0/1/1/1
2	MPO	B	502	-	-	3/7/15/15	0/1/1/1
2	MPO	C	501	-	-	3/7/15/15	0/1/1/1

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	502	MPO	C1-S1	3.58	1.82	1.77
2	B	502	MPO	C1-S1	3.41	1.82	1.77
2	C	501	MPO	C1-S1	3.33	1.82	1.77
2	D	501	MPO	C1-S1	3.18	1.82	1.77
2	C	503	MPO	C1-S1	3.14	1.82	1.77

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	MPO	O2-S1-C1	5.98	115.76	106.73
2	D	502	MPO	C7-N1-C4	5.60	120.91	108.84
2	C	502	MPO	C7-N1-C4	5.49	120.67	108.84
2	A	502	MPO	C7-N1-C4	5.25	120.16	108.84
2	A	503	MPO	C6-C7-N1	4.68	117.23	110.12

There are no chirality outliers.

5 of 38 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	MPO	C2-C1-S1-O1
2	A	501	MPO	C2-C1-S1-O2
2	A	502	MPO	C2-C1-S1-O1
2	B	501	MPO	C2-C1-S1-O1
2	C	501	MPO	C2-C1-S1-O1

There are no ring outliers.

10 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	503	MPO	2	0
2	D	502	MPO	3	0
2	B	501	MPO	1	0
2	C	503	MPO	2	0
2	D	503	MPO	1	0
2	A	502	MPO	2	0
2	A	501	MPO	2	0
2	C	502	MPO	5	0
2	B	502	MPO	2	0
2	C	501	MPO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/416 (94%)	-0.21	6 (1%) 73 70	22, 33, 50, 76	0
1	B	392/416 (94%)	-0.33	3 (0%) 86 84	21, 31, 46, 70	0
1	C	394/416 (94%)	-0.21	3 (0%) 86 84	21, 33, 51, 70	0
1	D	394/416 (94%)	-0.22	5 (1%) 77 74	20, 30, 47, 65	0
All	All	1575/1664 (94%)	-0.24	17 (1%) 80 78	20, 32, 49, 76	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	393	THR	4.9
1	A	393	THR	4.0
1	A	394	PRO	3.9
1	B	392	PRO	3.9
1	A	395	GLY	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MPO	B	503	13/13	0.73	0.24	45,50,80,81	13
2	MPO	C	502	13/13	0.74	0.22	36,43,52,64	13
2	MPO	D	502	13/13	0.78	0.22	28,39,51,54	13
2	MPO	D	503	13/13	0.78	0.20	38,42,64,68	13
2	MPO	B	502	13/13	0.81	0.21	35,42,55,59	13
2	MPO	C	503	13/13	0.81	0.25	44,51,81,83	13
2	MPO	A	502	13/13	0.83	0.20	32,42,49,58	13
2	MPO	A	503	13/13	0.83	0.19	42,46,71,73	13
2	MPO	C	501	13/13	0.93	0.16	35,41,52,56	13
2	MPO	A	501	13/13	0.94	0.11	37,40,50,55	13
2	MPO	D	501	13/13	0.94	0.10	28,36,49,49	13
2	MPO	B	501	13/13	0.95	0.13	31,35,48,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.