



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 10:06 AM EDT

PDB ID : 6GWV  
Title : Molybdenum storage protein without polymolybdate clusters and ATP  
Authors : Ermler, U.; Poppe, J.; Bruenle, S.  
Deposited on : 2018-06-26  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	269	
1	C	269	
1	E	269	
1	H	269	
1	I	269	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	269	
1	N	269	
1	O	269	
1	Q	269	
2	A	275	
2	D	275	
2	F	275	
2	G	275	
2	J	275	
2	L	275	
2	M	275	
2	P	275	
2	R	275	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	E	301	-	-	X	-
5	CL	M	302	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 69375 atoms, of which 35057 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Molybdenum storage protein subunit beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	268	Total	C	H	N	O	S	0	0	0
			4012	1255	2033	347	369	8			
1	C	268	Total	C	H	N	O	S	0	0	0
			4012	1255	2033	347	369	8			
1	E	268	Total	C	H	N	O	S	0	1	0
			4027	1260	2043	347	369	8			
1	H	268	Total	C	H	N	O	S	0	0	0
			4010	1255	2031	347	369	8			
1	I	268	Total	C	H	N	O	S	0	0	0
			4011	1255	2032	347	369	8			
1	K	268	Total	C	H	N	O	S	0	0	0
			4011	1255	2032	347	369	8			
1	N	268	Total	C	H	N	O	S	0	0	0
			4011	1255	2032	347	369	8			
1	O	268	Total	C	H	N	O	S	0	0	0
			4012	1255	2033	347	369	8			
1	Q	268	Total	C	H	N	O	S	0	0	0
			4012	1255	2033	347	369	8			

- Molecule 2 is a protein called Molybdenum storage protein subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	A	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			
2	D	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			
2	F	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			
2	G	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			
2	J	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			
2	M	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			
2	P	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			
2	R	244	Total	C	H	N	O	S	0	0	0
			3680	1152	1859	339	327	3			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	N	1	Total	O	S	0	0
			5	4	1		
3	N	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	M	1	Total	O	S	0	0
			5	4	1		
3	Q	1	Total	O	S	0	0
			5	4	1		
3	R	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	0	0
			43	10	12	5	13	3		
4	P	1	Total	C	H	N	O	P	0	0
			43	10	12	5	13	3		


- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	M	1	Total	Cl	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Molybdenum storage protein subunit beta




- Molecule 1: Molybdenum storage protein subunit beta




- Molecule 1: Molybdenum storage protein subunit beta








• Molecule 1: Molybdenum storage protein subunit beta




• Molecule 1: Molybdenum storage protein subunit beta




• Molecule 1: Molybdenum storage protein subunit beta









● Molecule 1: Molybdenum storage protein subunit beta



● Molecule 1: Molybdenum storage protein subunit beta




● Molecule 1: Molybdenum storage protein subunit beta




- Molecule 2: Molybdenum storage protein subunit alpha

Chain A: 




- Molecule 2: Molybdenum storage protein subunit alpha

Chain D: 




- Molecule 2: Molybdenum storage protein subunit alpha


Chain F: 




- Molecule 2: Molybdenum storage protein subunit alpha

Chain G: 






• Molecule 2: Molybdenum storage protein subunit alpha




• Molecule 2: Molybdenum storage protein subunit alpha




• Molecule 2: Molybdenum storage protein subunit alpha






• Molecule 2: Molybdenum storage protein subunit alpha



• Molecule 2: Molybdenum storage protein subunit alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.71Å 147.30Å 192.32Å 90.00° 107.62° 90.00°	Depositor
Resolution (Å)	47.79 – 2.80 47.79 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.3 (47.79-2.80) 97.1 (47.79-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 2.81Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.233 , 0.259 0.237 , 0.252	Depositor DCC
$R_{free}$ test set	7465 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.6	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 29.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.36$ , $\langle L^2 \rangle = 0.19$	Xtriage
Estimated twinning fraction	0.368 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	69375	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ATP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.34	0/2013	0.60	0/2735
1	C	0.32	0/2013	0.63	2/2735 (0.1%)
1	E	0.35	0/2021	0.61	0/2746
1	H	0.68	5/2013 (0.2%)	0.93	12/2735 (0.4%)
1	I	0.54	3/2013 (0.1%)	0.76	5/2735 (0.2%)
1	K	0.36	0/2013	0.68	1/2735 (0.0%)
1	N	0.38	0/2013	0.66	2/2735 (0.1%)
1	O	0.33	0/2013	0.62	0/2735
1	Q	0.35	0/2013	0.59	0/2735
2	A	0.29	0/1861	0.53	0/2538
2	D	0.31	0/1861	0.52	0/2538
2	F	0.29	0/1861	0.54	0/2538
2	G	0.30	0/1861	0.58	0/2538
2	J	0.29	0/1861	0.60	0/2538
2	L	0.42	1/1861 (0.1%)	0.68	3/2538 (0.1%)
2	M	0.32	0/1861	0.58	2/2538 (0.1%)
2	P	0.28	0/1861	0.50	0/2538
2	R	0.29	0/1861	0.53	0/2538
All	All	0.37	9/34874 (0.0%)	0.63	27/47468 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3
1	C	0	4
1	E	0	2
1	H	0	6
1	I	0	3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	#Chirality outliers	#Planarity outliers
1	K	0	2
1	N	0	2
1	O	0	2
1	Q	0	3
2	A	0	1
2	D	0	3
2	F	0	1
2	G	0	1
2	J	0	1
2	L	0	1
2	M	0	1
2	P	0	1
2	R	0	3
All	All	0	40

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	219	LYS	CB-CG	14.47	1.91	1.52
1	H	220	GLY	CA-C	12.75	1.72	1.51
1	I	187	PHE	CE1-CZ	-11.75	1.15	1.37
1	I	187	PHE	CG-CD1	-9.89	1.24	1.38
1	H	220	GLY	N-CA	8.79	1.59	1.46

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	219	LYS	C-N-CA	-16.30	88.08	122.30
2	L	205	ARG	NE-CZ-NH2	-11.22	114.69	120.30
1	H	219	LYS	N-CA-C	-10.58	82.44	111.00
2	L	205	ARG	NE-CZ-NH1	9.97	125.29	120.30
2	M	205	ARG	NE-CZ-NH2	-9.33	115.63	120.30

There are no chirality outliers.

5 of 40 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	155	TYR	Peptide
1	B	152	TYR	Peptide
1	B	204	ASP	Peptide
1	B	4	SER	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	C	6	ALA	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1979	2033	2032	84	2
1	C	1979	2033	2032	89	0
1	E	1984	2043	2043	82	0
1	H	1979	2031	2031	179	0
1	I	1979	2032	2032	160	0
1	K	1979	2032	2032	149	0
1	N	1979	2032	2032	89	0
1	O	1979	2033	2032	88	2
1	Q	1979	2033	2032	83	0
2	A	1821	1859	1859	45	0
2	D	1821	1859	1859	55	0
2	F	1821	1859	1859	39	0
2	G	1821	1859	1859	121	0
2	J	1821	1859	1859	95	0
2	L	1821	1859	1859	75	0
2	M	1821	1859	1859	38	0
2	P	1821	1859	1859	41	0
2	R	1821	1859	1859	49	0
3	B	5	0	0	1	0
3	D	5	0	0	0	0
3	E	5	0	0	2	0
3	F	5	0	0	0	0
3	G	5	0	0	0	0
3	M	5	0	0	0	0
3	N	10	0	0	1	0
3	Q	5	0	0	0	0
3	R	5	0	0	0	0
4	A	31	12	11	3	0
4	P	31	12	12	5	0
5	M	1	0	0	2	0
All	All	34318	35057	35052	1377	2



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 1377 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:219:LYS:CG	1:H:219:LYS:CB	1.91	1.45
1:O:187:PHE:CE1	1:O:243:VAL:HG11	1.84	1.12
1:N:13:MET:O	2:M:49:ARG:NH2	1.89	1.06
2:J:224:GLY:O	2:J:230:ARG:NH2	1.89	1.05
1:H:219:LYS:CB	1:H:221:LEU:HD22	1.87	1.04

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:200:LYS:NZ	1:O:204:ASP:OD2[2_354]	2.07	0.13
1:B:200:LYS:HZ1	1:O:204:ASP:OD2[2_354]	1.56	0.04

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	266/269 (99%)	260 (98%)	6 (2%)	0	100	100
1	C	266/269 (99%)	253 (95%)	13 (5%)	0	100	100
1	E	267/269 (99%)	263 (98%)	4 (2%)	0	100	100
1	H	266/269 (99%)	255 (96%)	10 (4%)	1 (0%)	34	66
1	I	266/269 (99%)	252 (95%)	14 (5%)	0	100	100
1	K	266/269 (99%)	255 (96%)	11 (4%)	0	100	100
1	N	266/269 (99%)	257 (97%)	9 (3%)	0	100	100
1	O	266/269 (99%)	260 (98%)	6 (2%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	266/269 (99%)	261 (98%)	5 (2%)	0	100	100
2	A	242/275 (88%)	240 (99%)	2 (1%)	0	100	100
2	D	242/275 (88%)	239 (99%)	3 (1%)	0	100	100
2	F	242/275 (88%)	238 (98%)	4 (2%)	0	100	100
2	G	242/275 (88%)	241 (100%)	1 (0%)	0	100	100
2	J	242/275 (88%)	239 (99%)	3 (1%)	0	100	100
2	L	242/275 (88%)	238 (98%)	4 (2%)	0	100	100
2	M	242/275 (88%)	239 (99%)	3 (1%)	0	100	100
2	P	242/275 (88%)	241 (100%)	1 (0%)	0	100	100
2	R	242/275 (88%)	233 (96%)	9 (4%)	0	100	100
All	All	4573/4896 (93%)	4464 (98%)	108 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	220	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	205/205 (100%)	199 (97%)	6 (3%)	42	76
1	C	205/205 (100%)	190 (93%)	15 (7%)	14	38
1	E	206/205 (100%)	198 (96%)	8 (4%)	32	66
1	H	205/205 (100%)	188 (92%)	17 (8%)	11	32
1	I	205/205 (100%)	187 (91%)	18 (9%)	10	29
1	K	205/205 (100%)	189 (92%)	16 (8%)	12	35
1	N	205/205 (100%)	198 (97%)	7 (3%)	37	71
1	O	205/205 (100%)	194 (95%)	11 (5%)	22	53

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Q	205/205 (100%)	200 (98%)	5 (2%)	49	81
2	A	187/215 (87%)	182 (97%)	5 (3%)	44	78
2	D	187/215 (87%)	183 (98%)	4 (2%)	53	84
2	F	187/215 (87%)	184 (98%)	3 (2%)	62	88
2	G	187/215 (87%)	177 (95%)	10 (5%)	22	54
2	J	187/215 (87%)	177 (95%)	10 (5%)	22	54
2	L	187/215 (87%)	178 (95%)	9 (5%)	25	58
2	M	187/215 (87%)	177 (95%)	10 (5%)	22	54
2	P	187/215 (87%)	183 (98%)	4 (2%)	53	84
2	R	187/215 (87%)	180 (96%)	7 (4%)	34	68
All	All	3529/3780 (93%)	3364 (95%)	165 (5%)	26	59

5 of 165 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	139	ILE
1	O	137	SER
2	L	209	ARG
2	M	67	LYS
2	P	144	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	G	158	HIS
2	J	241	HIS
1	K	222	HIS
1	O	222	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	N	302	-	4,4,4	0.21	0	6,6,6	0.13	0
3	SO4	E	301	-	4,4,4	0.24	0	6,6,6	0.12	0
3	SO4	D	301	-	4,4,4	0.23	0	6,6,6	0.07	0
3	SO4	N	301	-	4,4,4	0.22	0	6,6,6	0.09	0
4	ATP	A	301	-	28,33,33	3.39	14 (50%)	34,52,52	3.24	5 (14%)
3	SO4	M	301	-	4,4,4	0.23	0	6,6,6	0.09	0
3	SO4	G	301	-	4,4,4	0.21	0	6,6,6	0.10	0
3	SO4	R	301	-	4,4,4	0.24	0	6,6,6	0.09	0
3	SO4	B	301	-	4,4,4	0.22	0	6,6,6	0.11	0
3	SO4	Q	301	-	4,4,4	0.24	0	6,6,6	0.11	0
3	SO4	F	301	-	4,4,4	0.21	0	6,6,6	0.12	0
4	ATP	P	301	-	28,33,33	3.36	14 (50%)	34,52,52	3.68	6 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	P	301	-	-	0/18/38/38	0/3/3/3
4	ATP	A	301	-	-	8/18/38/38	0/3/3/3

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	ATP	C2'-C3'	-10.54	1.24	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	P	301	ATP	C2'-C3'	-9.92	1.26	1.53
4	A	301	ATP	PB-O3A	6.34	1.66	1.59
4	A	301	ATP	PA-O3A	6.30	1.66	1.59
4	P	301	ATP	PA-O3A	6.30	1.66	1.59

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	301	ATP	C5-C6-N6	14.00	141.64	120.31
4	A	301	ATP	C5-C6-N6	13.50	140.88	120.31
4	P	301	ATP	N6-C6-N1	-9.52	97.98	118.33
4	A	301	ATP	N6-C6-N1	-9.21	98.65	118.33
4	P	301	ATP	C4'-O4'-C1'	-9.02	101.66	109.92

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	301	ATP	C5'-O5'-PA-O1A
4	A	301	ATP	C5'-O5'-PA-O2A
4	A	301	ATP	C5'-O5'-PA-O3A
4	A	301	ATP	O4'-C4'-C5'-O5'
4	A	301	ATP	C3'-C4'-C5'-O5'


There are no ring outliers.


5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	N	302	SO4	1	0
3	E	301	SO4	2	0
4	A	301	ATP	3	0
3	B	301	SO4	1	0
4	P	301	ATP	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	268/269 (99%)	0.15	1 (0%) 92 91	20, 34, 58, 73	0
1	C	268/269 (99%)	0.38	9 (3%) 45 35	26, 44, 73, 88	0
1	E	268/269 (99%)	0.28	6 (2%) 62 52	22, 39, 67, 84	0
1	H	268/269 (99%)	1.17	52 (19%) 1 0	58, 81, 102, 108	0
1	I	268/269 (99%)	1.18	56 (20%) 1 0	60, 79, 96, 111	0
1	K	268/269 (99%)	0.95	47 (17%) 1 1	49, 72, 90, 99	0
1	N	268/269 (99%)	0.22	4 (1%) 73 68	17, 33, 61, 81	0
1	O	268/269 (99%)	0.19	6 (2%) 62 52	22, 39, 63, 78	0
1	Q	268/269 (99%)	0.23	7 (2%) 56 46	17, 33, 64, 83	0
2	A	244/275 (88%)	0.06	0 100 100	13, 22, 43, 55	0
2	D	244/275 (88%)	0.15	3 (1%) 79 73	14, 30, 63, 78	0
2	F	244/275 (88%)	0.14	1 (0%) 92 91	14, 29, 55, 74	0
2	G	244/275 (88%)	0.78	32 (13%) 3 2	48, 65, 83, 92	0
2	J	244/275 (88%)	0.92	34 (13%) 2 1	41, 65, 90, 99	0
2	L	244/275 (88%)	0.80	28 (11%) 4 2	43, 63, 83, 99	0
2	M	244/275 (88%)	0.09	0 100 100	13, 25, 49, 65	0
2	P	244/275 (88%)	0.11	1 (0%) 92 91	13, 24, 46, 62	0
2	R	244/275 (88%)	0.10	2 (0%) 86 81	13, 27, 53, 67	0
All	All	4608/4896 (94%)	0.44	289 (6%) 20 12	13, 42, 85, 111	0

The worst 5 of 289 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	5	THR	7.3
1	K	107	ALA	6.9
2	G	272	GLY	6.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	I	163	VAL	6.7
1	K	199	PRO	6.5

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	E	301	5/5	0.80	0.32	102,106,113,114	0
3	SO4	F	301	5/5	0.83	0.28	88,92,93,99	0
3	SO4	B	301	5/5	0.84	0.26	92,93,98,99	0
3	SO4	D	301	5/5	0.87	0.30	144,145,149,151	0
4	ATP	P	301	31/31	0.90	0.27	67,129,164,168	0
3	SO4	G	301	5/5	0.91	0.20	112,117,118,119	0
3	SO4	N	302	5/5	0.93	0.20	38,38,38,38	0
3	SO4	M	301	5/5	0.94	0.26	67,72,75,90	0
4	ATP	A	301	31/31	0.96	0.18	33,41,53,54	0
3	SO4	Q	301	5/5	0.96	0.29	97,103,105,106	0
3	SO4	N	301	5/5	0.97	0.20	64,73,79,80	0
3	SO4	R	301	5/5	0.97	0.18	44,44,44,44	0
5	CL	M	302	1/1	0.97	0.22	38,38,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ATP P 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ATP A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.