



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2025 – 01:37 pm BST

PDB ID : 9GWG / pdb_00009gwg
Title : Human PPAR-gamma ligand binding domain in complex with LW88
Authors : Oerlemans, G.J.M.; Koops, A.A.; Brunsveld, L.
Deposited on : 2024-09-26
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>9%</div> <div>7%</div> </div> </div>
1	B	275	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>5%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			1977	1281	321	368	7			
1	B	262	Total	C	N	O	S	0	0	0
			2011	1295	326	380	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	231	GLY	-	expression tag	UNP P37231
A	232	SER	-	expression tag	UNP P37231
A	233	HIS	-	expression tag	UNP P37231
A	234	MET	-	expression tag	UNP P37231
B	231	GLY	-	expression tag	UNP P37231
B	232	SER	-	expression tag	UNP P37231
B	233	HIS	-	expression tag	UNP P37231
B	234	MET	-	expression tag	UNP P37231

- Molecule 2 is 4-[5-chloranyl-1-[2-chloranyl-6-(trifluoromethyl)phenyl]carbonyl-indazol-3-yl]-3-[[1-[(2-methylpropan-2-yl)oxycarbonyl]piperidin-4-yl]methoxy]benzoic acid (CCD ID: A1IP1) (formula: C₃₃H₃₀Cl₂F₃N₃O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	Cl	F	N	O	0	0
			47	33	2	3	3	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	6	Total	O	0
			6	6	
3	B	7	Total	O	0
			7	7	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.70Å 61.62Å 119.24Å 90.00° 102.05° 90.00°	Depositor
Resolution (Å)	46.04 – 2.30 46.04 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.04-2.30) 99.6 (46.04-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487, PDB-REDO	Depositor
R, R_{free}	0.220 , 0.252 0.220 , 0.252	Depositor DCC
R_{free} test set	1517 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	50.0	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4048	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1IP1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2010	0.58	0/2724
1	B	0.39	0/2044	0.57	0/2768
All	All	0.39	0/4054	0.57	0/5492

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1977	0	1958	12	0
1	B	2011	0	1947	23	0
2	B	47	0	0	2	0
3	A	6	0	0	0	0
3	B	7	0	0	1	0
All	All	4048	0	3905	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:LYS:NZ	1:A:502:LYS:O	2.22	0.72
1:B:287:GLU:OE1	1:B:308:ARG:NH2	2.26	0.68
1:A:381:LEU:HD22	1:A:392:MET:HG3	1.76	0.68
1:B:353:ILE:HG23	1:B:416:ILE:HD12	1.81	0.62
1:B:472:GLN:O	1:B:476:GLU:HG3	1.99	0.62
1:B:316:ARG:HE	2:B:601:A1IP1:C13	2.13	0.61
1:A:355:TYR:CE2	1:A:395:LYS:HG3	2.45	0.52
1:B:355:TYR:OH	1:B:395:LYS:HE3	2.11	0.51
1:B:304:GLU:HB3	1:B:307:ILE:HD12	1.92	0.50
1:B:494:HIS:HD2	1:B:496:LEU:H	1.61	0.48
1:A:353:ILE:HG23	1:A:416:ILE:HD12	1.96	0.48
1:A:412:LEU:O	1:A:416:ILE:HG12	2.14	0.47
1:B:275:PHE:CZ	1:B:285:MET:HG2	2.49	0.47
1:A:254:PHE:CG	1:A:357:MET:HE1	2.50	0.46
1:B:235:GLU:N	3:B:701:HOH:O	2.49	0.45
1:A:304:GLU:HG3	1:A:305:VAL:N	2.32	0.44
1:A:267:GLY:C	1:A:269:THR:H	2.25	0.44
1:B:280:MET:SD	1:B:305:VAL:HG11	2.58	0.44
1:B:461:ALA:O	1:B:465:GLN:HG3	2.18	0.44
1:A:351:HIS:CE1	1:A:501:TYR:OH	2.71	0.43
1:B:385:ARG:HH11	1:B:386:LYS:CB	2.32	0.43
1:A:350:VAL:HG21	1:A:500:ILE:HD13	2.01	0.42
1:B:265:LEU:HD21	1:B:368:LEU:HG	2.00	0.42
1:B:497:LEU:HD23	1:B:497:LEU:HA	1.65	0.42
1:B:316:ARG:HG2	2:B:601:A1IP1:C11	2.49	0.42
1:B:304:GLU:OE2	1:B:385:ARG:NE	2.38	0.42
1:B:386:LYS:CB	1:B:387:PRO:HD3	2.50	0.42
1:A:477:HIS:NE2	1:A:481:LEU:HD11	2.34	0.42
1:B:376:MET:HE3	1:B:376:MET:HB2	1.75	0.41
1:B:386:LYS:CB	1:B:387:PRO:CD	2.99	0.41
1:B:254:PHE:CG	1:B:357:MET:HE1	2.55	0.41
1:A:386:LYS:CB	1:A:387:PRO:HD3	2.51	0.41
1:B:355:TYR:CE2	1:B:395:LYS:HG3	2.56	0.41
1:B:488:GLU:O	1:B:491:MET:HG2	2.21	0.41
1:B:355:TYR:OH	1:B:477:HIS:HB2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/275 (92%)	250 (99%)	2 (1%)	1 (0%)	30	39
1	B	256/275 (93%)	250 (98%)	5 (2%)	1 (0%)	30	39
All	All	509/550 (92%)	500 (98%)	7 (1%)	2 (0%)	30	39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	386	LYS
1	B	386	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/247 (83%)	203 (98%)	3 (2%)	60	76
1	B	208/247 (84%)	207 (100%)	1 (0%)	86	93
All	All	414/494 (84%)	410 (99%)	4 (1%)	73	85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	400	VAL
1	A	462	LYS
1	A	486	LYS
1	B	496	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	351	HIS
1	A	465	GLN
1	A	472	GLN
1	B	438	GLN
1	B	443	GLN
1	B	494	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A1IP1	B	601	-	47,51,51	0.73	1 (2%)	66,77,77	1.00	6 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1IP1	B	601	-	-	5/30/46/46	0/5/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	A1IP1	C17-N2	-3.09	1.32	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	A1IP1	C31-C26-N3	-2.75	105.30	109.08
2	B	601	A1IP1	C19-C18-N3	2.31	118.77	116.14
2	B	601	A1IP1	O6-C18-C19	2.26	124.11	119.44
2	B	601	A1IP1	C20-C19-C18	-2.24	118.50	121.19
2	B	601	A1IP1	C15-C16-C17	-2.22	117.39	120.14
2	B	601	A1IP1	O3-C10-C11	-2.04	118.89	123.58

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	A1IP1	C7-C8-C9-O3
2	B	601	A1IP1	C8-C9-O3-C10
2	B	601	A1IP1	C32-C8-C9-O3
2	B	601	A1IP1	C11-C10-O3-C9
2	B	601	A1IP1	C16-C10-O3-C9


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	A1IP1	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/275 (93%)	0.25	14 (5%) 32 34	43, 60, 88, 105	0
1	B	262/275 (95%)	0.24	12 (4%) 38 39	37, 57, 96, 116	0
All	All	519/550 (94%)	0.24	26 (5%) 35 36	37, 59, 90, 116	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	504	LEU	4.2
1	B	294	HIS	3.7
1	B	292	PHE	3.7
1	B	293	LYS	3.1
1	A	504	LEU	3.0
1	A	268	LYS	2.9
1	B	295	ILE	2.9
1	A	355	TYR	2.7
1	B	290	ILE	2.7
1	A	493	LEU	2.6
1	A	392	MET	2.6
1	B	501	TYR	2.6
1	A	270	THR	2.6
1	B	288	ASP	2.6
1	B	296	THR	2.5
1	A	503	ASP	2.5
1	B	489	THR	2.4
1	A	269	THR	2.3
1	A	489	THR	2.3
1	B	503	ASP	2.3
1	A	315	PHE	2.2
1	A	490	ASP	2.2
1	A	391	PHE	2.1
1	A	281	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	494	HIS	2.0
1	A	288	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	A1IP1	B	601	47/47	0.90	0.12	60,64,77,82	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1IP1 B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.