



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 06:11 AM EDT

PDB ID : 6GY0
Title : mPI3Kd IN COMPLEX WITH AZ3
Authors : Petersen, J.
Deposited on : 2018-06-27
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	938	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6687 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit delta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	822	Total	C	N	O	S	0	0	0
			6633	4254	1126	1199	54			

- Molecule 2 is {N}-[4-methyl-5-(1-oxidanylidene-7-sulfamoyl-isoindol-5-yl)-1,3-thiazol-2-yl]ethanamide (three-letter code: FGE) (formula: C₁₄H₁₂N₄O₄S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			24	14	4	4	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	30	Total 30 O 30	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.91Å 63.70Å 115.87Å 90.00° 103.61° 90.00°	Depositor
Resolution (Å)	42.48 – 2.55 42.48 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.3 (42.48-2.55) 99.3 (42.48-2.55)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.251 , 0.307 0.252 , 0.305	Depositor DCC
R_{free} test set	1620 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	54.4	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 29.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6687	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/6776	0.61	0/9142

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6633	0	6615	45	0
2	A	24	0	0	1	0
3	A	30	0	0	2	0
All	All	6687	0	6615	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:367:SER:HB3	1:A:368:GLU:CA	2.01	0.90

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:367:SER:HB3	1:A:368:GLU:HB3	1.56	0.86
1:A:367:SER:HB3	1:A:368:GLU:CB	2.08	0.84
1:A:367:SER:HB3	1:A:368:GLU:C	1.99	0.81
1:A:367:SER:CB	1:A:368:GLU:HB3	2.11	0.79
1:A:366:CYS:HB3	3:A:1216:HOH:O	1.84	0.76
1:A:610:GLN:HE22	1:A:795:GLN:HE21	1.32	0.75
1:A:364:ASN:O	1:A:366:CYS:N	2.20	0.73
1:A:971:GLY:HA3	1:A:1004:LEU:HD21	1.77	0.66
1:A:325:ILE:HD11	1:A:375:LEU:HD12	1.78	0.65
1:A:365:VAL:HG12	1:A:365:VAL:O	2.01	0.59
1:A:367:SER:OG	1:A:368:GLU:HB3	2.03	0.59
1:A:808:LEU:HD11	1:A:963:ALA:HB2	1.84	0.57
1:A:955:ARG:NH2	1:A:959:TYR:OH	2.41	0.54
1:A:247:HIS:HB2	1:A:738:SER:HA	1.91	0.53
1:A:367:SER:HB3	1:A:368:GLU:O	2.10	0.51
1:A:741:LEU:HD13	1:A:763:TYR:CE2	2.47	0.50
1:A:383:ASP:OD1	1:A:558:HIS:ND1	2.38	0.50
1:A:634:LEU:HD13	1:A:667:ILE:HG23	1.94	0.49
1:A:366:CYS:O	1:A:369:PRO:HA	2.14	0.48
1:A:379:ILE:HD12	1:A:556:ASN:HB2	1.96	0.48
1:A:109:VAL:HG12	1:A:110:LYS:H	1.79	0.48
1:A:549:LEU:HG	1:A:564:MET:HE3	1.96	0.48
1:A:245:GLY:HA3	1:A:768:ALA:HB2	1.96	0.47
1:A:291:ASN:HD22	1:A:291:ASN:N	2.14	0.46
1:A:366:CYS:CB	3:A:1216:HOH:O	2.55	0.46
1:A:600:LEU:HD22	1:A:603:LEU:HD11	1.97	0.46
1:A:549:LEU:HG	1:A:564:MET:CE	2.46	0.46
1:A:366:CYS:O	1:A:367:SER:HB2	2.15	0.45
1:A:752:MET:SD	2:A:1101:FGE:N17	2.89	0.45
1:A:790:THR:HG21	1:A:912:PHE:CD1	2.52	0.45
1:A:856:LEU:HD11	1:A:872:ILE:HD11	1.99	0.45
1:A:806:LEU:HD11	1:A:808:LEU:HD21	2.00	0.44
1:A:368:GLU:N	1:A:369:PRO:CD	2.80	0.43
1:A:907:LEU:HD23	1:A:908:PHE:N	2.34	0.43
1:A:637:ALA:HB1	1:A:644:GLY:HA2	2.01	0.43
1:A:614:GLN:HG3	1:A:981:MET:HG2	2.01	0.43
1:A:192:VAL:HG11	1:A:216:LEU:HD11	2.00	0.43
1:A:750:THR:OG1	1:A:751:PHE:N	2.53	0.42
1:A:367:SER:CB	1:A:368:GLU:CA	2.84	0.41
1:A:246:ARG:NH1	1:A:248:GLU:OE1	2.53	0.41
1:A:334:ASN:HD22	1:A:341:LEU:HD21	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:CYS:O	1:A:366:CYS:SG	2.79	0.41
1:A:419:ALA:HB3	1:A:441:MET:HE3	2.02	0.41
1:A:366:CYS:O	1:A:367:SER:CB	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	800/938 (85%)	763 (95%)	31 (4%)	6 (1%)	19 27

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	365	VAL
1	A	367	SER
1	A	911	ASP
1	A	841	LYS
1	A	1026	SER
1	A	328	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	728/826 (88%)	692 (95%)	36 (5%)	25	34

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	LYS
1	A	119	LEU
1	A	219	CYS
1	A	263	CYS
1	A	270	LEU
1	A	281	SER
1	A	291	ASN
1	A	316	LEU
1	A	317	TRP
1	A	318	SER
1	A	319	LEU
1	A	331	ARG
1	A	332	LYS
1	A	340	LYS
1	A	356	LYS
1	A	367	SER
1	A	423	LEU
1	A	437	ARG
1	A	472	VAL
1	A	489	GLU
1	A	512	ARG
1	A	517	ARG
1	A	553	THR
1	A	559	GLU
1	A	590	CYS
1	A	731	LEU
1	A	764	SER
1	A	779	LYS
1	A	841	LYS
1	A	898	ASN
1	A	915	PHE
1	A	919	PHE
1	A	930	VAL
1	A	937	ASP
1	A	1004	LEU
1	A	1009	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	GLN
1	A	247	HIS
1	A	278	HIS
1	A	291	ASN
1	A	610	GLN
1	A	617	GLN
1	A	748	GLN
1	A	836	ASN
1	A	838	GLN
1	A	898	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FGE	A	1101	-	20,26,26	2.17	6 (30%)	24,40,40	2.39	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FGE	A	1101	-	-	4/12/23/23	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	FGE	S24-N17	5.36	1.71	1.60
2	A	1101	FGE	C5-C12	4.92	1.48	1.42
2	A	1101	FGE	C12-N16	3.67	1.44	1.33
2	A	1101	FGE	C3-C7	-2.48	1.45	1.48
2	A	1101	FGE	C4-C6	-2.42	1.38	1.41
2	A	1101	FGE	C13-C8	-2.14	1.46	1.50

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	FGE	C5-C12-N16	-5.78	100.71	112.15
2	A	1101	FGE	C6-S24-N17	-4.82	99.39	108.28
2	A	1101	FGE	C1-C5-C12	-4.54	128.09	135.50
2	A	1101	FGE	O20-C11-N18	3.46	127.60	123.04
2	A	1101	FGE	C12-N16-C10	3.25	113.98	107.59
2	A	1101	FGE	C4-C10-N16	-2.54	105.47	109.91
2	A	1101	FGE	C14-C11-N18	-2.50	111.32	114.98
2	A	1101	FGE	O19-C10-C4	2.30	125.90	122.65
2	A	1101	FGE	O19-C10-N16	2.09	128.62	124.41

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1101	FGE	C4-C6-S24-N17
2	A	1101	FGE	C1-C3-C7-C8
2	A	1101	FGE	C2-C3-C7-C8
2	A	1101	FGE	C2-C6-S24-N17

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	FGE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	822/938 (87%)	0.42	42 (5%)	28 33	36, 61, 92, 123	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	847	THR	5.9
1	A	225	ALA	5.7
1	A	396	ALA	5.5
1	A	913	GLY	5.2
1	A	226	THR	5.0
1	A	515	LEU	4.0
1	A	334	ASN	3.9
1	A	367	SER	3.9
1	A	203	PHE	3.7
1	A	395	TYR	3.6
1	A	1017	VAL	3.5
1	A	512	ARG	3.4
1	A	517	ARG	3.4
1	A	205	PHE	3.3
1	A	370	VAL	3.3
1	A	471	LEU	3.2
1	A	416	CYS	3.2
1	A	842	SER	2.8
1	A	898	ASN	2.8
1	A	341	LEU	2.8
1	A	228	PHE	2.7
1	A	371	TRP	2.7
1	A	196	PHE	2.6
1	A	418	ILE	2.6
1	A	484	TYR	2.5
1	A	342	VAL	2.5
1	A	1024	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	462	ASN	2.4
1	A	366	CYS	2.4
1	A	472	VAL	2.4
1	A	317	TRP	2.4
1	A	514	ILE	2.3
1	A	417	PRO	2.2
1	A	846	ALA	2.2
1	A	256	LEU	2.2
1	A	1023	LEU	2.1
1	A	136	VAL	2.1
1	A	214	LEU	2.1
1	A	526	HIS	2.1
1	A	486	PRO	2.0
1	A	897	ASP	2.0
1	A	934	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FGE	A	1101	24/24	0.95	0.16	54,57,58,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.