



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2024 – 11:39 PM EDT

PDB ID : 3HEI
Title : Ligand Recognition by A-Class Eph Receptors: Crystal Structures of the EphA2 Ligand-Binding Domain and the EphA2/ephrin-A1 Complex
Authors : Himanen, J.P.; Goldgur, Y.; Miao, H.; Myshkin, E.; Guo, H.; Buck, M.; Nguyen, M.; Rajashankar, K.R.; Wang, B.; Nikolov, D.B.
Deposited on : 2009-05-08
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	174	<div> <div>7%</div> <div>79%</div> <div>16%</div> <div>5%</div> </div>
1	C	174	<div> <div>2%</div> <div>78%</div> <div>18%</div> <div>.</div> </div>
1	E	174	<div> <div>4%</div> <div>79%</div> <div>18%</div> <div>..</div> </div>
1	G	174	<div> <div>77%</div> <div>17%</div> <div>5%</div> <div>.</div> </div>
1	I	174	<div> <div>20%</div> <div>77%</div> <div>21%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	174	
1	M	174	
1	O	174	
2	B	132	
2	D	132	
2	F	132	
2	H	132	
2	J	132	
2	L	132	
2	N	132	
2	P	132	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 22908 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ephrin type-A receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			
1	C	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			
1	E	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			
1	G	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			
1	I	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			
1	K	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			
1	M	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			
1	O	174	Total	C	N	O	S	0	0	0
			1399	896	231	262	10			

- Molecule 2 is a protein called Ephrin-A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			
2	D	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			
2	F	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			
2	H	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			
2	J	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			
2	L	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			
2	P	132	Total	C	N	O	S	0	0	0
			1120	712	197	206	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	148	LYS	-	expression tag	UNP P20827
B	149	ILE	-	expression tag	UNP P20827
D	148	LYS	-	expression tag	UNP P20827
D	149	ILE	-	expression tag	UNP P20827
F	148	LYS	-	expression tag	UNP P20827
F	149	ILE	-	expression tag	UNP P20827
H	148	LYS	-	expression tag	UNP P20827
H	149	ILE	-	expression tag	UNP P20827
J	148	LYS	-	expression tag	UNP P20827
J	149	ILE	-	expression tag	UNP P20827
L	148	LYS	-	expression tag	UNP P20827
L	149	ILE	-	expression tag	UNP P20827
N	148	LYS	-	expression tag	UNP P20827
N	149	ILE	-	expression tag	UNP P20827
P	148	LYS	-	expression tag	UNP P20827
P	149	ILE	-	expression tag	UNP P20827

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	200	Total	O	0	0
			200	200		
3	B	226	Total	O	0	0
			226	226		
3	C	225	Total	O	0	0
			225	225		
3	D	171	Total	O	0	0
			171	171		
3	E	178	Total	O	0	0
			178	178		
3	F	179	Total	O	0	0
			179	179		
3	G	219	Total	O	0	0
			219	219		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	203	Total 203	O 203	0	0
3	I	115	Total 115	O 115	0	0
3	J	145	Total 145	O 145	0	0
3	K	173	Total 173	O 173	0	0
3	L	149	Total 149	O 149	0	0
3	M	157	Total 157	O 157	0	0
3	N	173	Total 173	O 173	0	0
3	O	99	Total 99	O 99	0	0
3	P	144	Total 144	O 144	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ephrin type-A receptor 2




- Molecule 1: Ephrin type-A receptor 2




- Molecule 1: Ephrin type-A receptor 2






- Molecule 1: Ephrin type-A receptor 2









• Molecule 1: Ephrin type-A receptor 2






• Molecule 1: Ephrin type-A receptor 2



• Molecule 1: Ephrin type-A receptor 2




• Molecule 1: Ephrin type-A receptor 2




• Molecule 2: Ephrin-A1







• Molecule 2: Ephrin-A1



• Molecule 2: Ephrin-A1




• Molecule 2: Ephrin-A1




• Molecule 2: Ephrin-A1




- Molecule 2: Ephrin-A1



- Molecule 2: Ephrin-A1



- Molecule 2: Ephrin-A1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	58.48Å 102.11Å 135.25Å 84.25° 79.77° 73.94°	Depositor
Resolution (Å)	50.00 – 2.00 43.60 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.0 (50.00-2.00) 96.0 (43.60-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.40 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.172 , 0.233 0.209 , 0.265	Depositor DCC
R_{free} test set	9696 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for h,h-k,h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	22908	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.32	4/1433 (0.3%)	1.15	8/1940 (0.4%)
1	C	1.27	2/1433 (0.1%)	1.26	14/1940 (0.7%)
1	E	1.17	2/1433 (0.1%)	1.03	6/1940 (0.3%)
1	G	1.29	9/1433 (0.6%)	1.06	5/1940 (0.3%)
1	I	0.92	1/1433 (0.1%)	0.90	2/1940 (0.1%)
1	K	1.19	2/1433 (0.1%)	1.04	5/1940 (0.3%)
1	M	1.00	1/1433 (0.1%)	0.93	0/1940
1	O	0.90	1/1433 (0.1%)	0.91	0/1940
2	B	1.36	7/1154 (0.6%)	1.09	3/1562 (0.2%)
2	D	1.34	9/1154 (0.8%)	1.07	8/1562 (0.5%)
2	F	1.29	5/1154 (0.4%)	1.11	3/1562 (0.2%)
2	H	1.45	12/1154 (1.0%)	1.15	8/1562 (0.5%)
2	J	1.14	4/1154 (0.3%)	0.99	2/1562 (0.1%)
2	L	1.23	2/1154 (0.2%)	0.97	3/1562 (0.2%)
2	N	1.24	5/1154 (0.4%)	1.00	3/1562 (0.2%)
2	P	1.15	4/1154 (0.3%)	0.97	1/1562 (0.1%)
All	All	1.21	70/20696 (0.3%)	1.04	71/28016 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	G	0	1
2	B	0	2
2	F	0	1
2	J	0	1
All	All	0	6

The worst 5 of 70 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	26	ASN	CG-ND2	9.52	1.56	1.32
2	N	37	TYR	CD2-CE2	9.47	1.53	1.39
2	H	26	ASN	CA-CB	9.43	1.77	1.53
2	B	26	ASN	CB-CG	9.11	1.72	1.51
2	D	26	ASN	CG-ND2	8.73	1.54	1.32

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	66	ARG	NE-CZ-NH2	-16.20	112.20	120.30
1	E	66	ARG	NE-CZ-NH1	11.75	126.18	120.30
1	C	66	ARG	NE-CZ-NH1	10.44	125.52	120.30
1	A	102	ASP	CB-CG-OD1	10.05	127.34	118.30
1	A	102	ASP	CB-CG-OD2	-9.33	109.91	118.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	48	GLY	Peptide
2	B	135	GLN	Peptide
2	B	148	LYS	Peptide
2	F	148	LYS	Peptide
1	G	83	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1399	0	1348	23	0
1	C	1399	0	1348	30	0
1	E	1399	0	1348	33	0
1	G	1399	0	1348	30	0
1	I	1399	0	1348	31	0
1	K	1399	0	1348	27	0
1	M	1399	0	1348	43	0
1	O	1399	0	1348	27	0
2	B	1120	0	1058	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1120	0	1058	39	0
2	F	1120	0	1058	61	0
2	H	1120	0	1058	53	0
2	J	1120	0	1058	45	0
2	L	1120	0	1058	34	0
2	N	1120	0	1058	29	0
2	P	1120	0	1058	41	0
3	A	200	0	0	8	0
3	B	226	0	0	11	1
3	C	225	0	0	13	0
3	D	171	0	0	8	0
3	E	178	0	0	10	0
3	F	179	0	0	19	0
3	G	219	0	0	11	0
3	H	203	0	0	19	1
3	I	115	0	0	7	0
3	J	145	0	0	11	0
3	K	173	0	0	8	1
3	L	149	0	0	7	0
3	M	157	0	0	13	1
3	N	173	0	0	8	0
3	O	99	0	0	10	0
3	P	144	0	0	6	0
All	All	22908	0	19248	542	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 542 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:26:ASN:CA	2:H:26:ASN:CB	1.77	1.57
2:F:47:VAL:HB	3:F:1187:HOH:O	1.27	1.30
2:H:18:ALA:HB3	3:H:720:HOH:O	1.18	1.27
2:H:139:ARG:HD3	3:H:1411:HOH:O	1.29	1.23
2:P:18:ALA:O	3:P:1057:HOH:O	1.58	1.19

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:K:438:HOH:O	3:M:1257:HOH:O[1_565]	1.97	0.23
3:B:2558:HOH:O	3:H:661:HOH:O[1_544]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	172/174 (99%)	166 (96%)	5 (3%)	1 (1%)	25	19
1	C	172/174 (99%)	165 (96%)	6 (4%)	1 (1%)	25	19
1	E	172/174 (99%)	163 (95%)	7 (4%)	2 (1%)	13	7
1	G	172/174 (99%)	164 (95%)	5 (3%)	3 (2%)	9	4
1	I	172/174 (99%)	165 (96%)	5 (3%)	2 (1%)	13	7
1	K	172/174 (99%)	164 (95%)	5 (3%)	3 (2%)	9	4
1	M	172/174 (99%)	161 (94%)	11 (6%)	0	100	100
1	O	172/174 (99%)	159 (92%)	12 (7%)	1 (1%)	25	19
2	B	130/132 (98%)	125 (96%)	3 (2%)	2 (2%)	10	4
2	D	130/132 (98%)	125 (96%)	1 (1%)	4 (3%)	4	1
2	F	130/132 (98%)	125 (96%)	2 (2%)	3 (2%)	6	2
2	H	130/132 (98%)	125 (96%)	1 (1%)	4 (3%)	4	1
2	J	130/132 (98%)	124 (95%)	3 (2%)	3 (2%)	6	2
2	L	130/132 (98%)	126 (97%)	2 (2%)	2 (2%)	10	4
2	N	130/132 (98%)	126 (97%)	2 (2%)	2 (2%)	10	4
2	P	130/132 (98%)	122 (94%)	4 (3%)	4 (3%)	4	1
All	All	2416/2448 (99%)	2305 (95%)	74 (3%)	37 (2%)	10	4

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	135	GLN
2	D	58	SER
2	F	58	SER
2	F	139	ARG
2	H	56	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/150 (100%)	139 (93%)	11 (7%)	14	9
1	C	150/150 (100%)	142 (95%)	8 (5%)	22	18
1	E	150/150 (100%)	141 (94%)	9 (6%)	19	14
1	G	150/150 (100%)	145 (97%)	5 (3%)	38	37
1	I	150/150 (100%)	142 (95%)	8 (5%)	22	18
1	K	150/150 (100%)	140 (93%)	10 (7%)	16	11
1	M	150/150 (100%)	143 (95%)	7 (5%)	26	22
1	O	150/150 (100%)	142 (95%)	8 (5%)	22	18
2	B	124/124 (100%)	117 (94%)	7 (6%)	21	17
2	D	124/124 (100%)	118 (95%)	6 (5%)	25	22
2	F	124/124 (100%)	116 (94%)	8 (6%)	17	12
2	H	124/124 (100%)	115 (93%)	9 (7%)	14	9
2	J	124/124 (100%)	121 (98%)	3 (2%)	49	51
2	L	124/124 (100%)	116 (94%)	8 (6%)	17	12
2	N	124/124 (100%)	117 (94%)	7 (6%)	21	17
2	P	124/124 (100%)	116 (94%)	8 (6%)	17	12
All	All	2192/2192 (100%)	2070 (94%)	122 (6%)	21	17

5 of 122 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	58	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	O	130	GLU
2	J	56	ASP
1	O	101	LEU
2	P	84	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 86 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	26	ASN
2	N	78	GLN
2	L	66	GLN
1	M	106	ASN
1	O	19	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	174/174 (100%)	0.56	13 (7%) 14 13	2, 5, 24, 42	0
1	C	174/174 (100%)	0.10	4 (2%) 60 59	2, 5, 20, 28	0
1	E	174/174 (100%)	0.21	7 (4%) 38 37	2, 8, 22, 36	0
1	G	174/174 (100%)	-0.08	0 100 100	2, 6, 20, 30	0
1	I	174/174 (100%)	0.92	35 (20%) 1 0	4, 11, 25, 35	0
1	K	174/174 (100%)	0.20	7 (4%) 38 37	2, 7, 23, 32	0
1	M	174/174 (100%)	0.43	16 (9%) 9 8	3, 8, 22, 27	0
1	O	174/174 (100%)	1.04	40 (22%) 0 0	4, 11, 24, 35	0
2	B	132/132 (100%)	0.22	7 (5%) 26 25	2, 6, 30, 41	0
2	D	132/132 (100%)	0.19	8 (6%) 21 20	2, 5, 24, 38	0
2	F	132/132 (100%)	0.04	5 (3%) 40 39	2, 6, 24, 37	0
2	H	132/132 (100%)	0.27	4 (3%) 50 49	2, 5, 22, 29	0
2	J	132/132 (100%)	0.13	10 (7%) 13 13	2, 6, 32, 39	0
2	L	132/132 (100%)	0.00	7 (5%) 26 25	2, 6, 21, 41	0
2	N	132/132 (100%)	0.01	5 (3%) 40 39	2, 6, 24, 39	0
2	P	132/132 (100%)	0.37	13 (9%) 7 7	2, 6, 29, 38	0
All	All	2448/2448 (100%)	0.31	181 (7%) 14 13	2, 7, 24, 42	0

The worst 5 of 181 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	18	ALA	11.1
1	O	174	CYS	10.7
1	I	174	CYS	10.1
1	K	82	PRO	9.7
2	N	18	ALA	9.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.