



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2024 – 06:15 AM EST

PDB ID : 3HHS  
Title : Crystal Structure of Manduca sexta prophenoloxidase  
Authors : Li, Y.; Wang, Y.; Jiang, H.; Deng, J.  
Deposited on : 2009-05-17  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1356 (1.98-1.98)
Clashscore	180529	1437 (1.98-1.98)
Ramachandran outliers	177936	1426 (1.98-1.98)
Sidechain outliers	177891	1426 (1.98-1.98)
RSRZ outliers	164620	1356 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	694	
2	B	684	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenoloxidase subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	669	Total	C	N	O	S	41	3	0
			5456	3457	950	1019	30			

- Molecule 2 is a protein called Phenoloxidase subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	656	Total	C	N	O	S	36	1	0
			5349	3390	945	978	36			

- Molecule 3 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cu	0	0
			2	2		
3	B	2	Total	Cu	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	682	Total	O	0	0
			682	682		
4	B	513	Total	O	0	0
			513	513		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Phenoloxidase subunit 2



#### • Molecule 2: Phenoloxidase subunit 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	148.83Å 153.66Å 75.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.45 – 1.97 42.45 – 1.97	Depositor EDS
% Data completeness (in resolution range)	98.9 (42.45-1.97) 98.9 (42.45-1.97)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.154 , 0.194 0.165 , 0.199	Depositor DCC
$R_{free}$ test set	6106 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.7	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12004	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.92	3/5600 (0.1%)	0.78	6/7593 (0.1%)
2	B	0.92	3/5492 (0.1%)	0.72	3/7437 (0.0%)
All	All	0.92	6/11092 (0.1%)	0.75	9/15030 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	1
All	All	0	3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	78	TYR	CD1-CE1	-6.11	1.30	1.39
1	A	197	GLU	CD-OE2	-5.64	1.19	1.25
2	B	28	VAL	CB-CG1	-5.38	1.41	1.52
2	B	248	VAL	C-N	-5.33	1.21	1.34
1	A	373	SER	CB-OG	-5.15	1.35	1.42
1	A	562	GLU	CD-OE1	-5.10	1.20	1.25

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	161	LEU	CB-CA-C	9.17	127.62	110.20
1	A	161	LEU	C-N-CA	7.77	141.12	121.70
1	A	475	ARG	NE-CZ-NH2	-7.08	116.76	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	200	ARG	NE-CZ-NH2	-5.99	117.30	120.30
2	B	200	ARG	NE-CZ-NH1	5.80	123.20	120.30
2	B	468	ARG	NE-CZ-NH2	-5.70	117.45	120.30
1	A	321	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	202	LEU	CA-CB-CG	-5.32	103.06	115.30
1	A	161	LEU	N-CA-C	-5.20	96.96	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	161	LEU	Peptide
1	A	629	SER	Peptide
2	B	536	ARG	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5456	0	5303	54	0
2	B	5349	0	5183	56	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	682	0	0	10	0
4	B	513	0	0	5	0
All	All	12004	0	10486	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:LEU:HB2	1:A:206[B]:ARG:NH1	1.57	1.19
2:B:474:ARG:HG2	2:B:474:ARG:NH1	1.58	1.04

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:474:ARG:HH11	2:B:474:ARG:CG	1.72	1.03
1:A:202:LEU:CB	1:A:206[B]:ARG:HH11	1.77	0.97
1:A:311:ASN:HB2	4:A:771:HOH:O	1.65	0.96
2:B:474:ARG:HG2	2:B:474:ARG:HH11	0.79	0.95
1:A:202:LEU:HB2	1:A:206[B]:ARG:HH11	1.16	0.91
1:A:35:GLN:HE22	1:A:67:LEU:H	1.22	0.86
2:B:479:GLN:HE21	2:B:480:HIS:H	1.26	0.83
2:B:414:ASN:HD21	2:B:522:ARG:HH22	1.27	0.81
1:A:362:ASP:HB2	4:A:1167:HOH:O	1.83	0.77
1:A:161:LEU:O	1:A:282:LEU:CD2	2.33	0.77
1:A:553:THR:HA	1:A:592:GLN:HE22	1.50	0.76
1:A:214:HIS:HD2	1:A:589:GLY:H	1.34	0.74
1:A:202:LEU:CB	1:A:206[B]:ARG:NH1	2.39	0.73
1:A:206[B]:ARG:HH21	1:A:402:PRO:HB3	1.55	0.71
2:B:441:GLU:OE2	2:B:543:ARG:NH1	2.23	0.71
2:B:616:GLN:HE22	2:B:635:LYS:H	1.38	0.71
1:A:446:GLU:OE2	1:A:452:GLN:HG3	1.91	0.69
1:A:202:LEU:HB2	1:A:206[B]:ARG:HH12	1.57	0.68
2:B:534:ALA:O	2:B:535:LEU:HD23	1.94	0.68
1:A:247:GLN:HE22	1:A:533:GLU:H	1.41	0.68
2:B:238:MET:HG2	4:B:1022:HOH:O	1.92	0.67
1:A:199:GLU:OE1	1:A:206[B]:ARG:NH2	2.25	0.67
2:B:477:HIS:HD2	2:B:478:LEU:O	1.78	0.66
2:B:455:SER:OG	2:B:477:HIS:HE1	1.80	0.65
2:B:536:ARG:HB3	2:B:537:PRO:HD2	1.80	0.64
2:B:81:GLN:HE21	2:B:386:GLN:HE22	1.46	0.63
2:B:189:THR:HB	4:B:1042:HOH:O	1.99	0.62
4:A:1280:HOH:O	2:B:190:THR:HG21	1.98	0.62
1:A:202:LEU:HD12	1:A:202:LEU:N	2.14	0.62
1:A:56:GLU:HB2	4:A:1057:HOH:O	2.01	0.61
2:B:536:ARG:H	2:B:540:ASN:HD21	1.49	0.61
2:B:188:ASN:HB3	2:B:190:THR:HG22	1.82	0.60
1:A:500:ALA:HB3	4:A:1007:HOH:O	2.01	0.59
1:A:247:GLN:NE2	1:A:533:GLU:H	2.00	0.58
2:B:337:LEU:HD23	2:B:345:MET:HE3	1.85	0.58
2:B:188:ASN:OD1	2:B:189:THR:HG22	2.04	0.58
1:A:444:ALA:HA	1:A:453:ASN:HD21	1.70	0.56
1:A:135:THR:HG21	1:A:220:LEU:HD21	1.87	0.56
2:B:374:ALA:HB1	2:B:387:PHE:HB2	1.88	0.56
1:A:202:LEU:N	1:A:202:LEU:CD1	2.67	0.56
2:B:207:LEU:HD13	2:B:558:THR:HG21	1.87	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:441:GLU:HG3	2:B:486:VAL:CG2	2.36	0.55
2:B:474:ARG:NH1	2:B:474:ARG:CG	2.40	0.54
2:B:414:ASN:ND2	2:B:522:ARG:HH22	2.03	0.54
1:A:3:ASP:HA	4:A:1239:HOH:O	2.07	0.53
1:A:487:HIS:HE1	4:A:723:HOH:O	1.92	0.53
2:B:480:HIS:HE1	2:B:586:HIS:O	1.92	0.53
1:A:177:PRO:HG2	1:A:180:ILE:HD12	1.91	0.53
2:B:534:ALA:C	2:B:535:LEU:HD23	2.29	0.52
2:B:351:GLU:N	2:B:352:SER:HA	2.24	0.52
1:A:140:HIS:HE1	1:A:234:LYS:NZ	2.07	0.52
2:B:54:ASN:O	2:B:55:GLU:HB2	2.10	0.52
1:A:136:VAL:O	1:A:140:HIS:HD2	1.95	0.50
1:A:74:PRO:HA	1:A:77:LYS:HE3	1.94	0.50
1:A:206[B]:ARG:HH21	1:A:402:PRO:CB	2.24	0.50
1:A:161:LEU:O	1:A:282:LEU:HD22	2.08	0.50
1:A:205:TRP:CD1	1:A:255:GLU:HG3	2.47	0.50
1:A:379:HIS:H	1:A:379:HIS:CD2	2.30	0.50
1:A:464:VAL:HG13	1:A:482:PHE:HE2	1.77	0.49
1:A:11:LEU:HD21	1:A:119:VAL:HG22	1.93	0.49
1:A:557:LEU:HD13	1:A:635:MET:HE2	1.95	0.49
2:B:489:VAL:HG13	2:B:540:ASN:HB2	1.94	0.49
1:A:445:VAL:H	1:A:453:ASN:ND2	2.10	0.49
2:B:491:ASN:HB3	2:B:536:ARG:O	2.13	0.48
2:B:131:VAL:O	2:B:135:HIS:HD2	1.96	0.48
2:B:246:TYR:CE2	2:B:250:ARG:HD2	2.49	0.48
1:A:639:LEU:HD13	1:A:642:LYS:HE3	1.95	0.48
1:A:164:GLN:HA	2:B:282:SER:HB3	1.94	0.48
1:A:201:ARG:C	1:A:202:LEU:HD12	2.34	0.48
1:A:202:LEU:HB3	1:A:206[B]:ARG:HH11	1.73	0.48
2:B:37:TYR:O	2:B:42:LYS:NZ	2.47	0.47
2:B:557:ARG:NH2	2:B:624:ASP:OD2	2.41	0.47
2:B:40:LYS:HG2	4:B:1066:HOH:O	2.14	0.47
1:A:135:THR:CG2	1:A:220:LEU:HD21	2.45	0.47
1:A:362:ASP:CB	4:A:1167:HOH:O	2.53	0.47
1:A:447:SER:HB2	1:A:450:GLY:HA3	1.97	0.47
2:B:484:GLN:HE21	2:B:545:ARG:HG2	1.80	0.47
1:A:4:ILE:O	1:A:4:ILE:HG13	2.14	0.46
2:B:215:VAL:HG13	4:B:1097:HOH:O	2.15	0.46
2:B:189:THR:HG23	2:B:190:THR:N	2.31	0.45
2:B:369:GLY:HA3	2:B:405:TRP:CZ2	2.52	0.45
1:A:217:HIS:HE1	4:A:1020:HOH:O	1.99	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:616:GLN:HE22	2:B:635:LYS:N	2.10	0.45
2:B:231:ARG:HD3	2:B:348:ASN:HD21	1.82	0.45
2:B:491:ASN:O	2:B:538:GLY:HA2	2.17	0.45
1:A:259:ASN:HD21	1:A:598:LYS:HE2	1.82	0.44
2:B:86:VAL:HG13	2:B:302:ASP:HB3	1.99	0.44
1:A:247:GLN:HE22	1:A:532:ILE:HA	1.83	0.44
1:A:551:GLN:HB2	1:A:554:GLU:HG3	1.99	0.44
2:B:327:ILE:HG21	2:B:349:LEU:HD11	2.00	0.44
1:A:140:HIS:HE1	1:A:234:LYS:HZ1	1.64	0.44
2:B:34:ASP:OD2	2:B:35:HIS:N	2.51	0.43
1:A:161:LEU:O	1:A:282:LEU:HD23	2.15	0.43
2:B:536:ARG:HB3	2:B:537:PRO:CD	2.48	0.43
2:B:148:ALA:HB1	2:B:203:ILE:HD11	2.00	0.42
1:A:400:ARG:HD3	4:A:797:HOH:O	2.18	0.42
1:A:134:TYR:CZ	1:A:138:ILE:HD11	2.55	0.41
2:B:123:TYR:CE1	2:B:154:LYS:HE3	2.56	0.41
2:B:295:ARG:NH1	4:B:824:HOH:O	2.50	0.41
2:B:414:ASN:HD21	2:B:522:ARG:NH2	2.04	0.41
2:B:468:ARG:HD3	2:B:468:ARG:HA	1.89	0.41
1:A:14:ARG:HG3	1:A:475:ARG:NH2	2.35	0.41
2:B:185:TYR:OH	2:B:398:ARG:NH2	2.47	0.41
2:B:480:HIS:HD2	2:B:481:ASP:O	2.04	0.41
1:A:219:HIS:CD2	1:A:395[A]:GLU:OE1	2.75	0.40
2:B:121:ASN:HA	2:B:122:PRO:HD3	1.98	0.40
2:B:463:LEU:HG	2:B:466:THR:HG21	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	665/694 (96%)	650 (98%)	15 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	649/684 (95%)	634 (98%)	15 (2%)	0	100	100
All	All	1314/1378 (95%)	1284 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	601/618 (97%)	596 (99%)	5 (1%)	79	79
2	B	590/608 (97%)	584 (99%)	6 (1%)	73	72
All	All	1191/1226 (97%)	1180 (99%)	11 (1%)	75	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	161	LEU
1	A	230	LYS
1	A	237	ARG
1	A	562	GLU
2	B	183	ILE
2	B	231	ARG
2	B	444	ARG
2	B	474	ARG
2	B	522	ARG
2	B	596	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	140	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	150	GLN
1	A	214	HIS
1	A	217	HIS
1	A	246	GLN
1	A	247	GLN
1	A	259	ASN
1	A	379	HIS
1	A	453	ASN
1	A	487	HIS
1	A	498	ASN
1	A	592	GLN
2	B	22	GLN
2	B	50	ASN
2	B	64	ASN
2	B	135	HIS
2	B	206	ASN
2	B	253	ASN
2	B	348	ASN
2	B	386	GLN
2	B	414	ASN
2	B	477	HIS
2	B	479	GLN
2	B	480	HIS
2	B	484	GLN
2	B	496	ASN
2	B	540	ASN
2	B	596	GLN
2	B	616	GLN
2	B	673	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	669/694 (96%)	-0.41	14 (2%) 63 72	6, 13, 21, 36	21 (3%)
2	B	656/684 (95%)	-0.27	8 (1%) 76 83	8, 15, 22, 35	13 (1%)
All	All	1325/1378 (96%)	-0.34	22 (1%) 69 77	6, 14, 22, 36	34 (2%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	450	GLY	4.6
1	A	586	CYS	4.2
2	B	625	ALA	4.2
2	B	614	ILE	4.2
2	B	331	ASN	4.1
2	B	623	ASN	4.1
2	B	615	GLU	3.8
1	A	585	TYR	3.3
2	B	582	CYS	3.2
1	A	451	GLN	3.1
2	B	583	GLY	3.0
1	A	161	LEU	2.7
1	A	605	ALA	2.6
1	A	500	ALA	2.6
1	A	630	CYS	2.5
1	A	623	PRO	2.5
2	B	617	ASP	2.4
1	A	629	SER	2.4
1	A	602	GLY	2.3
1	A	603	GLY	2.3
1	A	452	GLN	2.2
1	A	604	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CU	A	697	1/1	0.99	0.04	33,33,33,33	0
3	CU	B	1	1/1	0.99	0.02	23,23,23,23	0
3	CU	B	686	1/1	0.99	0.02	25,25,25,25	0
3	CU	A	696	1/1	1.00	0.03	24,24,24,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.