



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 08:29 AM EDT

PDB ID : 2I3A
Title : Crystal structure of N-Acetyl-gamma-Glutamyl-Phosphate Reductase (Rv1652) from Mycobacterium tuberculosis
Authors : Cherney, L.T.; Cherney, M.M.; Garen, C.R.; Moraidin, F.; James, M.N.G.; Mycobacterium Tuberculosis Structural Proteomics Project (XMTB)
Deposited on : 2006-08-17
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	<div> <div>83%</div> <div>13%</div> <div>..</div> </div>
1	B	352	<div> <div>88%</div> <div>9%</div> <div>..</div> </div>
1	C	352	<div> <div>86%</div> <div>10%</div> <div>..</div> </div>
1	D	352	<div> <div>87%</div> <div>10%</div> <div>..</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-acetyl-gamma-glutamyl-phosphate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	0	0
			2496	1576	448	467	5			
1	B	346	Total	C	N	O	S	0	0	0
			2509	1583	451	470	5			
1	C	344	Total	C	N	O	S	0	0	0
			2496	1576	448	467	5			
1	D	347	Total	C	N	O	S	0	0	0
			2516	1588	452	471	5			

- Molecule 2 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: $C_8H_{19}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	203	Total	O	0	0
			203	203		
3	B	230	Total	O	0	0
			230	230		
3	C	227	Total	O	0	0
			227	227		
3	D	256	Total	O	0	0
			256	256		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase




- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase



- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase



- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.69Å 104.03Å 138.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.68 – 2.15 48.69 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.68-2.15) 98.6 (48.69-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.161 , 0.225 0.161 , 0.164	Depositor DCC
R_{free} test set	3541 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10989	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BTB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	0/2552	0.85	1/3494 (0.0%)
1	B	0.92	0/2565	0.85	2/3512 (0.1%)
1	C	0.93	0/2552	0.88	4/3494 (0.1%)
1	D	0.95	1/2572 (0.0%)	0.87	2/3522 (0.1%)
All	All	0.92	1/10241 (0.0%)	0.86	9/14022 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	158	CYS	CB-SG	5.37	1.91	1.82

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	ASP	CB-CG-OD1	6.37	124.03	118.30
1	B	248	ARG	NE-CZ-NH2	-5.97	117.31	120.30
1	C	228	ARG	NE-CZ-NH2	-5.63	117.48	120.30
1	D	41	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	C	243	LEU	CB-CG-CD1	-5.60	101.48	111.00
1	C	306	ASP	CB-CG-OD1	5.33	123.10	118.30
1	C	193	ARG	NE-CZ-NH2	-5.15	117.72	120.30
1	D	41	ARG	NE-CZ-NH2	-5.06	117.77	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	282	MET	CG-SD-CE	5.02	108.23	100.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	49	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2496	0	2497	42	0
1	B	2509	0	2508	23	0
1	C	2496	0	2497	20	0
1	D	2516	0	2517	26	0
2	A	14	0	19	1	0
2	B	14	0	19	1	0
2	C	14	0	19	2	0
2	D	14	0	19	0	0
3	A	203	0	0	9	0
3	B	230	0	0	3	0
3	C	227	0	0	2	0
3	D	256	0	0	5	0
All	All	10989	0	10095	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:310:GLN:HG3	3:A:1584:HOH:O	1.30	1.28
1:B:264:GLN:HG3	3:B:1710:HOH:O	1.37	1.22
1:C:101:SER:HB3	1:C:103:GLU:OE2	1.56	1.05

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:257:ARG:HD2	3:A:1541:HOH:O	1.55	1.04
1:B:39:ARG:HG3	2:B:1500:BTB:O1	1.84	0.76
1:C:39:ARG:HD3	2:C:1600:BTB:O1	1.85	0.75
1:A:11:LYS:HD2	3:A:1590:HOH:O	1.87	0.74
1:C:266:ARG:NH1	1:C:303:VAL:O	2.21	0.73
1:D:174:ASP:HB2	3:D:1930:HOH:O	1.87	0.73
1:A:68:ARG:NH1	3:A:1592:HOH:O	2.23	0.70
1:D:41:ARG:N	1:D:41:ARG:HD3	2.07	0.70
1:D:270:GLU:O	1:D:274:HIS:HB2	1.92	0.69
1:A:147:LEU:O	1:A:150:THR:HG22	1.93	0.68
1:B:257:ARG:NH1	1:B:310:GLN:OE1	2.26	0.68
1:C:101:SER:CB	1:C:103:GLU:OE2	2.40	0.66
1:B:250:ILE:HB	1:B:318:ILE:HD11	1.75	0.66
1:B:68:ARG:HD2	3:B:1645:HOH:O	1.95	0.65
1:B:102:PRO:HA	1:B:152:ARG:HH22	1.61	0.65
1:A:91:GLY:HA2	3:A:1589:HOH:O	1.97	0.65
1:C:39:ARG:NH1	3:C:1773:HOH:O	2.29	0.63
1:A:135:TYR:CZ	1:A:156:PRO:HB3	2.33	0.63
1:A:315:ILE:HD13	1:D:210:ALA:HB3	1.81	0.63
1:B:91:GLY:HA2	3:B:1546:HOH:O	1.99	0.62
1:A:88:LEU:HB3	1:A:91:GLY:HA3	1.81	0.62
1:D:211:TYR:CE2	1:D:243:LEU:HG	2.35	0.62
1:A:102:PRO:HA	1:A:152:ARG:HH22	1.65	0.61
1:A:11:LYS:NZ	1:A:41:ARG:HH11	1.98	0.61
1:A:11:LYS:HZ1	1:A:41:ARG:HH11	1.46	0.60
1:D:193:ARG:O	1:D:193:ARG:HG2	2.02	0.60
1:B:104:THR:HB	1:B:106:ILE:HD11	1.84	0.60
1:B:250:ILE:HB	1:B:318:ILE:CD1	2.33	0.57
1:D:261:PRO:HD2	1:D:264:GLN:HE21	1.69	0.57
1:B:118:ALA:O	1:B:122:GLU:HG3	2.04	0.57
1:A:284:GLU:HB3	3:A:1573:HOH:O	2.04	0.57
1:B:37:ASP:OD1	1:B:39:ARG:HG2	2.07	0.55
1:A:266:ARG:O	1:A:270:GLU:HG2	2.06	0.55
1:B:197:THR:O	1:B:203:GLU:HG3	2.07	0.55
1:A:39:ARG:NH2	3:A:1534:HOH:O	2.28	0.55
1:A:89:PRO:HD2	3:A:1427:HOH:O	2.07	0.54
1:C:107:ILE:HG12	1:C:153:ILE:HD12	1.90	0.54
1:C:172:ALA:HB2	1:C:230:VAL:HB	1.90	0.54
1:D:41:ARG:N	1:D:41:ARG:CD	2.70	0.53
1:A:11:LYS:HE2	1:A:41:ARG:NH1	2.22	0.53
1:D:71:GLU:HB3	1:D:72:PRO:HD2	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:CYS:HB2	1:A:250:ILE:CD1	2.39	0.53
1:A:137:LEU:HG	1:A:140:LEU:HG	1.90	0.53
1:D:273:TYR:CG	1:D:279:ILE:HG21	2.44	0.53
1:D:68:ARG:NH1	3:D:1942:HOH:O	2.29	0.52
1:C:100:LEU:HD12	1:C:106:ILE:HD12	1.91	0.52
1:B:250:ILE:HG13	1:B:320:ASN:HB3	1.92	0.52
1:C:197:THR:O	1:C:203:GLU:HG3	2.09	0.51
1:B:185:VAL:HG22	1:B:251:LEU:HB3	1.93	0.51
1:A:284:GLU:H	1:A:284:GLU:CD	2.14	0.51
1:D:273:TYR:CD2	1:D:279:ILE:HG21	2.46	0.51
1:A:88:LEU:CB	1:A:91:GLY:HA3	2.40	0.51
1:B:107:ILE:HG12	1:B:153:ILE:HD12	1.93	0.51
1:C:284:GLU:OE2	1:C:284:GLU:HA	2.11	0.49
1:D:49:THR:O	1:D:50:SER:C	2.50	0.49
1:A:273:TYR:CG	1:A:279:ILE:HG21	2.47	0.49
1:C:90:HIS:HA	1:C:112:ASP:OD1	2.13	0.49
1:C:147:LEU:O	1:C:150:THR:HG22	2.13	0.49
1:A:273:TYR:CD2	1:A:279:ILE:HG21	2.49	0.48
1:A:315:ILE:HD11	1:D:213:ILE:HG21	1.95	0.48
1:A:51:ALA:CB	1:A:72:PRO:HG3	2.44	0.48
1:C:257:ARG:HA	1:C:257:ARG:HD2	1.66	0.48
1:D:244:ILE:HB	1:D:245:PRO:CD	2.44	0.48
1:A:168:PHE:C	1:A:168:PHE:CD1	2.88	0.47
2:A:1400:BTB:H71	2:A:1400:BTB:O3	2.14	0.47
1:D:90:HIS:HD2	3:D:1734:HOH:O	1.97	0.47
1:A:185:VAL:O	1:A:250:ILE:HA	2.15	0.47
1:C:74:GLU:HG2	1:C:77:VAL:HG23	1.97	0.47
1:D:9:ALA:HB1	1:D:41:ARG:HH22	1.80	0.47
1:A:315:ILE:HD12	1:D:213:ILE:HD13	1.97	0.46
1:D:9:ALA:HB1	1:D:41:ARG:NH2	2.30	0.46
1:B:109:CYS:HA	1:B:155:VAL:HB	1.97	0.45
1:D:286:GLN:NE2	3:D:1938:HOH:O	2.49	0.45
1:B:104:THR:HB	1:B:106:ILE:CD1	2.47	0.45
1:A:171:LEU:HD23	1:A:227:LEU:HB3	1.99	0.45
1:A:229:ALA:HB3	3:A:1567:HOH:O	2.15	0.45
1:B:200:LEU:HD21	1:D:195:ALA:HA	1.99	0.45
1:A:102:PRO:HA	1:A:152:ARG:NH2	2.29	0.45
1:B:270:GLU:O	1:B:274:HIS:HB3	2.17	0.44
1:A:200:LEU:HD21	1:C:195:ALA:HA	1.98	0.44
1:D:140:LEU:HD13	3:D:1850:HOH:O	2.17	0.44
1:A:37:ASP:C	1:A:37:ASP:OD1	2.56	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:GLY:O	1:A:92:HIS:ND1	2.50	0.44
1:D:109:CYS:HA	1:D:155:VAL:HB	2.00	0.44
1:B:39:ARG:HE	1:B:39:ARG:HB3	1.64	0.44
1:B:95:VAL:O	1:B:99:GLN:HG3	2.18	0.43
1:C:12:VAL:HG22	1:C:83:ALA:HB3	2.00	0.43
1:D:6:VAL:HG23	1:D:338:GLY:HA3	2.00	0.43
1:A:101:SER:C	1:A:103:GLU:H	2.21	0.43
1:A:87:ALA:C	1:A:89:PRO:HD3	2.39	0.43
1:D:63:THR:N	1:D:64:PRO:CD	2.82	0.43
1:C:233:ARG:CD	3:C:1667:HOH:O	2.67	0.42
1:A:159:TYR:HB2	1:A:160:PRO:HD3	2.01	0.42
1:A:176:ILE:HG22	1:A:258:THR:HB	2.00	0.42
1:C:39:ARG:HD3	2:C:1600:BTB:HO1	1.84	0.42
1:B:133:TRP:CD2	1:B:152:ARG:HB3	2.55	0.41
1:A:153:ILE:HD12	1:A:332:SER:HB3	2.03	0.41
1:A:276:GLU:HA	1:A:277:PRO:HD2	1.94	0.41
1:C:37:ASP:OD1	1:C:39:ARG:HG3	2.19	0.41
1:A:39:ARG:HE	1:A:39:ARG:HB3	1.74	0.41
1:A:87:ALA:HA	1:A:109:CYS:HB2	2.03	0.40
1:D:301:ILE:HA	1:D:315:ILE:O	2.21	0.40
1:C:160:PRO:O	1:C:164:LEU:HG	2.21	0.40
1:B:168:PHE:C	1:B:168:PHE:CD1	2.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/352 (97%)	332 (97%)	9 (3%)	1 (0%)	41	37
1	B	344/352 (98%)	335 (97%)	9 (3%)	0	100	100
1	C	342/352 (97%)	332 (97%)	9 (3%)	1 (0%)	41	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	345/352 (98%)	336 (97%)	8 (2%)	1 (0%)	41	37
All	All	1373/1408 (98%)	1335 (97%)	35 (2%)	3 (0%)	47	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	48	ALA
1	A	284	GLU
1	D	9	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/254 (97%)	239 (97%)	8 (3%)	39	38
1	B	248/254 (98%)	240 (97%)	8 (3%)	39	38
1	C	247/254 (97%)	237 (96%)	10 (4%)	31	29
1	D	249/254 (98%)	240 (96%)	9 (4%)	35	33
All	All	991/1016 (98%)	956 (96%)	35 (4%)	36	34

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
1	A	112	ASP
1	A	137	LEU
1	A	153	ILE
1	A	168	PHE
1	A	196	THR
1	A	245	PRO
1	A	270	GLU
1	B	39	ARG
1	B	95	VAL
1	B	128	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	137	LEU
1	B	168	PHE
1	B	257	ARG
1	B	281	LEU
1	B	318	ILE
1	C	98	GLN
1	C	103	GLU
1	C	122	GLU
1	C	128	SER
1	C	137	LEU
1	C	151	ARG
1	C	168	PHE
1	C	257	ARG
1	C	266	ARG
1	C	284	GLU
1	D	6	VAL
1	D	10	THR
1	D	41	ARG
1	D	96	LEU
1	D	112	ASP
1	D	168	PHE
1	D	196	THR
1	D	203	GLU
1	D	301	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	264	GLN
1	A	310	GLN
1	D	90	HIS
1	D	264	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BTB	C	1600	-	13,13,13	1.21	1 (7%)	7,16,16	0.92	0
2	BTB	A	1400	-	13,13,13	1.11	1 (7%)	7,16,16	1.05	0
2	BTB	B	1500	-	13,13,13	0.98	1 (7%)	7,16,16	1.20	1 (14%)
2	BTB	D	1700	-	13,13,13	1.43	2 (15%)	7,16,16	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTB	C	1600	-	-	1/21/21/21	-
2	BTB	A	1400	-	-	1/21/21/21	-
2	BTB	B	1500	-	-	4/21/21/21	-
2	BTB	D	1700	-	-	1/21/21/21	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1700	BTB	C2-N	2.78	1.54	1.48
2	C	1600	BTB	C2-N	2.68	1.53	1.48
2	A	1400	BTB	C2-N	2.67	1.53	1.48
2	D	1700	BTB	C4-C2	2.54	1.56	1.53
2	B	1500	BTB	C2-N	2.21	1.52	1.48

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1500	BTB	C8-C7-N	-2.25	102.81	111.59

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1500	BTB	C3-C2-N-C7
2	A	1400	BTB	N-C7-C8-O8
2	C	1600	BTB	N-C7-C8-O8
2	D	1700	BTB	N-C7-C8-O8
2	B	1500	BTB	C1-C2-N-C7
2	B	1500	BTB	C4-C2-N-C7
2	B	1500	BTB	N-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1600	BTB	2	0
2	A	1400	BTB	1	0
2	B	1500	BTB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/352 (97%)	-0.35	3 (0%) 84 88	17, 27, 46, 55	0
1	B	346/352 (98%)	-0.49	2 (0%) 89 91	17, 27, 46, 56	0
1	C	344/352 (97%)	-0.37	2 (0%) 89 91	16, 26, 45, 54	0
1	D	347/352 (98%)	-0.41	2 (0%) 89 91	16, 25, 45, 53	0
All	All	1381/1408 (98%)	-0.41	9 (0%) 87 91	16, 26, 46, 56	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	92	HIS	3.8
1	A	102	PRO	3.2
1	A	88	LEU	2.7
1	C	98	GLN	2.6
1	A	89	PRO	2.5
1	D	6	VAL	2.3
1	D	9	ALA	2.3
1	B	89	PRO	2.2
1	B	126	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BTB	A	1400	14/14	0.82	0.23	52,58,59,61	0
2	BTB	C	1600	14/14	0.83	0.27	44,52,55,57	0
2	BTB	B	1500	14/14	0.88	0.20	33,42,50,52	0
2	BTB	D	1700	14/14	0.92	0.14	26,33,39,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.