



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 05:39 AM EDT

PDB ID : 6I5R
Title : BlMnBP1 binding protein of an ABC transporter from *Bifidobacterium animalis* subsp. *lactis* ATCC27673 in complex with mannobiose
Authors : Abou Hachem, M.; Ejby, M.; Guskov, A.; Slotboom, D.J.
Deposited on : 2018-11-14
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	
2	B	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ZN	A	514	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7163 atoms, of which 3254 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sugar ABC transporter substrate-binding protein, BMnBP1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	428	Total	C	H	N	O	S	0	2	0
			6608	2129	3232	539	692	16			

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-beta-D-mannopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	H	O	0	0	0
			45	12	22	11			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	14	Total	Zn	0	0
			14	14		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	496	Total	O	0	0
			496	496		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sugar ABC transporter substrate-binding protein, BlMnBP1



- Molecule 2: beta-D-mannopyranose-(1-4)-beta-D-mannopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.56Å 74.50Å 102.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.83 – 1.60 48.83 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.83-1.60) 100.0 (48.83-1.60)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.217 , 0.246 0.219 , 0.246	Depositor DCC
R_{free} test set	2840 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7163	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	2/3459 (0.1%)	0.70	3/4690 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	254	PRO	C-N	-9.03	1.13	1.34
1	A	255	ASP	C-N	-5.00	1.22	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	ASP	CB-CG-OD1	8.24	125.72	118.30
1	A	254	PRO	C-N-CA	6.24	137.31	121.70
1	A	155	ASP	CB-CG-OD2	5.72	123.45	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3376	3232	3223	41	0
2	B	23	22	21	0	0
3	A	14	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	496	0	0	22	9
All	All	3909	3254	3244	42	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:GLU:OE1	4:A:603:HOH:O	2.00	0.79
1:A:255:ASP:OD2	4:A:604:HOH:O	2.04	0.76
1:A:424:GLY:O	4:A:605:HOH:O	2.09	0.69
1:A:191:GLU:OE2	4:A:606:HOH:O	2.10	0.68
1:A:416:ASP:OD2	4:A:607:HOH:O	2.14	0.66
1:A:244:LYS:HE2	1:A:459:GLU:CG	2.28	0.63
1:A:83:ASP:OD1	4:A:608:HOH:O	2.16	0.62
1:A:175:LYS:NZ	4:A:626:HOH:O	2.31	0.62
1:A:328:ASP:OD2	4:A:609:HOH:O	2.16	0.59
1:A:364:GLU:OE2	4:A:610:HOH:O	2.17	0.57
1:A:257:HIS:HE1	4:A:724:HOH:O	1.84	0.54
1:A:345:LYS:HD2	1:A:346:ASN:N	2.22	0.54
1:A:244:LYS:HE2	1:A:459:GLU:HG3	1.91	0.53
1:A:150:ASP:O	4:A:613:HOH:O	2.19	0.52
1:A:460:ASP:OD1	4:A:611:HOH:O	2.18	0.52
1:A:109:LEU:HD22	1:A:132:TYR:CD1	2.46	0.51
1:A:171:VAL:HG11	1:A:308:MET:HG3	1.91	0.50
1:A:276:GLU:OE1	4:A:615:HOH:O	2.20	0.50
1:A:459:GLU:OE1	4:A:614:HOH:O	2.19	0.50
1:A:201:GLU:OE1	4:A:612:HOH:O	2.19	0.48
1:A:378:LYS:HE2	1:A:380:ASP:OD2	2.13	0.48
1:A:86:LYS:O	4:A:616:HOH:O	2.20	0.46
1:A:168:ALA:HB1	1:A:224:TYR:CZ	2.50	0.46
1:A:405:GLY:HA2	4:A:962:HOH:O	2.16	0.46
1:A:419:LEU:HD12	1:A:454:TRP:HE3	1.81	0.46
1:A:345:LYS:HD2	1:A:346:ASN:OD1	2.16	0.46
1:A:421:ILE:HD13	1:A:428:LYS:HE3	1.99	0.45
1:A:462:GLU:O	4:A:617:HOH:O	2.21	0.45
1:A:345:LYS:CD	1:A:345:LYS:C	2.85	0.44
1:A:439:ARG:NH1	4:A:639:HOH:O	2.41	0.44
1:A:410:LEU:O	1:A:414:ASN:HB2	2.17	0.44
1:A:67:MET:O	1:A:76:ASN:HB2	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:GLU:OE2	4:A:620:HOH:O	2.21	0.43
1:A:375:ILE:HG12	1:A:384:MET:SD	2.60	0.42
1:A:215:ASP:OD2	1:A:427:LYS:NZ	2.50	0.42
1:A:419:LEU:HD12	1:A:454:TRP:CE3	2.55	0.41
1:A:404:GLU:HB2	4:A:703:HOH:O	2.21	0.41
1:A:77:TRP:O	1:A:81[A]:ILE:HG12	2.21	0.40
1:A:81[B]:ILE:HD11	1:A:95:VAL:CG2	2.51	0.40
1:A:256:THR:O	1:A:257:HIS:HB2	2.21	0.40
1:A:266:TYR:CD2	1:A:436:ALA:HB2	2.57	0.40

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:606:HOH:O	4:A:836:HOH:O[2_764]	1.72	0.48
4:A:624:HOH:O	4:A:713:HOH:O[3_755]	1.77	0.43
4:A:918:HOH:O	4:A:1003:HOH:O[4_466]	1.81	0.39
4:A:1007:HOH:O	4:A:1034:HOH:O[4_556]	1.82	0.38
4:A:1021:HOH:O	4:A:1022:HOH:O[2_665]	1.89	0.31
4:A:854:HOH:O	4:A:918:HOH:O[4_566]	1.90	0.30
4:A:787:HOH:O	4:A:923:HOH:O[2_665]	1.93	0.27
4:A:762:HOH:O	4:A:916:HOH:O[4_566]	2.02	0.18
4:A:667:HOH:O	4:A:887:HOH:O[2_665]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	428/444 (96%)	416 (97%)	10 (2%)	2 (0%)	29 11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	PRO
1	A	257	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	364/369 (99%)	362 (100%)	2 (0%)	88 80

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	339	TYR
1	A	345	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BMA	B	1	2	12,12,12	0.99	0	17,17,17	1.10	1 (5%)
2	BMA	B	2	2	11,11,12	1.25	1 (9%)	15,15,17	1.10	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BMA	B	1	2	-	0/2/22/22	0/1/1/1
2	BMA	B	2	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	BMA	O5-C1	-3.28	1.38	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	BMA	O2-C2-C3	-2.83	104.28	110.15
2	B	1	BMA	C1-O5-C5	-2.03	109.73	113.65


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	254:PRO	C	255:ASP	N	1.13

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/444 (96%)	0.49	17 (3%) 38 35	16, 19, 26, 31	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	35	GLY	4.3
1	A	48	GLY	4.2
1	A	390	THR	3.8
1	A	383	SER	3.1
1	A	73	PRO	2.9
1	A	36	ASN	2.8
1	A	330	LYS	2.6
1	A	382	GLU	2.6
1	A	47	LEU	2.5
1	A	71	THR	2.5
1	A	389	GLU	2.4
1	A	55	LYS	2.3
1	A	171	VAL	2.3
1	A	388	TYR	2.3
1	A	423	ASN	2.2
1	A	52	LYS	2.1
1	A	450	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	B	1	12/12	0.93	0.10	16,20,25,29	0
2	BMA	B	2	11/12	0.96	0.10	16,18,20,21	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	A	516	1/1	0.11	0.18	96,96,96,96	0
3	ZN	A	511	1/1	0.39	0.17	106,106,106,106	0
3	ZN	A	514	1/1	0.70	0.43	117,117,117,117	0
3	ZN	A	515	1/1	0.71	0.17	91,91,91,91	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ZN	A	503	1/1	0.74	0.19	67,67,67,67	0
3	ZN	A	512	1/1	0.75	0.16	76,76,76,76	0
3	ZN	A	513	1/1	0.83	0.26	100,100,100,100	0
3	ZN	A	510	1/1	0.95	0.08	21,21,21,21	1
3	ZN	A	509	1/1	0.99	0.06	17,17,17,17	1
3	ZN	A	504	1/1	0.99	0.13	37,37,37,37	0
3	ZN	A	507	1/1	0.99	0.06	19,19,19,19	0
3	ZN	A	508	1/1	0.99	0.08	23,23,23,23	1
3	ZN	A	505	1/1	1.00	0.10	20,20,20,20	0
3	ZN	A	506	1/1	1.00	0.11	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.