



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2024 – 07:47 AM EST

PDB ID : 3IJ9
Title : Directed 'in situ' Elongation as a Strategy to Characterize the Covalent Glycosyl-Enzyme Catalytic Intermediate of Human Pancreatic α -Amylase
Authors : Li, C.; Zhang, R.; Withers, S.G.; Brayer, G.D.
Deposited on : 2009-08-04
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	<div> <div>4%</div> <div>85%</div> <div>13%</div> <div>.</div> </div>
2	B	2	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BGC	B	2	X	-	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pancreatic alpha-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	496	3946	2497	696	733	20	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-2)-5-fluoro-alpha-L-idopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	F	O			
2	B	2	23	12	1	10	0	0	0

- Molecule 3 is (2R,3S,4R,5R,6R)-2,6-difluoro-2-(hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol (three-letter code: B0D) (formula: C₆H₁₀F₂O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	O		
3	A	1	13	6	2	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	F	O	0	0
			13	6	2	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	307	Total	O	0	0
			307	307		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pancreatic alpha-amylase



- Molecule 2: beta-D-glucopyranose-(1-2)-5-fluoro-alpha-L-idopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.38Å 68.00Å 129.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.38 – 1.85 33.38 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (33.38-1.85) 99.7 (33.38-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	18.20 (at 1.85Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.184 , 0.207 0.180 , 0.205	Depositor DCC
R_{free} test set	2018 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.588	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4304	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: B9D, BGC, PCA, CA, B0D, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/4053	0.53	0/5506

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	PCA	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3946	0	3718	47	0
2	B	23	0	17	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	26	0	0	0	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	307	0	0	1	0
All	All	4304	0	3735	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ASN:HD21	1:A:220:ASN:HD22	1.27	0.79
1:A:218:ASN:HD22	1:A:220:ASN:H	1.31	0.74
1:A:150:ASN:HD22	1:A:152:ASN:H	1.33	0.74
1:A:216:ASN:HD21	1:A:227:LYS:NZ	1.84	0.74
1:A:218:ASN:ND2	1:A:220:ASN:H	1.93	0.66
1:A:33:ALA:HB3	1:A:34:PRO:HD3	1.78	0.65
1:A:260:ALA:O	1:A:264:THR:HG23	1.99	0.62
1:A:150:ASN:ND2	1:A:152:ASN:H	1.98	0.62
1:A:458:ILE:HG22	1:A:459:ASN:OD1	2.01	0.60
1:A:218:ASN:HD21	1:A:220:ASN:ND2	2.00	0.58
1:A:216:ASN:HD21	1:A:227:LYS:HZ2	1.56	0.54
1:A:449:TYR:CZ	1:A:495:LYS:HD3	2.42	0.54
1:A:200:LYS:HE2	1:A:240:GLU:OE2	2.09	0.53
1:A:216:ASN:HD21	1:A:227:LYS:HZ3	1.54	0.52
1:A:439:THR:C	1:A:440:LEU:HD12	2.30	0.52
1:A:308:GLY:HA3	1:A:312:ILE:HD11	1.93	0.51
1:A:458:ILE:HD12	1:A:463:THR:HG21	1.93	0.51
1:A:218:ASN:ND2	1:A:220:ASN:HB2	2.27	0.49
1:A:404:GLN:OE1	1:A:422:GLY:HA3	2.13	0.49
1:A:306:GLY:HA3	6:A:668:HOH:O	2.12	0.49
1:A:235:ILE:CG2	1:A:307:ALA:HB2	2.43	0.49
1:A:380:ASN:O	1:A:381:ASP:HB2	2.12	0.48
1:A:142:LYS:HZ1	1:A:142:LYS:H	1.63	0.47
1:A:147:ASP:OD1	1:A:161:ARG:NH1	2.48	0.46
1:A:4:PRO:HA	1:A:229:PHE:CG	2.51	0.46
1:A:4:PRO:O	1:A:5:ASN:HB2	2.16	0.45
1:A:241:PRO:O	1:A:243:LYS:HE3	2.17	0.45
1:A:103:CYS:SG	1:A:121:PRO:HG3	2.57	0.45
1:A:150:ASN:ND2	1:A:152:ASN:HB2	2.31	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:MET:H	1:A:415:ASN:HD22	1.64	0.44
1:A:2:TYR:CE1	1:A:251:GLY:HA2	2.53	0.44
1:A:375:ASP:C	1:A:376:THR:HG23	2.38	0.44
1:A:211:LEU:HA	1:A:214:LEU:HD22	2.01	0.43
1:A:273:LYS:HA	1:A:415:ASN:HD21	1.84	0.42
1:A:86:CYS:O	1:A:89:VAL:HG12	2.18	0.42
1:A:200:LYS:CE	1:A:240:GLU:OE2	2.68	0.42
1:A:3:SER:HA	1:A:4:PRO:HD3	1.88	0.42
1:A:142:LYS:H	1:A:142:LYS:NZ	2.17	0.42
1:A:200:LYS:NZ	1:A:240:GLU:OE2	2.51	0.41
1:A:31:TYR:C	1:A:34:PRO:HD2	2.41	0.41
1:A:349:GLN:O	1:A:352:ASN:ND2	2.53	0.41
1:A:126:PHE:HB2	1:A:131:TYR:HB2	2.02	0.41
1:A:261:LYS:HD2	1:A:261:LYS:HA	1.75	0.41
1:A:235:ILE:HG21	1:A:307:ALA:HB2	2.02	0.41
1:A:235:ILE:HG23	1:A:307:ALA:HB2	2.03	0.41
1:A:31:TYR:O	1:A:35:LYS:HB2	2.20	0.40
1:A:27:GLU:HG3	1:A:32:LEU:HG	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	494/496 (100%)	476 (96%)	17 (3%)	1 (0%)	44 32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	239	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	418/418 (100%)	393 (94%)	25 (6%)	16 4

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LEU
1	A	89	VAL
1	A	115	CYS
1	A	142	LYS
1	A	150	ASN
1	A	165	LEU
1	A	208	LYS
1	A	214	LEU
1	A	218	ASN
1	A	243	LYS
1	A	270	ASN
1	A	273	LYS
1	A	303	ARG
1	A	305	HIS
1	A	323	MET
1	A	341	SER
1	A	347	GLN
1	A	352	ASN
1	A	369	GLU
1	A	411	ASP
1	A	444	LEU
1	A	459	ASN
1	A	472	ASP
1	A	495	LYS
1	A	496	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	150	ASN
1	A	152	ASN
1	A	185	HIS
1	A	216	ASN
1	A	218	ASN
1	A	270	ASN
1	A	364	ASN
1	A	415	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	1	1	7,8,9	2.03	3 (42%)	9,10,12	1.96	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	CB-CG	-3.38	1.45	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	CD-N	2.89	1.41	1.34
1	A	1	PCA	OE-CD	2.24	1.27	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CB-CG-CD	4.41	111.24	104.41
1	A	1	PCA	O-C-CA	-2.59	118.11	124.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	B9D	B	1	1,2	9,12,13	1.90	2 (22%)	14,18,20	0.92	1 (7%)
2	BGC	B	2	2	11,11,12	1.51	2 (18%)	15,15,17	1.15	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B9D	B	1	1,2	-	0/2/23/26	0/1/1/1
2	BGC	B	2	2	1/1/4/5	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	B9D	O5-C1	4.29	1.46	1.37
2	B	2	BGC	C2-C3	2.93	1.57	1.52
2	B	1	B9D	O5-C5	2.79	1.47	1.43
2	B	2	BGC	C4-C5	2.37	1.58	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	BGC	C1-O5-C5	3.62	117.04	112.19
2	B	1	B9D	O5-C1-C6	2.39	111.81	106.62

All (1) chirality outliers are listed below:


Mol	Chain	Res	Type	Atom
2	B	2	BGC	C1

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	B0D	A	504	-	10,13,13	3.33	6 (60%)	17,20,20	1.21	2 (11%)
3	B0D	A	503	-	10,13,13	3.77	4 (40%)	17,20,20	1.23	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	B0D	A	504	-	-	0/2/26/26	0/1/1/1
3	B0D	A	503	-	-	0/2/26/26	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	503	B0D	C2-C1	7.80	1.56	1.52
3	A	504	B0D	C2-C1	6.50	1.56	1.52
3	A	503	B0D	O5-C1	6.05	1.48	1.39
3	A	503	B0D	O5-C5	5.76	1.49	1.37
3	A	504	B0D	O5-C5	4.89	1.47	1.37
3	A	504	B0D	O5-C1	4.25	1.45	1.39
3	A	504	B0D	F1-C1	-2.94	1.35	1.39
3	A	504	B0D	C3-C4	2.73	1.58	1.53
3	A	504	B0D	C3-C2	2.12	1.57	1.52
3	A	503	B0D	F1-C1	-2.03	1.36	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	B0D	F1-C1-C2	2.92	111.49	108.32
3	A	504	B0D	O5-C5-C6	2.78	112.88	106.35
3	A	503	B0D	O5-C1-C2	-2.30	109.17	112.47
3	A	504	B0D	O5-C1-C2	-2.10	109.46	112.47
3	A	503	B0D	O5-C5-C6	2.08	111.25	106.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	495/496 (99%)	0.00	19 (3%)	44 47	12, 18, 31, 43	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	GLY	6.7
1	A	239	GLY	4.5
1	A	352	ASN	4.3
1	A	496	LEU	4.1
1	A	308	GLY	4.1
1	A	459	ASN	3.8
1	A	270	ASN	3.7
1	A	306	GLY	2.9
1	A	458	ILE	2.7
1	A	484	GLU	2.7
1	A	271	GLY	2.6
1	A	237	LEU	2.4
1	A	307	ALA	2.4
1	A	240	GLU	2.4
1	A	351	GLY	2.4
1	A	375	ASP	2.3
1	A	305	HIS	2.2
1	A	374	PRO	2.0
1	A	77	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	PCA	A	1	8/9	0.87	0.11	16,22,24,29	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BGC	B	2	11/12	0.74	0.16	30,31,34,34	0
2	B9D	B	1	12/13	0.85	0.11	23,24,26,28	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	B0D	A	503	13/13	0.52	0.21	44,45,46,47	0
3	B0D	A	504	13/13	0.74	0.17	32,40,44,44	0
4	CA	A	497	1/1	0.99	0.02	13,13,13,13	0
5	CL	A	498	1/1	0.99	0.02	15,15,15,15	0

6.5 Other polymers ⓘ

There are no such residues in this entry.