



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 8, 2025 – 12:12 PM JST

PDB ID : 9IQA / pdb\_00009iqa  
Title : structure of the oleate hydratase V206L-mutant from Staphylococcus aureus  
Authors : Xue, S.; Feng, T.  
Deposited on : 2024-07-12  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	592	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 89%, yellow 89%, yellow 99%, green 99%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>89%</span> <span>9%</span> <span>.</span> </div> </div>
1	B	592	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, orange 3%, orange 92%, yellow 92%, yellow 98%, green 98%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>3%</span> <span>92%</span> <span>6%</span> <span>..</span> </div> </div>
1	C	592	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 2%, orange 2%, orange 92%, yellow 92%, yellow 98%, green 98%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>92%</span> <span>6%</span> <span>.</span> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oleate hydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	579	Total	C	N	O	S	0	0	0
			4687	2991	772	902	22			
1	B	583	Total	C	N	O	S	0	0	0
			4719	3013	777	906	23			
1	C	582	Total	C	N	O	S	0	0	0
			4711	3008	776	905	22			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP A0A0D6GJV1
A	206	LEU	VAL	engineered mutation	UNP A0A0D6GJV1
B	0	HIS	-	expression tag	UNP A0A0D6GJV1
B	206	LEU	VAL	engineered mutation	UNP A0A0D6GJV1
C	0	HIS	-	expression tag	UNP A0A0D6GJV1
C	206	LEU	VAL	engineered mutation	UNP A0A0D6GJV1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	521	Total	O	0	0
			521	521		
2	B	522	Total	O	0	0
			522	522		
2	C	476	Total	O	0	0
			476	476		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Oleate hydratase



#### • Molecule 1: Oleate hydratase



#### • Molecule 1: Oleate hydratase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	189.18Å 113.06Å 118.97Å 90.00° 117.12° 90.00°	Depositor
Resolution (Å)	42.10 – 2.09 42.10 – 2.09	Depositor EDS
% Data completeness (in resolution range)	96.3 (42.10-2.09) 96.9 (42.10-2.09)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.16 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.162 , 0.188 0.163 , 0.188	Depositor DCC
$R_{free}$ test set	6649 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15636	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.10	0/4795	0.30	0/6497
1	B	0.10	0/4828	0.29	0/6541
1	C	0.11	0/4820	0.30	0/6530
All	All	0.10	0/14443	0.30	0/19568

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4687	0	4578	29	0
1	B	4719	0	4618	23	0
1	C	4711	0	4610	23	0
2	A	521	0	0	1	0
2	B	522	0	0	2	0
2	C	476	0	0	2	0
All	All	15636	0	13806	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:ARG:NH2	1:B:225:GLN:HE21	1.77	0.82
1:A:15:LYS:NZ	1:B:117:GLU:OE2	2.13	0.81
1:C:554:ASP:HB2	1:C:557:GLU:HG3	1.67	0.77
1:C:82:GLU:HG2	1:C:218:LEU:HD13	1.67	0.75
1:A:534:ASN:HD21	1:B:534:ASN:HD21	1.37	0.72
1:A:28:ILE:HD12	1:A:250:VAL:HG21	1.71	0.72
1:A:447:THR:HA	1:A:450:ILE:HD12	1.71	0.72
1:B:261:GLN:NE2	2:B:604:HOH:O	2.29	0.65
1:A:82:GLU:HG2	1:A:218:LEU:HD13	1.78	0.65
1:A:116:LYS:NZ	1:B:526:ASP:OD1	2.31	0.63
1:C:243:GLN:NE2	2:C:605:HOH:O	2.32	0.61
1:A:266:GLU:OE1	1:A:277:SER:OG	2.20	0.59
1:A:112:TYR:CZ	1:A:116:LYS:HE3	2.38	0.59
1:A:225:GLN:HG3	1:A:229:LEU:HD12	1.87	0.57
1:C:447:THR:HA	1:C:450:ILE:HD12	1.87	0.57
1:C:28:ILE:HD12	1:C:250:VAL:HG21	1.87	0.56
1:B:28:ILE:HD12	1:B:250:VAL:HG21	1.87	0.56
1:A:186:MET:HE3	1:A:215:PHE:HE1	1.71	0.55
1:A:554:ASP:HB2	1:A:557:GLU:HG3	1.89	0.55
1:A:113:TRP:O	1:A:117:GLU:HG2	2.06	0.54
1:C:331:ASN:OD1	2:C:601:HOH:O	2.18	0.54
1:A:82:GLU:HB3	1:A:218:LEU:HD22	1.90	0.53
1:B:81:ARG:NE	1:B:225:GLN:HG2	2.23	0.53
1:A:74:GLY:N	2:A:616:HOH:O	2.41	0.53
1:C:225:GLN:HG3	1:C:229:LEU:HD12	1.90	0.53
1:C:254:LYS:HA	1:C:254:LYS:HE2	1.91	0.51
1:C:32:LEU:HD22	1:C:233:MET:HE1	1.93	0.51
1:B:30:SER:HB2	1:B:56:GLU:OE1	2.11	0.50
1:B:554:ASP:HB2	1:B:557:GLU:HG3	1.94	0.48
1:C:341:LYS:HD3	1:C:341:LYS:HA	1.59	0.47
1:B:81:ARG:HE	1:B:225:GLN:HG2	1.79	0.47
1:C:556:ARG:HG3	1:C:556:ARG:HH11	1.80	0.47
1:C:562:SER:HB3	1:C:565:GLN:HB2	1.97	0.47
1:A:387:TRP:HB3	1:A:413:ALA:HB1	1.98	0.45
1:B:452:ASP:OD1	1:B:456:HIS:HD2	1.99	0.45
1:A:90:LEU:HA	1:A:512:VAL:HG11	1.98	0.45
1:B:77:VAL:HG23	1:B:410:TRP:CD1	2.53	0.44
1:C:319:TRP:CD2	1:C:332:PRO:HB3	2.52	0.44
1:C:112:TYR:CZ	1:C:116:LYS:HE3	2.51	0.44
1:B:320:LYS:NZ	2:B:628:HOH:O	2.51	0.44
1:A:389:MET:HE1	1:A:438:GLU:HG3	1.99	0.44
1:C:562:SER:O	1:C:566:LYS:HB2	2.18	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:GLU:H	1:A:21:GLU:CD	2.26	0.43
1:B:82:GLU:HG3	1:B:220:PHE:CZ	2.53	0.43
1:A:209:ILE:HG21	1:A:545:ALA:HA	1.99	0.43
1:B:32:LEU:HD22	1:B:233:MET:HE1	2.01	0.43
1:C:159:GLU:H	1:C:159:GLU:CD	2.27	0.43
1:A:32:LEU:HD22	1:A:233:MET:HE1	2.01	0.43
1:C:21:GLU:H	1:C:21:GLU:CD	2.27	0.43
1:B:87:PHE:CG	1:B:509:GLU:HB2	2.54	0.43
1:A:372:GLY:HA2	1:A:396:GLN:O	2.19	0.43
1:C:78:ARG:HB2	1:C:81:ARG:NH1	2.34	0.42
1:B:346:VAL:HA	1:B:463:VAL:O	2.19	0.42
1:B:319:TRP:CD2	1:B:332:PRO:HB3	2.55	0.42
1:C:201:TYR:HA	1:C:504:THR:HG21	2.02	0.42
1:A:319:TRP:CD2	1:A:332:PRO:HB3	2.55	0.41
1:A:209:ILE:HG23	1:A:548:GLU:HG3	2.02	0.41
1:C:90:LEU:HA	1:C:512:VAL:HG11	2.02	0.41
1:A:367:ARG:HD2	1:A:374:THR:O	2.20	0.41
1:B:158:GLU:HG3	1:B:196:MET:HG3	2.03	0.41
1:B:341:LYS:HA	1:B:341:LYS:HD3	1.84	0.41
1:A:82:GLU:HG3	1:A:220:PHE:CE2	2.55	0.41
1:A:87:PHE:CG	1:A:509:GLU:HB2	2.55	0.41
1:A:517:GLU:HG2	1:A:529:ILE:HD13	2.02	0.41
1:B:447:THR:HA	1:B:450:ILE:HD12	2.02	0.41
1:B:502:ARG:NH2	1:B:532:VAL:O	2.52	0.41
1:A:170:PHE:CE2	1:A:198:MET:HE2	2.56	0.41
1:B:535:SER:N	1:B:536:PRO:HD2	2.36	0.41
1:C:85:ASN:OD1	1:C:85:ASN:N	2.54	0.41
1:C:372:GLY:HA2	1:C:396:GLN:O	2.22	0.40
1:A:30:SER:HB2	1:A:56:GLU:OE1	2.20	0.40
1:C:222:LYS:HG2	1:C:223:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	575/592 (97%)	561 (98%)	14 (2%)	0	100	100
1	B	579/592 (98%)	566 (98%)	13 (2%)	0	100	100
1	C	578/592 (98%)	564 (98%)	14 (2%)	0	100	100
All	All	1732/1776 (98%)	1691 (98%)	41 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	519/528 (98%)	516 (99%)	3 (1%)	84	89
1	B	523/528 (99%)	516 (99%)	7 (1%)	65	72
1	C	522/528 (99%)	519 (99%)	3 (1%)	84	89
All	All	1564/1584 (99%)	1551 (99%)	13 (1%)	79	84

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	84	GLU
1	A	506	PHE
1	B	30	SER
1	B	32	LEU
1	B	84	GLU
1	B	477	ILE
1	B	535	SER
1	B	567	LEU
1	B	577	LYS
1	C	32	LEU
1	C	212	LEU
1	C	535	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	261	GLN
1	A	398	GLN
1	A	437	GLN
1	A	489	GLN
1	A	524	ASN
1	A	534	ASN
1	B	243	GLN
1	B	261	GLN
1	B	437	GLN
1	B	456	HIS
1	B	521	GLN
1	C	274	ASN
1	C	331	ASN
1	C	337	GLN
1	C	383	ASN
1	C	394	ASN
1	C	437	GLN
1	C	521	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	579/592 (97%)	-0.34	7 (1%) 76 77	17, 25, 41, 56	0
1	B	583/592 (98%)	-0.25	17 (2%) 54 55	16, 24, 48, 73	0
1	C	582/592 (98%)	-0.30	13 (2%) 62 64	16, 24, 47, 77	0
All	All	1744/1776 (98%)	-0.30	37 (2%) 63 65	16, 24, 45, 77	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	564	MET	5.1
1	B	29	GLY	5.1
1	B	80	GLY	4.6
1	A	80	GLY	3.9
1	C	80	GLY	3.8
1	A	554	ASP	3.7
1	B	72	LEU	3.5
1	B	567	LEU	3.4
1	C	71	PRO	3.4
1	B	578	GLY	3.3
1	C	29	GLY	3.3
1	B	246	TYR	3.3
1	B	563	LYS	3.2
1	B	554	ASP	3.2
1	C	564	MET	3.2
1	A	79	GLY	3.1
1	A	29	GLY	3.0
1	C	554	ASP	3.0
1	C	72	LEU	2.8
1	B	456	HIS	2.8
1	A	30	SER	2.7
1	B	565	GLN	2.7
1	C	578	GLY	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	30	SER	2.6
1	C	575	LYS	2.6
1	A	578	GLY	2.5
1	B	70	MET	2.5
1	B	71	PRO	2.5
1	C	567	LEU	2.4
1	C	30	SER	2.4
1	C	574	LYS	2.3
1	B	577	LYS	2.2
1	C	577	LYS	2.1
1	C	563	LYS	2.1
1	A	78	ARG	2.1
1	B	58	LEU	2.1
1	B	299	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.