



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 8, 2025 – 12:21 PM JST

PDB ID : 9IQZ / pdb_00009iqz
Title : phage HY126 glycosyltransferase
Authors : Yu, H.
Deposited on : 2024-07-13
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4623 (2.64-2.60)
Clashscore	180529	5071 (2.64-2.60)
Ramachandran outliers	177936	5006 (2.64-2.60)
Sidechain outliers	177891	5006 (2.64-2.60)
RSRZ outliers	164620	4622 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	 2% 72% 23% . .
1	H	369	 83% 13% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyltransferase subfamily 4-like N-terminal domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	366	Total	C	N	O	S	0	0	0
			2924	1885	487	538	14			
1	A	366	Total	C	N	O	S	0	0	0
			2924	1885	487	538	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	-1	GLY	-	expression tag	UNP A0A386K5C4
H	0	SER	-	expression tag	UNP A0A386K5C4
H	1	HIS	-	expression tag	UNP A0A386K5C4
H	316	PHE	SER	conflict	UNP A0A386K5C4
A	-1	GLY	-	expression tag	UNP A0A386K5C4
A	0	SER	-	expression tag	UNP A0A386K5C4
A	1	HIS	-	expression tag	UNP A0A386K5C4
A	316	PHE	SER	conflict	UNP A0A386K5C4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	109	Total	O	0	0
			109	109		
2	A	58	Total	O	0	0
			58	58		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycosyltransferase subfamily 4-like N-terminal domain-containing protein



- Molecule 1: Glycosyltransferase subfamily 4-like N-terminal domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.12Å 82.72Å 139.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.95 – 2.61 49.95 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.95-2.61) 99.9 (49.95-2.61)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.23 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.196 , 0.248 0.195 , 0.247	Depositor DCC
R_{free} test set	1248 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	44.7	Xtriage
Anisotropy	0.262	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 26.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6015	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/3001	0.55	0/4079
1	H	0.42	0/3001	0.61	0/4079
All	All	0.37	0/6002	0.58	0/8158

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2924	0	2916	51	0
1	H	2924	0	2916	39	0
2	A	58	0	0	1	0
2	H	109	0	0	2	0
All	All	6015	0	5832	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:68:SER:H	1:H:71:MET:HE3	1.30	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:111:LYS:HD3	1:H:111:LYS:H	1.43	0.82
1:H:148:PRO:HG3	1:H:275:THR:HG21	1.69	0.75
1:A:201:THR:O	1:A:203:LEU:N	2.23	0.72
1:H:272:ASP:OD1	1:H:273:THR:N	2.26	0.68
1:A:122:GLU:HG3	1:A:124:VAL:HG22	1.75	0.68
1:A:135:THR:HG22	1:A:180:LEU:HD13	1.79	0.64
1:H:213:LYS:HD2	1:H:213:LYS:H	1.63	0.63
1:H:125:SER:O	1:H:129:VAL:HG13	1.99	0.62
1:H:111:LYS:HD3	1:H:111:LYS:N	2.14	0.61
1:H:272:ASP:OD1	1:H:273:THR:HG22	2.02	0.59
1:H:196:GLU:H	1:H:196:GLU:CD	2.11	0.59
1:A:148:PRO:HD2	1:A:152:PHE:HD2	1.67	0.59
1:A:268:ASN:HD21	1:A:277:GLU:HG2	1.67	0.59
1:A:122:GLU:HG2	1:A:214:MET:HE2	1.87	0.57
1:H:148:PRO:CG	1:H:275:THR:HG21	2.34	0.56
1:A:236:PRO:HG2	1:A:239:TRP:HB2	1.88	0.56
1:A:209:ARG:NE	1:A:236:PRO:HD3	2.21	0.56
1:A:121:PHE:HE1	1:A:129:VAL:HA	1.71	0.55
1:H:274:GLY:HA3	1:H:300:HIS:CE1	2.41	0.54
1:A:10:ARG:HD2	1:A:67:PHE:CD2	2.42	0.54
1:A:223:ALA:O	1:A:227:LEU:HD22	2.08	0.54
1:A:22:GLY:O	1:A:26:VAL:HG23	2.07	0.54
1:A:215:ARG:HD2	1:A:270:CYS:SG	2.48	0.54
1:A:205:ILE:HG21	1:A:256:ILE:HD11	1.91	0.53
1:A:209:ARG:HD3	1:A:236:PRO:HG3	1.90	0.53
1:H:20:GLN:HG3	2:H:468:HOH:O	2.09	0.53
1:A:151:ALA:N	1:A:304:GLU:OE2	2.39	0.53
1:A:227:LEU:HD13	1:A:332:LEU:HD23	1.91	0.52
1:A:45:GLY:O	1:A:60:PRO:HA	2.10	0.52
1:H:213:LYS:H	1:H:213:LYS:CD	2.23	0.52
1:A:273:THR:O	1:A:275:THR:N	2.43	0.52
1:H:2:MET:HE2	1:H:90:PHE:HE2	1.75	0.51
1:H:275:THR:HB	2:H:415:HOH:O	2.10	0.51
1:A:269:THR:O	1:A:271:PRO:HD3	2.11	0.51
1:A:274:GLY:HA2	1:A:277:GLU:OE1	2.10	0.51
1:A:296:LYS:HD3	1:A:313:GLU:HG2	1.93	0.50
1:H:7:ILE:HB	1:H:93:THR:HG22	1.94	0.50
1:A:209:ARG:HE	1:A:236:PRO:HD3	1.76	0.50
1:H:27:HIS:CD2	1:H:27:HIS:C	2.89	0.50
1:A:27:HIS:HB2	1:A:94:MET:HE3	1.93	0.49
1:A:310:VAL:HB	2:A:449:HOH:O	2.11	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:209:ARG:HB2	1:H:239:TRP:HB2	1.93	0.49
1:H:212:THR:HB	1:H:213:LYS:HE3	1.94	0.48
1:H:111:LYS:H	1:H:111:LYS:CD	2.18	0.48
1:H:19:VAL:HG13	1:H:253:HIS:CD2	2.48	0.48
1:A:270:CYS:HB3	1:A:273:THR:OG1	2.14	0.48
1:H:155:PHE:HB3	1:H:159:TYR:CZ	2.49	0.47
1:H:208:GLN:NE2	1:H:215:ARG:HB3	2.30	0.47
1:A:315:ASP:OD2	1:A:317:SER:OG	2.26	0.47
1:A:270:CYS:O	1:A:300:HIS:NE2	2.43	0.46
1:A:157:ARG:NE	1:A:157:ARG:HA	2.31	0.46
1:A:200:ALA:HB1	1:A:264:LYS:HA	1.97	0.46
1:A:125:SER:O	1:A:129:VAL:HG12	2.17	0.45
1:A:218:ASP:OD1	1:A:218:ASP:N	2.49	0.45
1:A:241:PRO:HB2	1:A:245:TYR:HB2	1.99	0.45
1:A:295:PHE:HA	1:A:314:ILE:O	2.17	0.45
1:H:94:MET:HB2	1:H:94:MET:HE2	1.52	0.44
1:A:72:SER:OG	1:A:99:LEU:HD12	2.18	0.44
1:A:198:LEU:HD12	1:A:261:LYS:HB3	2.00	0.44
1:H:269:THR:O	1:H:271:PRO:HD3	2.17	0.43
1:A:196:GLU:H	1:A:196:GLU:CD	2.25	0.43
1:H:236:PRO:HG2	1:H:239:TRP:HE3	1.84	0.43
1:H:267:ILE:HG23	1:H:327:TYR:OH	2.19	0.43
1:A:242:PRO:HB2	1:A:244:LYS:HG2	2.00	0.43
1:H:30:VAL:O	1:H:34:LEU:HG	2.19	0.42
1:H:153:ARG:NH2	1:H:185:ASP:OD1	2.49	0.42
1:H:157:ARG:NE	1:H:157:ARG:HA	2.35	0.42
1:H:68:SER:N	1:H:71:MET:HE3	2.14	0.42
1:H:296:LYS:HG3	1:H:313:GLU:HB3	2.01	0.42
1:A:14:PHE:CZ	1:A:31:LEU:HD23	2.55	0.42
1:A:27:HIS:CD2	1:A:27:HIS:C	2.98	0.42
1:A:320:LYS:O	1:A:323:VAL:HG22	2.20	0.42
1:A:215:ARG:NH1	1:A:270:CYS:SG	2.83	0.42
1:H:321:LYS:HD3	1:H:321:LYS:C	2.45	0.42
1:A:89:ASP:O	1:A:112:VAL:HB	2.20	0.42
1:H:110:SER:HB3	1:H:141:ASN:O	2.20	0.41
1:H:111:LYS:N	1:H:111:LYS:CD	2.82	0.41
1:A:209:ARG:H	1:A:209:ARG:HG2	1.56	0.41
1:A:312:VAL:HG21	1:A:327:TYR:CD1	2.55	0.41
1:H:215:ARG:HG2	1:H:270:CYS:SG	2.61	0.41
1:A:83:ALA:CB	1:A:107:ALA:HB1	2.49	0.41
1:A:61:VAL:O	1:A:71:MET:HG2	2.20	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:SER:HB3	1:A:128:ILE:HB	2.03	0.40
1:A:163:THR:HG22	1:A:172:TYR:CZ	2.56	0.40
1:A:321:LYS:HA	1:A:321:LYS:HE3	2.03	0.40
1:H:66:LYS:HA	1:H:66:LYS:HD2	1.95	0.40
1:H:148:PRO:HB3	1:H:186:TYR:CZ	2.57	0.40
1:H:201:THR:O	1:H:264:LYS:HB2	2.22	0.40
1:A:46:PHE:CE2	1:A:64:LEU:HD22	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/369 (99%)	349 (96%)	12 (3%)	3 (1%)	16	32
1	H	364/369 (99%)	349 (96%)	14 (4%)	1 (0%)	37	57
All	All	728/738 (99%)	698 (96%)	26 (4%)	4 (0%)	25	45

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	300	HIS
1	A	202	ASP
1	A	274	GLY
1	A	250	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/318 (100%)	286 (90%)	31 (10%)	6	12
1	H	317/318 (100%)	298 (94%)	19 (6%)	16	32
All	All	634/636 (100%)	584 (92%)	50 (8%)	10	20

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	19	VAL
1	H	87	GLU
1	H	94	MET
1	H	105	LYS
1	H	111	LYS
1	H	129	VAL
1	H	170	ILE
1	H	196	GLU
1	H	209	ARG
1	H	211	ASP
1	H	213	LYS
1	H	215	ARG
1	H	225	LYS
1	H	229	GLU
1	H	273	THR
1	H	275	THR
1	H	335	THR
1	H	348	VAL
1	H	354	ARG
1	A	12	VAL
1	A	17	ASP
1	A	19	VAL
1	A	23	LEU
1	A	28	LEU
1	A	32	LYS
1	A	55	LYS
1	A	56	VAL
1	A	72	SER
1	A	87	GLU
1	A	94	MET
1	A	99	LEU
1	A	162	MET
1	A	168	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	169	LYS
1	A	196	GLU
1	A	208	GLN
1	A	209	ARG
1	A	227	LEU
1	A	250	ASP
1	A	256	ILE
1	A	259	ARG
1	A	273	THR
1	A	276	VAL
1	A	296	LYS
1	A	315	ASP
1	A	321	LYS
1	A	324	VAL
1	A	335	THR
1	A	343	LYS
1	A	354	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	20	GLN
1	H	63	HIS
1	H	134	GLN
1	H	137	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	366/369 (99%)	0.12	8 (2%) 62 57	33, 54, 88, 104	0
1	H	366/369 (99%)	-0.53	0 100 100	29, 35, 47, 61	0
All	All	732/738 (99%)	-0.20	8 (1%) 77 74	29, 40, 83, 104	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	256	ILE	4.2
1	A	165	ALA	3.7
1	A	1	HIS	2.8
1	A	166	LEU	2.5
1	A	224	ILE	2.4
1	A	249	ILE	2.2
1	A	253	HIS	2.1
1	A	214	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.