



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2025 – 12:38 AM JST

PDB ID : 9JLV / pdb_00009jlv
Title : Crystal structure of GH57 family amylopullulanase from Aquifex aeolicus mutant E256Q in complex with alpha-cyclodextrin
Authors : Zhu, Z.M.; Wang, W.W.; Li, M.J.; Xu, Q.; Zhou, H.; Huang, L.Q.; Wang, Q.S.; Yu, F.
Deposited on : 2024-09-19
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5293 (1.54-1.50)
Clashscore	180529	5759 (1.54-1.50)
Ramachandran outliers	177936	5653 (1.54-1.50)
Sidechain outliers	177891	5650 (1.54-1.50)
RSRZ outliers	164620	5293 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	477	<div> <div></div> <div>92%8%</div> </div>
2	B	4	<div> <div>25%25%50%</div> </div>
3	C	6	<div> <div>83%17%</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4555 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase family 57 N-terminal domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	477	Total	C	N	O	S	0	2	0
			4066	2658	652	746	10			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	256	GLN	GLU	engineered mutation	UNP O66934

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	4	Total	C	O	0	0	0
			45	24	21			

- Molecule 3 is an oligosaccharide called Cyclohexakis-(1-4)-(alpha-D-glucopyranose).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	6	Total	C	O	0	0	0
			66	36	30			

- Molecule 4 is CITRATE ANION (CCD ID: FLC) (formula: C₆H₅O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		
4	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	340	Total	O	0	0
			340	340		

- Molecule 1: Glycoside hydrolase family 57 N-terminal domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	199.97Å 40.57Å 62.62Å 90.00° 102.82° 90.00°	Depositor
Resolution (Å)	25.22 – 1.51 25.22 – 1.51	Depositor EDS
% Data completeness (in resolution range)	99.2 (25.22-1.51) 99.2 (25.22-1.51)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.51Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.164 , 0.184 0.164 , 0.183	Depositor DCC
R_{free} test set	3887 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4555	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, GLC, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	1/4179 (0.0%)	0.50	1/5652 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	256	GLN	C-O	-6.79	1.14	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	379	GLU	CB-CG-CD	-5.49	103.26	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4066	0	3998	40	0
2	B	45	0	39	5	0
3	C	66	0	54	1	0
4	A	26	0	10	0	0
5	A	12	0	16	0	0
6	A	340	0	0	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4555	0	4117	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:LYS:NZ	6:A:604:HOH:O	2.09	0.84
1:A:181:ARG:NH1	6:A:602:HOH:O	2.02	0.83
1:A:149:ASP:OD1	6:A:601:HOH:O	2.00	0.78
1:A:396:LYS:HE3	6:A:906:HOH:O	1.84	0.77
1:A:181:ARG:NH2	6:A:602:HOH:O	2.12	0.75
1:A:1:MET:HE3	1:A:3:LYS:HE2	1.67	0.75
1:A:46:LYS:NZ	6:A:608:HOH:O	2.22	0.72
1:A:325:ASP:OD1	6:A:605:HOH:O	2.09	0.70
1:A:469:ARG:NH1	6:A:609:HOH:O	2.25	0.65
1:A:29:TRP:CD2	2:B:3:GLC:H2	2.33	0.63
1:A:474:ARG:NE	6:A:611:HOH:O	2.31	0.62
1:A:420:LYS:HE2	1:A:420:LYS:H	1.67	0.58
1:A:181:ARG:CZ	6:A:602:HOH:O	2.38	0.58
1:A:459:PRO:O	6:A:606:HOH:O	2.18	0.54
1:A:378:GLU:OE2	6:A:607:HOH:O	2.19	0.53
1:A:65:GLN:O	1:A:69:GLN:HG3	2.09	0.52
1:A:420:LYS:H	1:A:420:LYS:CE	2.23	0.51
1:A:148:LYS:NZ	6:A:616:HOH:O	2.39	0.51
1:A:1:MET:HE3	1:A:3:LYS:CE	2.39	0.51
2:B:1:GLC:O1	3:C:1:GLC:H61	2.10	0.50
1:A:120:GLU:H	1:A:120:GLU:CD	2.20	0.50
1:A:148:LYS:NZ	1:A:149:ASP:OD1	2.44	0.49
1:A:335:LYS:HG2	1:A:380:TRP:CZ2	2.49	0.48
1:A:115:GLU:OE2	1:A:155:ARG:NH1	2.39	0.48
1:A:256:GLN:HE22	2:B:1:GLC:C1	2.27	0.47
1:A:282:LEU:HB2	1:A:305:PHE:CD2	2.50	0.46
1:A:101:LEU:HD13	1:A:459:PRO:HG2	1.97	0.46
1:A:420:LYS:H	1:A:420:LYS:CD	2.29	0.46
1:A:115:GLU:CD	1:A:155:ARG:HH12	2.21	0.45
1:A:182:GLU:O	1:A:185:GLU:HG2	2.16	0.45
1:A:251:TYR:OH	1:A:387:GLU:HA	2.17	0.44
1:A:251:TYR:OH	1:A:387:GLU:CD	2.61	0.43
1:A:256:GLN:HE22	2:B:1:GLC:H1	1.83	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:LEU:O	1:A:383:THR:HA	2.19	0.42
1:A:331:ILE:O	1:A:335:LYS:HG3	2.19	0.42
1:A:171:GLU:O	1:A:175:LYS:HG3	2.19	0.42
1:A:29:TRP:CE2	2:B:3:GLC:H2	2.55	0.42
1:A:380:TRP:CD1	1:A:380:TRP:H	2.39	0.41
1:A:111:GLU:OE1	1:A:114:ARG:NH2	2.52	0.41
1:A:65:GLN:O	1:A:69:GLN:CG	2.70	0.40
1:A:145:LYS:NZ	6:A:628:HOH:O	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	477/477 (100%)	472 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	439/437 (100%)	439 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	476	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	B	1	2	12,12,12	0.88	0	17,17,17	1.12	1 (5%)
2	GLC	B	2	2	11,11,12	0.46	0	15,15,17	0.95	1 (6%)
2	GLC	B	3	2	11,11,12	0.47	0	15,15,17	0.98	1 (6%)
2	GLC	B	4	2	11,11,12	0.63	0	15,15,17	0.76	0
3	GLC	C	1	3	11,11,12	1.43	2 (18%)	15,15,17	1.81	3 (20%)
3	GLC	C	2	3	11,11,12	0.85	0	15,15,17	1.60	4 (26%)
3	GLC	C	3	3	11,11,12	1.34	2 (18%)	15,15,17	1.30	3 (20%)
3	GLC	C	4	3	11,11,12	1.23	2 (18%)	15,15,17	1.55	4 (26%)
3	GLC	C	5	3	11,11,12	1.45	1 (9%)	15,15,17	1.46	3 (20%)
3	GLC	C	6	3	11,11,12	1.48	3 (27%)	15,15,17	2.08	8 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	0/2/19/22	0/1/1/1
2	GLC	B	4	2	-	2/2/19/22	0/1/1/1
3	GLC	C	1	3	-	0/2/19/22	0/1/1/1
3	GLC	C	2	3	-	0/2/19/22	0/1/1/1
3	GLC	C	3	3	-	0/2/19/22	0/1/1/1
3	GLC	C	4	3	-	0/2/19/22	0/1/1/1
3	GLC	C	5	3	-	2/2/19/22	0/1/1/1
3	GLC	C	6	3	-	0/2/19/22	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	GLC	O5-C1	-3.42	1.38	1.43
3	C	3	GLC	O5-C1	-3.39	1.38	1.43
3	C	5	GLC	O5-C1	-3.18	1.38	1.43
3	C	4	GLC	O5-C1	-3.06	1.38	1.43
3	C	6	GLC	O4-C4	-2.52	1.37	1.43
3	C	3	GLC	O5-C5	-2.29	1.38	1.43
3	C	6	GLC	O5-C5	-2.10	1.39	1.43
3	C	1	GLC	O2-C2	-2.07	1.39	1.43
3	C	6	GLC	O5-C1	-2.06	1.40	1.43
3	C	4	GLC	O5-C5	-2.00	1.39	1.43

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	GLC	O5-C1-C2	-4.33	104.08	110.77
3	C	6	GLC	O4-C4-C3	-4.23	100.57	110.35
2	B	3	GLC	C1-O5-C5	3.35	116.73	112.19
3	C	5	GLC	C1-O5-C5	3.30	116.67	112.19
3	C	2	GLC	C1-O5-C5	3.20	116.52	112.19
3	C	6	GLC	C1-O5-C5	3.08	116.37	112.19
3	C	1	GLC	C1-C2-C3	-2.96	106.03	109.67
3	C	6	GLC	O5-C1-C2	-2.83	106.40	110.77
3	C	1	GLC	C6-C5-C4	-2.74	106.60	113.00
3	C	4	GLC	O2-C2-C1	2.70	114.68	109.15
3	C	5	GLC	O2-C2-C3	-2.57	104.98	110.14
3	C	6	GLC	C6-C5-C4	-2.54	107.06	113.00
3	C	3	GLC	O6-C6-C5	-2.54	102.59	111.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	4	GLC	O4-C4-C3	-2.53	104.50	110.35
3	C	4	GLC	C1-O5-C5	2.50	115.58	112.19
3	C	4	GLC	O2-C2-C3	-2.41	105.31	110.14
2	B	1	GLC	C1-O5-C5	2.36	118.12	113.66
3	C	6	GLC	O3-C3-C4	-2.35	104.91	110.35
3	C	2	GLC	C6-C5-C4	-2.32	107.56	113.00
2	B	2	GLC	C1-O5-C5	2.26	115.25	112.19
3	C	2	GLC	O5-C1-C2	-2.23	107.32	110.77
3	C	2	GLC	C3-C4-C5	-2.17	106.36	110.24
3	C	3	GLC	O4-C4-C3	-2.16	105.35	110.35
3	C	5	GLC	C1-C2-C3	-2.14	107.04	109.67
3	C	6	GLC	O3-C3-C2	2.08	113.98	109.99
3	C	3	GLC	O5-C5-C6	-2.05	103.98	107.20
3	C	6	GLC	C3-C4-C5	2.02	113.85	110.24
3	C	6	GLC	O5-C5-C6	-2.01	104.05	107.20

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	4	GLC	O5-C5-C6-O6
2	B	4	GLC	C4-C5-C6-O6
3	C	5	GLC	C4-C5-C6-O6
3	C	5	GLC	O5-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	GLC	2	0
2	B	1	GLC	3	0
3	C	1	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	504	-	5,5,5	1.03	0	5,5,5	1.10	0
4	FLC	A	501	-	12,12,12	1.29	1 (8%)	17,17,17	1.41	3 (17%)
5	GOL	A	503	-	5,5,5	0.88	0	5,5,5	1.02	0
4	FLC	A	502	-	12,12,12	1.11	1 (8%)	17,17,17	1.22	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	504	-	-	0/4/4/4	-
4	FLC	A	501	-	-	4/16/16/16	-
5	GOL	A	503	-	-	2/4/4/4	-
4	FLC	A	502	-	-	0/16/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	FLC	CB-CBC	-3.05	1.50	1.53
4	A	502	FLC	CB-CBC	-2.08	1.51	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	FLC	OB2-CBC-CB	3.12	118.46	113.05
4	A	501	FLC	OB1-CBC-CB	-2.86	118.21	122.25
4	A	502	FLC	OB1-CBC-CB	-2.67	118.48	122.25
4	A	501	FLC	OB2-CBC-CB	2.64	117.63	113.05
4	A	501	FLC	OA1-CAC-CA	-2.14	116.69	122.94

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	GOL	O1-C1-C2-C3
5	A	503	GOL	O1-C1-C2-O2
4	A	501	FLC	CB-CG-CGC-OG2
4	A	501	FLC	CB-CG-CGC-OG1
4	A	501	FLC	CAC-CA-CB-CG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	501	FLC	CAC-CA-CB-CBC

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	477/477 (100%)	-0.20	5 (1%) 79 82	8, 19, 38, 65	2 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	458	ALA	4.4
1	A	460	PHE	3.5
1	A	459	PRO	2.8
1	A	457	LYS	2.6
1	A	461	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLC	C	3	11/12	0.89	0.10	22,25,38,42	0
3	GLC	C	4	11/12	0.91	0.09	24,27,35,39	0
3	GLC	C	5	11/12	0.91	0.09	20,29,36,41	0
2	GLC	B	4	11/12	0.92	0.09	19,27,43,50	0
3	GLC	C	2	11/12	0.95	0.07	14,17,25,30	0
3	GLC	C	1	11/12	0.95	0.07	15,18,25,38	0
3	GLC	C	6	11/12	0.95	0.07	16,19,23,30	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	B	1	12/12	0.96	0.07	11,13,19,30	0
2	GLC	B	2	11/12	0.97	0.04	10,11,12,12	0
2	GLC	B	3	11/12	0.98	0.04	11,13,16,16	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	A	503	6/6	0.80	0.15	37,42,51,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	FLC	A	501	13/13	0.84	0.12	29,37,49,50	0
4	FLC	A	502	13/13	0.91	0.08	26,30,32,34	0
5	GOL	A	504	6/6	0.94	0.10	14,28,30,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.