



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 11:26 PM EDT

PDB ID : 5JX5
Title : GH6 Orpinomyces sp. Y102 enzyme
Authors : Tsai, L.C.; Huang, H.C.
Deposited on : 2016-05-12
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	322	<div> <div>8%</div> <div>82%</div> <div>15%</div> <div>.</div> </div>
1	B	322	<div> <div>9%</div> <div>85%</div> <div>13%</div> <div>.</div> </div>
1	C	322	<div> <div>9%</div> <div>77%</div> <div>18%</div> <div>..</div> </div>
1	D	322	<div> <div>8%</div> <div>84%</div> <div>13%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	C	501	-	-	X	-
3	GOL	C	502	-	-	X	-
4	EDO	A	507	-	-	X	-
4	EDO	A	510	-	-	X	-
4	EDO	B	502	-	-	X	-
4	EDO	B	507	-	-	X	-
7	PEG	C	506	-	-	X	-
7	PEG	C	507	-	-	X	-

2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 11278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	5	0
			2507	1560	449	488	10			
1	B	322	Total	C	N	O	S	0	3	0
			2503	1559	448	486	10			
1	C	322	Total	C	N	O	S	0	3	0
			2501	1558	446	486	11			
1	D	322	Total	C	N	O	S	0	6	0
			2509	1562	448	489	10			

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	O P	0	0
			5	4 1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is AMMONIUM ION (three-letter code: NH4) (formula: H₄N).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	N	0	0
			1	1		
5	A	1	Total	N	0	0
			1	1		
5	A	1	Total	N	0	0
			1	1		
5	A	1	Total	N	0	0
			1	1		
5	B	1	Total	N	0	0
			1	1		
5	B	1	Total	N	0	0
			1	1		
5	C	1	Total	N	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total N 1 1	0	0
5	C	1	Total N 1 1	0	0
5	C	1	Total N 1 1	0	0
5	C	1	Total N 1 1	0	0
5	D	1	Total N 1 1	0	0
5	D	1	Total N 1 1	0	0

- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C O 8 6 2	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			7	4	3		
7	C	1	Total	C	O	0	0
			7	4	3		
7	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	275	Total 275	O 275	0	0
9	B	274	Total 274	O 274	0	0
9	C	273	Total 273	O 273	0	0
9	D	271	Total 271	O 271	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucanase




- Molecule 1: Glucanase




- Molecule 1: Glucanase



- Molecule 1: Glucanase





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	309.86Å 87.35Å 82.55Å 90.00° 93.95° 90.00°	Depositor
Resolution (Å)	27.76 – 1.80 27.76 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.3 (27.76-1.80) 98.3 (27.76-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.209 , 0.239 0.209 , 0.239	Depositor DCC
R_{free} test set	10048 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11278	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, EDO, MPD, GOL, PEG, ACT, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	6/2592 (0.2%)	1.10	5/3523 (0.1%)
1	B	1.30	3/2577 (0.1%)	1.10	4/3503 (0.1%)
1	C	1.25	5/2574 (0.2%)	1.07	7/3498 (0.2%)
1	D	1.25	4/2598 (0.2%)	1.06	1/3532 (0.0%)
All	All	1.29	18/10341 (0.2%)	1.09	17/14056 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	1
All	All	0	3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	173	TRP	CD2-CE2	5.96	1.48	1.41
1	C	176	TRP	CD2-CE2	5.95	1.48	1.41
1	C	296	TRP	CD2-CE2	5.75	1.48	1.41
1	C	330	TYR	CG-CD2	5.74	1.46	1.39
1	A	337	TYR	CE1-CZ	5.66	1.46	1.38

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	174	LEU	CB-CG-CD1	5.97	121.15	111.00
1	B	214	GLY	N-CA-C	5.76	127.51	113.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	330	TYR	N-CA-CB	-5.73	100.29	110.60
1	C	390	ASP	CB-CG-OD2	-5.72	113.15	118.30
1	C	384	ARG	NE-CZ-NH2	-5.62	117.49	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	216	ALA	Peptide
1	A	257	ASN	Peptide
1	C	207	CYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2507	0	2391	99	0
1	B	2503	0	2390	80	0
1	C	2501	0	2396	75	0
1	D	2509	0	2400	52	0
2	A	5	0	0	1	0
3	A	12	0	16	4	0
3	B	6	0	8	2	0
3	C	12	0	16	13	0
3	D	12	0	16	3	0
4	A	40	0	60	36	0
4	B	12	0	18	19	0
4	C	12	0	18	3	0
4	D	8	0	12	2	0
5	A	4	0	0	0	0
5	B	2	0	0	0	0
5	C	5	0	0	0	0
5	D	2	0	0	0	0
6	B	8	0	14	2	0
7	C	14	0	20	11	0
7	D	7	0	10	2	0
8	C	4	0	3	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	275	0	0	16	0
9	B	274	0	0	12	0
9	C	273	0	0	11	0
9	D	271	0	0	5	0
All	All	11278	0	9788	320	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 320 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:THR:HG22	1:B:213:ASN:CB	1.65	1.23
1:A:211:GLY:CA	1:A:214:GLY:HA2	1.70	1.22
1:A:211:GLY:HA2	1:A:213:ASN:HA	1.25	1.18
1:A:248:ASP:HA	9:A:665:HOH:O	1.47	1.15
1:D:240:ARG:HG3	1:D:240:ARG:HH11	1.12	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/322 (101%)	313 (96%)	12 (4%)	0	100	100
1	B	323/322 (100%)	306 (95%)	17 (5%)	0	100	100
1	C	323/322 (100%)	309 (96%)	13 (4%)	1 (0%)	41	27
1	D	326/322 (101%)	315 (97%)	11 (3%)	0	100	100
All	All	1297/1288 (101%)	1243 (96%)	53 (4%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	205	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/266 (102%)	258 (95%)	13 (5%)	25	11
1	B	269/266 (101%)	258 (96%)	11 (4%)	30	16
1	C	269/266 (101%)	253 (94%)	16 (6%)	19	7
1	D	272/266 (102%)	259 (95%)	13 (5%)	25	11
All	All	1081/1064 (102%)	1028 (95%)	53 (5%)	27	11

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	205	ARG
1	C	265	HIS
1	D	313[B]	ARG
1	C	206	ASP
1	C	240	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	386	GLN
1	D	263	ASN
1	C	420	ASN
1	D	218	ASN
1	D	331	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 44 ligands modelled in this entry, 13 are modelled with single atom - leaving 31 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	C	501	-	5,5,5	0.45	0	5,5,5	0.59	0
3	GOL	D	504	-	5,5,5	0.52	0	5,5,5	0.65	0
4	EDO	A	508	-	3,3,3	0.56	0	2,2,2	0.18	0
4	EDO	A	512	-	3,3,3	0.28	0	2,2,2	1.56	1 (50%)
4	EDO	A	509	-	3,3,3	0.68	0	2,2,2	1.45	0
4	EDO	C	505	-	3,3,3	0.47	0	2,2,2	0.62	0
4	EDO	A	517	-	3,3,3	0.57	0	2,2,2	1.23	0
4	EDO	A	504	-	3,3,3	0.71	0	2,2,2	0.64	0
7	PEG	C	507	-	6,6,6	1.23	0	5,5,5	1.87	1 (20%)
4	EDO	D	502	-	3,3,3	0.93	0	2,2,2	0.32	0
3	GOL	C	502	-	5,5,5	0.45	0	5,5,5	0.68	0
4	EDO	C	503	-	3,3,3	0.79	0	2,2,2	0.10	0
7	PEG	C	506	-	6,6,6	0.63	0	5,5,5	0.81	0
8	ACT	C	513	-	3,3,3	0.93	0	3,3,3	0.53	0
4	EDO	A	506	-	3,3,3	0.67	0	2,2,2	0.64	0
4	EDO	B	507	-	3,3,3	0.58	0	2,2,2	0.87	0
4	EDO	B	502	-	3,3,3	0.73	0	2,2,2	0.78	0
6	MPD	B	504	-	7,7,7	1.53	1 (14%)	9,10,10	1.82	3 (33%)
3	GOL	D	501	-	5,5,5	0.54	0	5,5,5	1.25	0
4	EDO	A	511	-	3,3,3	0.37	0	2,2,2	1.36	0
4	EDO	B	503	-	3,3,3	0.60	0	2,2,2	0.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	502	-	5,5,5	0.70	0	5,5,5	0.59	0
2	PO4	A	501	-	4,4,4	0.38	0	6,6,6	1.26	0
3	GOL	A	503	-	5,5,5	0.78	0	5,5,5	1.49	1 (20%)
4	EDO	A	505	-	3,3,3	0.29	0	2,2,2	0.52	0
4	EDO	A	507	-	3,3,3	1.39	1 (33%)	2,2,2	1.33	0
4	EDO	D	503	-	3,3,3	0.56	0	2,2,2	0.54	0
7	PEG	D	505	-	6,6,6	0.94	0	5,5,5	1.65	2 (40%)
3	GOL	B	501	-	5,5,5	0.58	0	5,5,5	0.93	0
4	EDO	A	510	-	3,3,3	0.42	0	2,2,2	0.41	0
4	EDO	C	504	-	3,3,3	0.47	0	2,2,2	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	501	-	-	3/4/4/4	-
3	GOL	D	504	-	-	2/4/4/4	-
4	EDO	A	508	-	-	1/1/1/1	-
4	EDO	A	512	-	-	1/1/1/1	-
4	EDO	A	509	-	-	1/1/1/1	-
4	EDO	C	505	-	-	1/1/1/1	-
4	EDO	A	517	-	-	1/1/1/1	-
4	EDO	A	504	-	-	1/1/1/1	-
7	PEG	C	507	-	-	3/4/4/4	-
4	EDO	D	502	-	-	0/1/1/1	-
3	GOL	C	502	-	-	2/4/4/4	-
4	EDO	C	503	-	-	0/1/1/1	-
7	PEG	C	506	-	-	3/4/4/4	-
4	EDO	A	506	-	-	1/1/1/1	-
4	EDO	B	507	-	-	0/1/1/1	-
4	EDO	B	502	-	-	1/1/1/1	-
6	MPD	B	504	-	-	2/5/5/5	-
3	GOL	D	501	-	-	2/4/4/4	-
4	EDO	A	511	-	-	0/1/1/1	-
4	EDO	B	503	-	-	0/1/1/1	-
3	GOL	A	502	-	-	4/4/4/4	-
3	GOL	A	503	-	-	3/4/4/4	-
4	EDO	A	505	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	507	-	-	1/1/1/1	-
4	EDO	D	503	-	-	1/1/1/1	-
7	PEG	D	505	-	-	2/4/4/4	-
3	GOL	B	501	-	-	0/4/4/4	-
4	EDO	A	510	-	-	1/1/1/1	-
4	EDO	C	504	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	504	MPD	O2-C2	3.40	1.53	1.44
4	A	507	EDO	O2-C2	2.39	1.54	1.42

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	504	MPD	CM-C2-C1	-3.65	102.98	110.57
7	C	507	PEG	O2-C2-C1	3.45	125.21	110.07
6	B	504	MPD	O4-C4-C3	2.84	122.83	111.36
3	A	503	GOL	O1-C1-C2	2.40	121.69	110.20
7	D	505	PEG	O2-C3-C4	2.36	120.45	110.07

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	GOL	O1-C1-C2-C3
3	A	502	GOL	C1-C2-C3-O3
3	A	503	GOL	O1-C1-C2-C3
3	C	501	GOL	O1-C1-C2-O2
3	C	501	GOL	O1-C1-C2-C3

There are no ring outliers.

25 monomers are involved in 99 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	501	GOL	5	0
4	A	512	EDO	2	0
4	A	509	EDO	2	0
4	C	505	EDO	1	0
4	A	504	EDO	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	507	PEG	4	0
3	C	502	GOL	8	0
7	C	506	PEG	7	0
8	C	513	ACT	1	0
4	B	507	EDO	10	0
4	B	502	EDO	6	0
6	B	504	MPD	2	0
3	D	501	GOL	3	0
4	A	511	EDO	1	0
4	B	503	EDO	3	0
3	A	502	GOL	3	0
2	A	501	PO4	1	0
3	A	503	GOL	1	0
4	A	505	EDO	3	0
4	A	507	EDO	22	0
4	D	503	EDO	2	0
7	D	505	PEG	2	0
3	B	501	GOL	2	0
4	A	510	EDO	5	0
4	C	504	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	322/322 (100%)	0.30	27 (8%) 11 8	17, 26, 61, 86	13 (4%)
1	B	322/322 (100%)	0.44	30 (9%) 8 6	16, 26, 66, 80	17 (5%)
1	C	322/322 (100%)	0.55	28 (8%) 10 8	17, 28, 70, 86	16 (4%)
1	D	322/322 (100%)	0.25	26 (8%) 12 9	17, 28, 57, 85	8 (2%)
All	All	1288/1288 (100%)	0.39	111 (8%) 10 8	16, 27, 66, 86	54 (4%)

The worst 5 of 111 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	211	GLY	18.6
1	B	209	ALA	17.3
1	C	214	GLY	17.2
1	C	209	ALA	16.6
1	C	216	ALA	15.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	MPD	B	504	8/8	0.70	0.23	28,43,49,50	0
4	EDO	A	517	4/4	0.71	0.16	47,51,57,59	0
8	ACT	C	513	4/4	0.79	0.21	39,46,51,53	0
7	PEG	C	507	7/7	0.80	0.36	35,36,49,51	0
4	EDO	A	506	4/4	0.81	0.17	39,40,49,59	0
4	EDO	D	503	4/4	0.81	0.53	46,50,51,52	0
5	NH4	C	510	1/1	0.81	0.38	31,31,31,31	0
4	EDO	C	503	4/4	0.82	0.17	48,55,56,57	0
3	GOL	D	504	6/6	0.83	0.38	52,57,60,62	0
3	GOL	A	502	6/6	0.85	0.16	40,56,59,69	0
7	PEG	D	505	7/7	0.86	0.39	33,40,49,51	0
3	GOL	C	501	6/6	0.86	0.31	37,45,48,50	0
4	EDO	A	510	4/4	0.88	0.18	40,41,46,48	0
3	GOL	C	502	6/6	0.88	0.39	37,40,55,61	0
4	EDO	A	509	4/4	0.88	0.34	38,39,40,51	0
4	EDO	C	505	4/4	0.89	0.28	46,47,52,58	0
7	PEG	C	506	7/7	0.89	0.20	35,47,53,54	0
3	GOL	A	503	6/6	0.89	0.45	44,55,58,63	0
3	GOL	D	501	6/6	0.89	0.22	43,56,56,59	0
5	NH4	C	511	1/1	0.89	0.47	34,34,34,34	0
4	EDO	A	508	4/4	0.90	0.12	34,48,51,62	0
4	EDO	B	503	4/4	0.90	0.20	32,40,43,57	0
5	NH4	D	507	1/1	0.91	0.13	32,32,32,32	0
5	NH4	A	513	1/1	0.91	0.24	39,39,39,39	0
5	NH4	B	505	1/1	0.92	0.09	23,23,23,23	0
4	EDO	B	502	4/4	0.92	0.33	32,34,36,49	0
4	EDO	A	504	4/4	0.92	0.15	31,33,44,53	0
5	NH4	C	512	1/1	0.92	0.22	15,15,15,15	0
3	GOL	B	501	6/6	0.92	0.11	35,43,53,57	0
4	EDO	C	504	4/4	0.93	0.31	38,38,40,41	0
4	EDO	A	512	4/4	0.94	0.21	28,38,41,48	0
5	NH4	A	515	1/1	0.94	0.17	31,31,31,31	0
4	EDO	A	505	4/4	0.94	0.29	28,38,39,47	0
2	PO4	A	501	5/5	0.95	0.15	57,57,61,67	0
4	EDO	A	511	4/4	0.95	0.20	29,29,43,43	0
4	EDO	A	507	4/4	0.96	0.16	28,28,29,33	0
4	EDO	D	502	4/4	0.96	0.12	22,22,24,29	0
5	NH4	B	506	1/1	0.96	0.15	32,32,32,32	0
5	NH4	C	509	1/1	0.97	0.14	22,22,22,22	0
4	EDO	B	507	4/4	0.97	0.12	29,31,35,47	0
5	NH4	D	506	1/1	0.97	0.06	32,32,32,32	0
5	NH4	A	514	1/1	0.98	0.14	15,15,15,15	0
5	NH4	C	508	1/1	0.99	0.18	16,16,16,16	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NH4	A	516	1/1	0.99	0.16	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.