



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2025 – 12:10 PM JST

PDB ID : 9K2S / pdb_00009k2s
Title : Killer cell immunoglobulin-like receptor 2DL2 in complex with HLA-C*1202 and IY10 peptide
Authors : Chikata, T.; Kuroki, K.; Kuse, N.; Kliszczak, A.; Paes, W.; Tomioka, N.; Parker, R.; Aflalo, A.; Akahoshi, T.; Yamashita, R.; Sakata, R.; Kusaka, H.; Watanabe, Y.; Nicastrì, A.; Matsubara, H.; Ose, T.; Kita, S.; Oka, S.; Gatanaga, H.; Lin, Z.; Ternette, N.; Borrow, P.; Maenaka, K.; Takiguchi, M.
Deposited on : 2024-10-18
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engl & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	10	<div> <div>11%</div> <div>90%10%</div> </div>
2	A	276	<div> <div>6%</div> <div>80%18%.</div> </div>
3	B	119	<div> <div>6%</div> <div>65%18%17%</div> </div>
4	C	201	<div> <div>15%</div> <div>72%18%.9%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4711 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called peptide from p51 RT.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	D	10	Total	C	N	O	0	0	0
			82	55	13	14			

- Molecule 2 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	272	Total	C	N	O	S	0	0	0
			2237	1392	410	429	6			

- Molecule 3 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	VAL	LEU	conflict	UNP P61769

- Molecule 4 is a protein called KIR2DL2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	C	183	Total	C	N	O	S	0	0	0
			1419	896	244	271	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP Q8N742

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	2	Total 2	O 2	0	0
5	A	74	Total 74	O 74	0	0
5	B	27	Total 27	O 27	0	0
5	C	41	Total 41	O 41	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: peptide from p51 RT

Chain D:  90% 10%




- Molecule 2: MHC class I antigen

Chain A:  11% 80% 18%




- Molecule 3: Beta-2-microglobulin


Chain B:  6% 65% 18% 17%



- Molecule 4: KIR2DL2

Chain C:  15% 72% 18% 9%





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.49Å 44.59Å 88.24Å 90.00° 111.45° 90.00°	Depositor
Resolution (Å)	48.61 – 2.40 48.61 – 2.40	Depositor EDS
% Data completeness (in resolution range)	93.4 (48.61-2.40) 93.4 (48.61-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.245 , 0.299 0.244 , 0.299	Depositor DCC
R_{free} test set	1163 reflections (2.98%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.008 for l,-k,h	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4711	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.55	0/84	0.68	0/112
2	A	0.39	0/2300	0.65	0/3124
3	B	0.38	0/852	0.61	0/1152
4	C	0.48	0/1456	0.79	0/1971
All	All	0.42	0/4692	0.69	0/6359

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	82	0	87	3	0
2	A	2237	0	2076	35	0
3	B	829	0	791	13	0
4	C	1419	0	1356	22	0
5	A	74	0	0	3	0
5	B	27	0	0	2	0
5	C	41	0	0	1	0
5	D	2	0	0	0	0
All	All	4711	0	4310	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:117:THR:HG22	4:C:198:SER:HB2	1.66	0.77
2:A:34:PHE:HD2	2:A:53:VAL:HG11	1.57	0.69
2:A:193:HIS:NE2	3:B:99:ASP:OD2	2.23	0.69
2:A:109:ARG:HD3	5:A:342:HOH:O	1.95	0.67
4:C:102:THR:HG22	4:C:188:TRP:HB2	1.77	0.66
4:C:155:LYS:HD2	4:C:157:ASN:H	1.61	0.66
2:A:34:PHE:CD2	2:A:53:VAL:HG11	2.33	0.63
2:A:234:THR:OG1	2:A:244:LYS:HE2	2.00	0.62
4:C:17:LEU:HD12	4:C:100:VAL:HG22	1.83	0.61
2:A:56:GLU:OE1	2:A:171:ARG:NH2	2.29	0.60
2:A:259:THR:HG22	2:A:274:ARG:HG3	1.85	0.58
2:A:262:VAL:HB	2:A:271:LEU:HB2	1.84	0.58
4:C:40:HIS:NE2	4:C:42:GLU:OE2	2.27	0.57
2:A:36:ARG:HD3	2:A:49:ARG:HD3	1.87	0.57
2:A:155:GLU:HG3	2:A:158:ARG:HH22	1.70	0.57
4:C:155:LYS:HD3	4:C:156:VAL:HG23	1.87	0.57
3:B:96:TRP:CZ2	3:B:98:ARG:HG2	2.40	0.56
4:C:176:ARG:HD3	4:C:194:PRO:HA	1.89	0.54
2:A:185:HIS:HB3	2:A:266:GLY:O	2.08	0.53
2:A:146:ARG:HD2	4:C:133:SER:OG	2.09	0.53
3:B:26:CYS:HB2	3:B:40:LEU:HD21	1.92	0.52
2:A:251:PRO:HB2	2:A:254:GLU:HG3	1.93	0.50
2:A:15:ARG:HB3	2:A:18:ARG:HB2	1.93	0.50
2:A:160:TYR:CD1	2:A:164:THR:HG23	2.47	0.50
2:A:53:VAL:C	2:A:55:GLN:H	2.20	0.50
2:A:274:ARG:NH2	5:A:307:HOH:O	2.35	0.50
1:D:2:LEU:HD23	2:A:67:LYS:HB3	1.94	0.49
3:B:37:GLU:HB3	3:B:84:ASN:HB3	1.95	0.48
3:B:25:ASN:HB3	3:B:66:LEU:HD11	1.94	0.48
4:C:38:LEU:O	4:C:79:CYS:HA	2.13	0.47
3:B:95:LYS:HD2	5:B:213:HOH:O	2.14	0.47
4:C:124:VAL:HG23	4:C:166:LEU:HB2	1.95	0.47
4:C:138:HIS:O	4:C:177:CYS:HA	2.15	0.47
2:A:6:MET:HB2	2:A:169:LEU:HD13	1.96	0.46
2:A:168:TRP:CE3	2:A:171:ARG:HD3	2.50	0.46
2:A:4:HIS:HA	2:A:30:ASP:OD1	2.16	0.46
2:A:134:TRP:CH2	2:A:154:ALA:HB2	2.51	0.46
4:C:16:ARG:HD2	4:C:98:ASP:O	2.16	0.46
4:C:17:LEU:HD12	4:C:100:VAL:CG2	2.46	0.46
2:A:15:ARG:NH1	2:A:22:ARG:HB2	2.31	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:67:GLY:O	4:C:68:PRO:C	2.57	0.45
3:B:32:HIS:ND1	5:B:201:HOH:O	2.27	0.45
2:A:214:ILE:HD12	2:A:264:HIS:HB2	1.98	0.45
1:D:2:LEU:HD23	2:A:67:LYS:CB	2.47	0.45
2:A:155:GLU:O	2:A:156:GLN:C	2.60	0.45
2:A:104:LEU:HG	2:A:169:LEU:HD23	1.98	0.44
2:A:112:ARG:HE	2:A:114:TYR:HD2	1.65	0.44
4:C:106:GLU:OE2	4:C:132:SER:OG	2.29	0.44
2:A:188:THR:HA	2:A:205:TRP:O	2.16	0.44
3:B:8:ILE:HD12	3:B:92:LYS:HD3	1.99	0.44
4:C:10:LEU:HD11	4:C:26:LEU:HB3	1.98	0.44
4:C:23:THR:HG23	4:C:66:ILE:O	2.19	0.43
4:C:140:SER:HB3	4:C:145:ALA:HB2	2.00	0.43
3:B:3:GLN:HG2	3:B:33:PRO:HD3	2.01	0.43
4:C:29:TRP:HA	4:C:60:SER:O	2.19	0.43
3:B:7:LYS:HE3	3:B:7:LYS:HB2	1.46	0.42
2:A:110:LEU:HD22	2:A:162:GLU:HA	2.01	0.42
4:C:78:ARG:HG2	4:C:94:SER:O	2.19	0.42
2:A:256:GLN:H	2:A:256:GLN:CD	2.27	0.42
3:B:52:HIS:HB3	3:B:67:TYR:CD2	2.54	0.42
4:C:105:TYR:OH	5:C:301:HOH:O	2.21	0.42
1:D:2:LEU:HB3	2:A:67:LYS:HD2	2.01	0.42
2:A:160:TYR:CD2	2:A:161:LEU:HD23	2.55	0.42
3:B:5:THR:HG22	3:B:6:PRO:HD2	2.01	0.41
4:C:90:LEU:HD12	4:C:90:LEU:HA	1.64	0.41
2:A:220:ARG:HD3	2:A:257:ARG:CZ	2.51	0.41
2:A:244:LYS:HE3	5:A:309:HOH:O	2.21	0.41
3:B:56:SER:HB3	3:B:64:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	8/10 (80%)	5 (62%)	3 (38%)	0	100	100
2	A	268/276 (97%)	256 (96%)	12 (4%)	0	100	100
3	B	97/119 (82%)	97 (100%)	0	0	100	100
4	C	175/201 (87%)	172 (98%)	3 (2%)	0	100	100
All	All	548/606 (90%)	530 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	9/9 (100%)	9 (100%)	0	100	100
2	A	229/231 (99%)	228 (100%)	1 (0%)	89	95
3	B	94/109 (86%)	94 (100%)	0	100	100
4	C	159/173 (92%)	156 (98%)	3 (2%)	52	72
All	All	491/522 (94%)	487 (99%)	4 (1%)	79	90

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	263	GLN
4	C	149	ARG
4	C	155	LYS
4	C	156	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	71	GLN
2	A	181	GLN
4	C	113	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	10/10 (100%)	0.47	0 100 100	23, 33, 39, 41	0
2	A	272/276 (98%)	0.81	31 (11%) 11 10	14, 28, 55, 84	0
3	B	99/119 (83%)	0.67	7 (7%) 23 21	15, 29, 51, 71	0
4	C	183/201 (91%)	1.17	30 (16%) 5 5	18, 32, 60, 87	0
All	All	564/606 (93%)	0.90	68 (12%) 10 8	14, 30, 56, 87	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	C	56	HIS	6.9
4	C	155	LYS	5.2
4	C	94	SER	5.0
2	A	43	SER	4.9
4	C	171	HIS	4.8
4	C	156	VAL	4.8
3	B	2	ILE	4.6
4	C	84	THR	4.2
4	C	67	GLY	4.1
2	A	155	GLU	4.1
2	A	226	THR	4.0
4	C	83	VAL	3.9
2	A	129	GLU	3.9
3	B	48	GLU	3.8
3	B	75	GLU	3.7
2	A	228	ASP	3.7
2	A	182	ARG	3.7
2	A	18	ARG	3.6
4	C	121	GLY	3.6
4	C	50	HIS	3.6
4	C	200	THR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	C	52	ILE	3.5
4	C	57	ASP	3.4
4	C	157	ASN	3.4
4	C	21	GLU	3.3
4	C	31	ASP	3.1
2	A	17	GLY	3.1
4	C	35	GLU	3.1
2	A	45	ARG	3.0
2	A	178	GLU	3.0
2	A	221	ASP	3.0
4	C	116	PRO	2.9
4	C	131	ARG	2.9
2	A	164	THR	2.9
2	A	196	SER	2.9
2	A	266	GLY	2.9
4	C	141	ARG	2.9
2	A	137	ALA	2.8
3	B	90	GLN	2.8
3	B	59	LYS	2.8
4	C	118	VAL	2.8
2	A	222	GLY	2.7
4	C	144	GLU	2.7
4	C	115	GLY	2.7
2	A	227	GLN	2.7
2	A	265	GLU	2.6
2	A	112	ARG	2.6
2	A	54	GLU	2.5
2	A	198	HIS	2.4
4	C	55	HIS	2.4
2	A	59	GLU	2.4
3	B	49	LYS	2.4
2	A	217	THR	2.4
2	A	109	ARG	2.4
4	C	46	LYS	2.3
4	C	119	LEU	2.3
2	A	220	ARG	2.3
4	C	145	ALA	2.2
2	A	152	ARG	2.2
2	A	177	LYS	2.2
3	B	58	SER	2.2
2	A	252	SER	2.1
4	C	124	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	C	37	PHE	2.1
2	A	40	ASP	2.1
2	A	90	GLU	2.1
4	C	113	GLN	2.1
2	A	73	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.