



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 14, 2025 – 12:34 PM JST

PDB ID : 9KDS / pdb_00009kds
Title : The crystal structure of human AURKA kinase domain in complex with RA1
Authors : Zhang, Z.M.; Wang, L.
Deposited on : 2024-11-04
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	<div> <div>6%</div> <div>82%</div> <div>14%</div> <div>.</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aurora kinase A.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	257	Total	C	Cl	F	N	O	S	0	0	0
			2016	1308	1	1	340	362	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	290	ALA	CYS	engineered mutation	UNP O14965


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	23	Total	O	0	0
			23	23		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aurora kinase A



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	81.94Å 81.94Å 172.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.74 – 2.50 44.74 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.9 (44.74-2.50) 96.9 (44.74-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.9_1692+SVN	Depositor
R, R_{free}	0.213 , 0.253 0.232 , 0.270	Depositor DCC
R_{free} test set	1249 reflections (9.71%)	wwPDB-VP
Wilson B-factor (Å ²)	52.3	Xtriage
Anisotropy	0.320	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 64.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2039	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EFT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/2016	0.74	0/2742

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2016	0	1866	21	0
2	A	23	0	0	1	0
All	All	2039	0	1866	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:ARG:O	2:A:501:HOH:O	1.92	0.87
1:A:185:GLN:OE1	1:A:277:TRP:NE1	2.29	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:ALA:HB1	1:A:204:THR:HA	1.83	0.60
1:A:202:ASP:HB3	1:A:203:ALA:C	2.26	0.59
1:A:222:LEU:HD21	1:A:321:GLU:HG2	1.90	0.53
1:A:317:VAL:HG13	1:A:328:PRO:HD2	1.93	0.51
1:A:333:THR:HG22	1:A:335:GLN:H	1.75	0.50
1:A:254:HIS:NE2	1:A:274:ASP:OD1	2.45	0.50
1:A:340:ARG:NH1	1:A:345:GLU:O	2.43	0.49
1:A:335:GLN:OE1	1:A:335:GLN:N	2.46	0.49
1:A:256:ASP:HB3	1:A:296:LEU:HD23	1.94	0.48
1:A:139:LEU:HD13	1:A:220:A1EFT:C9	2.45	0.47
1:A:202:ASP:N	1:A:205:ARG:O	2.48	0.46
1:A:333:THR:HG22	1:A:335:GLN:N	2.30	0.45
1:A:327:PRO:HG2	1:A:330:GLU:HB2	1.98	0.45
1:A:296:LEU:HA	1:A:297:PRO:HD3	1.87	0.45
1:A:190:HIS:CG	1:A:191:PRO:HD2	2.54	0.43
1:A:246:TYR:O	1:A:249:SER:OG	2.28	0.42
1:A:213:ALA:HA	1:A:214:PRO:HD3	1.87	0.41
1:A:182:VAL:HG13	1:A:196:LEU:HD21	2.03	0.40
1:A:215:LEU:HB2	1:A:264:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	252/266 (95%)	243 (96%)	8 (3%)	1 (0%)	30 49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	276	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	195/233 (84%)	191 (98%)	4 (2%)	48 74

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	206	VAL
1	A	226	SER
1	A	260	GLU
1	A	292	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	A1EFT	A	220	1	49,53,54	3.73	13 (26%)	53,75,77	2.79	17 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	A1EFT	A	220	1	-	9/24/50/52	0/5/6/6

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	A1EFT	C7-C8	-14.94	1.33	1.53
1	A	220	A1EFT	O2-C8	10.92	1.41	1.23
1	A	220	A1EFT	C16-N4	10.91	1.42	1.28
1	A	220	A1EFT	C7-NH2	-10.64	1.34	1.46
1	A	220	A1EFT	C15-C16	3.96	1.55	1.49
1	A	220	A1EFT	C12-C11	3.53	1.45	1.37
1	A	220	A1EFT	CZ-NH1	-3.08	1.28	1.33
1	A	220	A1EFT	C9-N1	3.08	1.42	1.36
1	A	220	A1EFT	C14-C10	-2.94	1.45	1.49
1	A	220	A1EFT	O1-C5	2.75	1.41	1.37
1	A	220	A1EFT	O3-C27	2.71	1.41	1.37
1	A	220	A1EFT	C22-C16	2.67	1.53	1.49
1	A	220	A1EFT	C10-N2	2.66	1.38	1.34

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	220	A1EFT	C17-N4-C16	-13.55	102.11	117.48
1	A	220	A1EFT	O1-C5-C6	6.27	121.82	115.83
1	A	220	A1EFT	N3-C9-N2	-5.22	121.61	126.55
1	A	220	A1EFT	C11-C10-N2	-4.65	117.00	122.67
1	A	220	A1EFT	C15-C16-N4	-4.45	117.65	125.08
1	A	220	A1EFT	C23-C22-C16	3.65	124.98	122.04
1	A	220	A1EFT	O1-C5-C4	-3.34	118.37	124.12
1	A	220	A1EFT	C12-N3-C9	3.03	120.44	115.88
1	A	220	A1EFT	C13-O1-C5	-2.68	113.49	117.53
1	A	220	A1EFT	C21-C15-C14	2.59	122.52	119.96
1	A	220	A1EFT	C27-C22-C16	-2.55	115.44	119.83
1	A	220	A1EFT	C11-C12-N3	-2.49	119.67	123.82
1	A	220	A1EFT	NE-CZ-NH2	2.46	130.29	122.17
1	A	220	A1EFT	C15-C14-C10	2.30	128.58	122.33
1	A	220	A1EFT	C18-C14-C10	-2.27	112.82	117.75
1	A	220	A1EFT	C15-C16-C22	-2.25	115.48	118.10
1	A	220	A1EFT	C24-C23-C22	-2.11	119.81	123.58

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	220	A1EFT	N3-C9-N1-C3
1	A	220	A1EFT	N2-C9-N1-C3
1	A	220	A1EFT	C5-C6-C7-NH2
1	A	220	A1EFT	N4-C16-C22-C23
1	A	220	A1EFT	C4-C5-O1-C13
1	A	220	A1EFT	C6-C5-O1-C13
1	A	220	A1EFT	C1-C6-C7-NH2
1	A	220	A1EFT	C26-C27-O3-C28
1	A	220	A1EFT	C22-C27-O3-C28

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	220	A1EFT	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/266 (96%)	0.24	17 (6%) 26 24	37, 57, 99, 117	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	280	HIS	3.5
1	A	203	ALA	3.3
1	A	291	GLY	3.2
1	A	165	PHE	3.2
1	A	204	THR	3.1
1	A	128	TRP	3.1
1	A	281	ALA	2.8
1	A	305	MET	2.7
1	A	173	GLY	2.5
1	A	144	PHE	2.4
1	A	127	GLN	2.3
1	A	253	ILE	2.3
1	A	172	ALA	2.3
1	A	391	SER	2.3
1	A	176	HIS	2.2
1	A	390	PRO	2.1
1	A	277	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	A1EFT	A	220	48/49	0.84	0.20	46,82,112,142	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.