



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 29, 2024 – 09:07 PM EDT

PDB ID : 4KKU
Title : Structure of BesA (Selenomethionone derivative - P212121)
Authors : Greene, N.P.; Hinchliffe, P.; Crow, A.; Ababou, A.; Hughes, C.; Koronakis, V.
Deposited on : 2013-05-06
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1460 (2.36-2.36)
Clashscore	180529	1571 (2.36-2.36)
Ramachandran outliers	177936	1559 (2.36-2.36)
Sidechain outliers	177891	1559 (2.36-2.36)
RSRZ outliers	164620	1460 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	 10% 78% 9% • 10%
1	B	296	 6% 77% 11% • 10%
1	C	296	 13% 79% 8% • 11%
1	D	296	 19% 78% 9% • 10%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 8454 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane fusion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	266	2075	1331	340	399	5	0	3	0
1	B	267	2084	1339	342	398	5	0	3	0
1	C	264	2043	1311	337	391	4	0	1	0
1	D	266	2074	1330	340	400	4	0	3	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLY	-	expression tag	UNP O51166
A	23	SER	-	expression tag	UNP O51166
A	24	HIS	-	expression tag	UNP O51166
A	25	MSE	-	expression tag	UNP O51166
B	22	GLY	-	expression tag	UNP O51166
B	23	SER	-	expression tag	UNP O51166
B	24	HIS	-	expression tag	UNP O51166
B	25	MSE	-	expression tag	UNP O51166
C	22	GLY	-	expression tag	UNP O51166
C	23	SER	-	expression tag	UNP O51166
C	24	HIS	-	expression tag	UNP O51166
C	25	MSE	-	expression tag	UNP O51166
D	22	GLY	-	expression tag	UNP O51166
D	23	SER	-	expression tag	UNP O51166
D	24	HIS	-	expression tag	UNP O51166
D	25	MSE	-	expression tag	UNP O51166


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	50	Total 50	O 50	0	0
2	B	47	Total 47	O 47	0	0
2	C	37	Total 37	O 37	0	0
2	D	44	Total 44	O 44	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Membrane fusion protein



- Molecule 1: Membrane fusion protein




- Molecule 1: Membrane fusion protein



- Molecule 1: Membrane fusion protein

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.74Å 152.13Å 155.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	77.74 – 2.35 77.74 – 2.35	Depositor EDS
% Data completeness (in resolution range)	98.5 (77.74-2.35) 98.5 (77.74-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.236 , 0.264 0.240 , 0.266	Depositor DCC
R_{free} test set	3697 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtrriage
Anisotropy	0.049	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 23.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.000 for -h,l,k	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8454	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.5035e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.89	0/2094	0.93	3/2814 (0.1%)
1	B	0.92	0/2103	0.92	4/2825 (0.1%)
1	C	0.89	2/2060 (0.1%)	0.94	6/2766 (0.2%)
1	D	0.84	1/2093 (0.0%)	0.98	8/2813 (0.3%)
All	All	0.88	3/8350 (0.0%)	0.94	21/11218 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	271	GLU	CD-OE1	7.57	1.33	1.25
1	C	281	GLU	CG-CD	7.45	1.63	1.51
1	D	271	GLU	CD-OE1	6.02	1.32	1.25

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	220	MSE	CA-CB-CG	13.70	136.59	113.30
1	D	220	MSE	CG-SE-CE	-9.66	77.64	98.90
1	D	220	MSE	N-CA-CB	-9.15	94.13	110.60
1	A	146	VAL	CB-CA-C	-7.55	97.05	111.40
1	B	146	VAL	CB-CA-C	-7.46	97.23	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2075	0	2199	34	0
1	B	2084	0	2219	30	0
1	C	2043	0	2180	23	0
1	D	2074	0	2197	25	0
2	A	50	0	0	7	0
2	B	47	0	0	4	0
2	C	37	0	0	5	0
2	D	44	0	0	5	0
All	All	8454	0	8795	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:230:ARG:HG3	2:D:420:HOH:O	1.49	1.11
1:D:245:ARG:NH2	1:D:263:MSE:HE1	1.91	0.86
1:A:142:ASN:ND2	2:A:412:HOH:O	2.09	0.84
1:B:245:ARG:NH2	1:B:263:MSE:HE1	1.93	0.83
1:D:69:ASP:OD2	2:D:441:HOH:O	1.99	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/296 (90%)	256 (97%)	9 (3%)	0	100	100
1	B	266/296 (90%)	257 (97%)	9 (3%)	0	100	100
1	C	261/296 (88%)	254 (97%)	7 (3%)	0	100	100
1	D	265/296 (90%)	257 (97%)	8 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1057/1184 (89%)	1024 (97%)	33 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/257 (93%)	223 (94%)	15 (6%)	15	16
1	B	239/257 (93%)	228 (95%)	11 (5%)	23	28
1	C	234/257 (91%)	224 (96%)	10 (4%)	25	31
1	D	238/257 (93%)	226 (95%)	12 (5%)	20	25
All	All	949/1028 (92%)	901 (95%)	48 (5%)	20	24

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	142	ASN
1	C	297	SER
1	C	181	ASN
1	C	220	MSE
1	D	91	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	181	ASN
1	C	274	ASN
1	D	153	GLN
1	B	96	GLN
1	A	181	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/296 (88%)	0.50	30 (11%) 11 13	11, 35, 72, 115	2 (0%)
1	B	263/296 (88%)	0.40	19 (7%) 23 27	12, 33, 68, 98	2 (0%)
1	C	260/296 (87%)	0.62	38 (14%) 7 8	18, 37, 96, 112	1 (0%)
1	D	262/296 (88%)	0.97	56 (21%) 3 3	18, 39, 107, 130	3 (1%)
All	All	1047/1184 (88%)	0.62	143 (13%) 8 9	11, 36, 90, 130	8 (0%)

The worst 5 of 143 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	114	LEU	8.1
1	D	106	PRO	8.0
1	A	106	PRO	7.0
1	D	91	LEU	6.8
1	D	81	VAL	6.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.