



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 10, 2025 – 02:40 PM JST

PDB ID : 9KK9 / pdb\_00009kk9  
Title : Structure of the transaminase PhnW from *Vibrio vulnificus* in complex with PLP and AEP  
Authors : Ma, Q.; Han, M.  
Deposited on : 2024-11-13  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 11%, green 86%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>86%</span> <span>11% ..</span> </div> </div>
1	B	369	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 8%, green 89%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>89%</span> <span>8% ..</span> </div> </div>
1	C	369	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 8%, green 89%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>89%</span> <span>8% ..</span> </div> </div>




In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-aminoethylphosphonate--pyruvate transaminase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	362	Total 2812	C 1786	N 472	O 538	P 1	S 15	0	0	0
1	B	362	Total 2812	C 1786	N 472	O 538	P 1	S 15	0	0	0
1	C	362	Total 2812	C 1786	N 472	O 538	P 1	S 15	0	0	0

- Molecule 2 is (2-aminoethyl)phosphonic acid (CCD ID: P7I) (formula:  $\text{C}_2\text{H}_8\text{NO}_3\text{P}$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 7	C 2	N 1	O 3	P 1	0	0
2	B	1	Total 7	C 2	N 1	O 3	P 1	0	0
2	C	1	Total 7	C 2	N 1	O 3	P 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	102	Total 102	O 102	0	0
3	B	100	Total 100	O 100	0	0
3	C	113	Total 113	O 113	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-aminoethylphosphonate--pyruvate transaminase



- Molecule 1: 2-aminoethylphosphonate--pyruvate transaminase



- Molecule 1: 2-aminoethylphosphonate--pyruvate transaminase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	158.25Å 94.94Å 91.18Å 90.00° 90.51° 90.00°	Depositor
Resolution (Å)	60.87 – 2.15 60.87 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.4 (60.87-2.15) 99.4 (60.87-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.14Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.167 , 0.189 0.169 , 0.199	Depositor DCC
$R_{free}$ test set	3534 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.4	Xtriage
Anisotropy	0.634	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 35.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.006 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.000 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.055 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.038 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.018 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8772	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: P7I, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.82	1/2850 (0.0%)	1.20	8/3867 (0.2%)
1	B	0.82	1/2850 (0.0%)	1.18	4/3867 (0.1%)
1	C	0.85	1/2850 (0.0%)	1.19	3/3867 (0.1%)
All	All	0.83	3/8550 (0.0%)	1.19	15/11601 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	23	MET	SD-CE	-7.29	1.61	1.79
1	A	23	MET	SD-CE	-6.91	1.62	1.79
1	B	23	MET	SD-CE	-6.49	1.63	1.79

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	109	VAL	N-CA-CB	6.26	119.23	111.41
1	A	109	VAL	N-CA-C	-5.75	98.14	106.88
1	C	109	VAL	N-CA-C	-5.41	98.66	106.88
1	A	259	GLU	CA-C-N	5.33	127.96	120.28
1	A	259	GLU	C-N-CA	5.33	127.96	120.28
1	C	87	ASP	CA-CB-CG	5.19	117.79	112.60
1	B	141	GLU	CA-C-N	5.17	127.93	120.38
1	B	141	GLU	C-N-CA	5.17	127.93	120.38
1	A	87	ASP	CA-CB-CG	5.16	117.76	112.60
1	C	297	SER	N-CA-C	-5.10	103.52	110.36
1	B	112	GLY	N-CA-C	-5.04	106.23	112.33
1	A	122	VAL	CA-C-N	5.03	127.02	120.28
1	A	122	VAL	C-N-CA	5.03	127.02	120.28
1	A	141	GLU	CA-C-N	5.03	127.72	120.38
1	A	141	GLU	C-N-CA	5.03	127.72	120.38

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2812	0	2757	24	0
1	B	2812	0	2757	13	0
1	C	2812	0	2757	17	0
2	A	7	0	4	0	0
2	B	7	0	4	0	0
2	C	7	0	4	0	0
3	A	102	0	0	1	0
3	B	100	0	0	0	0
3	C	113	0	0	1	0
All	All	8772	0	8283	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:VAL:HG12	1:A:153:PHE:HZ	1.55	0.70
1:A:135:VAL:HG12	1:A:153:PHE:CZ	2.31	0.66
1:A:20:ARG:HA	1:A:23:MET:HE2	1.79	0.64
1:C:268:HIS:HD2	3:C:513:HOH:O	1.86	0.58
1:C:117:PRO:HD2	1:C:149:PRO:HD2	1.89	0.55
1:C:312:GLN:HE21	1:C:314:LYS:HB3	1.71	0.55
1:A:312:GLN:HE21	1:A:314:LYS:HB3	1.72	0.55
1:A:354:LEU:O	1:A:358:ILE:HG12	2.07	0.54
1:C:11:GLY:HA2	1:C:12:PRO:C	2.33	0.54
1:A:135:VAL:CG1	1:A:153:PHE:CZ	2.92	0.53
1:A:11:GLY:HA2	1:A:12:PRO:C	2.35	0.52
1:A:117:PRO:HD2	1:A:149:PRO:HD2	1.90	0.52
1:A:268:HIS:HD2	3:A:512:HOH:O	1.92	0.52
1:B:31:ASP:OD2	1:B:33:GLU:HB2	2.11	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:11:GLY:HA2	1:B:12:PRO:C	2.36	0.50
1:B:116:GLN:HG2	1:B:149:PRO:HD3	1.94	0.50
1:B:179:ILE:HB	1:B:184:ILE:HB	1.94	0.49
1:A:179:ILE:HB	1:A:184:ILE:HB	1.94	0.49
1:C:179:ILE:HB	1:C:184:ILE:HB	1.95	0.48
1:A:239:TRP:CD2	1:A:243:SER:HB3	2.48	0.48
1:B:239:TRP:CD2	1:B:243:SER:HB3	2.49	0.48
1:B:40:GLU:OE1	1:B:43:ARG:HD3	2.14	0.47
1:A:37:ASP:CG	1:C:112:GLY:HA3	2.39	0.47
1:B:84:LEU:HB3	1:B:135:VAL:HG12	1.97	0.47
1:C:40:GLU:OE1	1:C:43:ARG:HD3	2.15	0.47
1:C:135:VAL:HG21	1:C:157:ALA:HB2	1.97	0.47
1:C:239:TRP:CD2	1:C:243:SER:HB3	2.50	0.46
1:A:88:ASN:HB3	1:A:111:PRO:HD2	1.98	0.46
1:C:158:LYS:HA	1:C:158:LYS:HD2	1.70	0.45
1:C:195:ILE:HG21	1:C:251:PHE:CE1	2.51	0.45
1:A:230:HIS:O	1:A:234:VAL:HG22	2.17	0.45
1:A:82:LYS:HG3	1:A:105:PRO:HB2	1.97	0.45
1:A:84:LEU:HB3	1:A:135:VAL:HG13	1.99	0.45
1:B:167:ASP:OD1	1:B:169:MET:HB2	2.16	0.44
1:B:282:MET:SD	1:B:358:ILE:HD13	2.57	0.44
1:C:59:VAL:HG22	1:C:229:TRP:HB2	1.99	0.44
1:A:44:THR:HB	1:C:334:ASN:HD22	1.83	0.43
1:B:86:VAL:HG11	1:B:117:PRO:HG3	1.99	0.43
1:A:139:HIS:HB2	1:A:166:LEU:HD11	2.00	0.43
1:A:158:LYS:HA	1:A:158:LYS:HD2	1.90	0.43
1:C:135:VAL:HG21	1:C:157:ALA:CB	2.48	0.43
1:A:275:GLN:HE21	1:A:275:GLN:HB3	1.69	0.42
1:B:358:ILE:O	1:B:362:MET:HG3	2.20	0.42
1:C:84:LEU:HB3	1:C:135:VAL:HG12	2.03	0.41
1:C:86:VAL:HG11	1:C:117:PRO:HG3	2.03	0.41
1:A:167:ASP:OD1	1:A:169:MET:HB2	2.21	0.41
1:B:139:HIS:HB2	1:B:166:LEU:HD11	2.01	0.41
1:B:135:VAL:HG21	1:B:157:ALA:HB2	2.02	0.40
1:A:158:LYS:HD3	1:A:164:VAL:HG23	2.02	0.40
1:A:33:GLU:OE1	1:C:118:HIS:CE1	2.75	0.40
1:A:235:ASN:HB2	1:A:238:LYS:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/369 (97%)	348 (97%)	11 (3%)	0	100	100
1	B	359/369 (97%)	351 (98%)	8 (2%)	0	100	100
1	C	359/369 (97%)	350 (98%)	9 (2%)	0	100	100
All	All	1077/1107 (97%)	1049 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/302 (99%)	293 (98%)	5 (2%)	56	61
1	B	298/302 (99%)	292 (98%)	6 (2%)	50	55
1	C	298/302 (99%)	291 (98%)	7 (2%)	45	49
All	All	894/906 (99%)	876 (98%)	18 (2%)	50	55

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	GLU
1	A	109	VAL
1	A	135	VAL
1	A	141	GLU
1	A	352	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	21	GLU
1	B	59	VAL
1	B	109	VAL
1	B	141	GLU
1	B	290	LEU
1	B	342	ASN
1	C	84	LEU
1	C	109	VAL
1	C	110	SER
1	C	141	GLU
1	C	158	LYS
1	C	275	GLN
1	C	352	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	A	192	ASN
1	A	275	GLN
1	A	312	GLN
1	A	342	ASN
1	B	120	ASN
1	B	192	ASN
1	B	236	HIS
1	B	268	HIS
1	B	269	ASN
1	B	274	ASN
1	B	312	GLN
1	B	342	ASN
1	C	118	HIS
1	C	120	ASN
1	C	192	ASN
1	C	235	ASN
1	C	269	ASN
1	C	275	GLN
1	C	312	GLN
1	C	334	ASN
1	C	342	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	LLP	C	193	1,2	23,24,25	1.22	2 (8%)	25,32,34	1.38	2 (8%)
1	LLP	A	193	1,2	23,24,25	1.31	1 (4%)	25,32,34	1.46	4 (16%)
1	LLP	B	193	1,2	23,24,25	1.36	1 (4%)	25,32,34	1.35	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	C	193	1,2	-	4/16/17/19	0/1/1/1
1	LLP	A	193	1,2	-	4/16/17/19	0/1/1/1
1	LLP	B	193	1,2	-	6/16/17/19	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	193	LLP	C4'-NZ	5.25	1.44	1.27
1	B	193	LLP	C4'-NZ	5.09	1.44	1.27
1	C	193	LLP	C4'-NZ	4.50	1.42	1.27
1	C	193	LLP	OP4-C5'	-2.01	1.37	1.45

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	193	LLP	C4-C4'-NZ	-4.56	103.39	124.31
1	C	193	LLP	C4-C4'-NZ	-4.36	104.29	124.31
1	B	193	LLP	C4-C4'-NZ	-3.93	106.27	124.31
1	B	193	LLP	C3-C4-C5	-3.08	115.90	118.26
1	A	193	LLP	C3-C4-C5	-2.96	115.99	118.26
1	C	193	LLP	C3-C4-C5	-2.63	116.24	118.26
1	A	193	LLP	C4-C3-C2	2.20	121.55	120.19
1	B	193	LLP	C4-C3-C2	2.14	121.51	120.19
1	A	193	LLP	OP4-P-OP1	2.06	112.24	106.47
1	B	193	LLP	OP4-P-OP1	2.00	112.09	106.47

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	193	LLP	O-C-CA-CB
1	B	193	LLP	C6-C5-C5'-OP4
1	B	193	LLP	O-C-CA-CB
1	C	193	LLP	O-C-CA-CB
1	B	193	LLP	C4-C4'-NZ-CE
1	C	193	LLP	C4-C4'-NZ-CE
1	A	193	LLP	C4-C4'-NZ-CE
1	A	193	LLP	CG-CD-CE-NZ
1	B	193	LLP	CG-CD-CE-NZ
1	C	193	LLP	CG-CD-CE-NZ
1	A	193	LLP	C3-C4-C4'-NZ
1	B	193	LLP	C3-C4-C4'-NZ
1	C	193	LLP	C3-C4-C4'-NZ
1	B	193	LLP	C5'-OP4-P-OP3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	P7I	C	401	1	6,6,6	1.21	1 (16%)	8,8,8	1.83	2 (25%)
2	P7I	B	401	1	6,6,6	1.18	1 (16%)	8,8,8	1.88	2 (25%)
2	P7I	A	401	1	6,6,6	1.10	0	8,8,8	2.25	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P7I	C	401	1	-	1/4/4/4	-
2	P7I	B	401	1	-	1/4/4/4	-
2	P7I	A	401	1	-	1/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	P7I	PAG-CAF	2.31	1.81	1.78
2	B	401	P7I	PAG-CAF	2.28	1.81	1.78

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	P7I	PAG-CAF-CAE	-5.81	107.86	114.99
2	B	401	P7I	PAG-CAF-CAE	-4.64	109.30	114.99
2	C	401	P7I	PAG-CAF-CAE	-4.39	109.60	114.99
2	C	401	P7I	OAD-PAG-CAF	-2.34	107.06	111.40
2	B	401	P7I	OAD-PAG-CAF	-2.25	107.23	111.40
2	A	401	P7I	OAD-PAG-CAF	-2.22	107.29	111.40

There are no chirality outliers.


All (3) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	401	P7I	NAA-CAE-CAF-PAG
2	B	401	P7I	NAA-CAE-CAF-PAG
2	C	401	P7I	NAA-CAE-CAF-PAG

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	361/369 (97%)	-0.23	4 (1%) 77 81	35, 48, 75, 100	0
1	B	361/369 (97%)	-0.23	4 (1%) 77 81	35, 49, 74, 96	0
1	C	361/369 (97%)	-0.28	4 (1%) 77 81	33, 45, 69, 89	0
All	All	1083/1107 (97%)	-0.24	12 (1%) 77 81	33, 47, 74, 100	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	110	SER	4.4
1	C	111	PRO	4.0
1	C	110	SER	3.9
1	A	153	PHE	3.9
1	A	111	PRO	3.0
1	C	119	LEU	2.8
1	C	153	PHE	2.8
1	B	153	PHE	2.7
1	B	365	GLN	2.2
1	A	365	GLN	2.2
1	B	5	TYR	2.0
1	B	159	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	LLP	A	193	24/25	0.97	0.07	34,40,50,56	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	LLP	B	193	24/25	0.97	0.07	35,44,51,55	0
1	LLP	C	193	24/25	0.98	0.08	33,37,47,52	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	P7I	B	401	7/7	0.96	0.08	44,48,53,57	0
2	P7I	A	401	7/7	0.97	0.07	45,47,52,57	0
2	P7I	C	401	7/7	0.97	0.08	41,43,51,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around P7I B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around P7I A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around P7I C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.