



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 25, 2024 – 06:31 PM EST

PDB ID : 3LHK
Title : Crystal structure of putative DNA binding protein from *Methanocaldococcus jannaschii*.
Authors : Chang, C.; Chhor, G.; Cobb, G.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2010-01-22
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	154	 84% 10% • 5%
1	B	154	 86% 8% 6%
1	C	154	 10% 81% 12% • 6%
1	D	154	 5% 82% 10% • 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative DNA binding protein MJ0014.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	Se	0	0	0
			1202	776	208	215	3			
1	B	145	Total	C	N	O	Se	0	0	0
			1193	771	207	212	3			
1	C	145	Total	C	N	O	Se	0	0	0
			1176	761	202	210	3			
1	D	145	Total	C	N	O	Se	0	0	0
			1187	767	205	212	3			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	SER	-	expression tag	UNP Q60329
A	61	ASN	-	expression tag	UNP Q60329
A	62	ALA	-	expression tag	UNP Q60329
B	60	SER	-	expression tag	UNP Q60329
B	61	ASN	-	expression tag	UNP Q60329
B	62	ALA	-	expression tag	UNP Q60329
C	60	SER	-	expression tag	UNP Q60329
C	61	ASN	-	expression tag	UNP Q60329
C	62	ALA	-	expression tag	UNP Q60329
D	60	SER	-	expression tag	UNP Q60329
D	61	ASN	-	expression tag	UNP Q60329
D	62	ALA	-	expression tag	UNP Q60329

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	82	Total	O	0	0
			82	82		
2	B	68	Total	O	0	0
			68	68		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	71	Total	O	0	0
			71	71		
2	D	61	Total	O	0	0
			61	61		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: putative DNA binding protein MJ0014




- Molecule 1: putative DNA binding protein MJ0014



- Molecule 1: putative DNA binding protein MJ0014



- Molecule 1: putative DNA binding protein MJ0014



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.35Å 86.16Å 115.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 50.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.20) 99.8 (50.00-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.09 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.203 , 0.249 0.215 , 0.251	Depositor DCC
R_{free} test set	2000 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.690	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5040	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9137e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/1219	0.72	1/1625 (0.1%)
1	B	0.70	0/1210	0.69	0/1613
1	C	0.66	0/1192	0.66	0/1590
1	D	0.58	0/1204	0.66	0/1607
All	All	0.66	0/4825	0.68	1/6435 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	ARG	CG-CD-NE	-8.26	94.45	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1202	0	1252	12	0
1	B	1193	0	1246	8	0
1	C	1176	0	1219	15	0
1	D	1187	0	1228	19	0
2	A	82	0	0	1	0
2	B	68	0	0	2	0
2	C	71	0	0	3	0
2	D	61	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5040	0	4945	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:194:LYS:O	1:C:198:LYS:HG2	1.66	0.95
1:C:63:LYS:HE3	1:C:98:GLN:OE1	1.70	0.91
1:C:192:LYS:HA	2:C:218:HOH:O	1.84	0.78
1:D:78:ASP:OD2	2:D:278:HOH:O	2.03	0.76
1:A:117:LEU:HA	1:A:120:MSE:HE2	1.65	0.76
1:C:119:LYS:HA	1:C:122:MSE:HE3	1.68	0.74
1:C:63:LYS:NZ	2:C:227:HOH:O	2.32	0.62
1:C:127:GLU:HG2	1:C:128:LYS:HG3	1.81	0.62
1:A:160:ASN:HB3	1:A:163:HIS:HB2	1.83	0.61
1:D:87:LYS:HD3	2:D:279:HOH:O	2.01	0.61
1:D:65:ILE:HD13	1:D:120:MSE:CE	2.32	0.59
1:C:195:LYS:HB3	2:C:218:HOH:O	2.02	0.59
1:A:191:HIS:HB2	2:A:234:HOH:O	2.03	0.58
1:D:77:ASP:HB2	2:D:241:HOH:O	2.02	0.58
1:C:177:ILE:HG21	1:D:177:ILE:HG21	1.86	0.57
1:D:78:ASP:OD1	1:D:78:ASP:C	2.43	0.57
1:B:127:GLU:HG2	1:B:128:LYS:HG3	1.86	0.57
1:A:78:ASP:O	1:A:81:ARG:HB3	2.06	0.56
1:B:88:SER:O	1:B:92:GLU:HG3	2.09	0.53
1:A:177:ILE:HG21	1:B:177:ILE:HG21	1.90	0.52
1:D:65:ILE:HD13	1:D:120:MSE:HE1	1.91	0.52
1:A:153:GLY:O	1:D:124:ARG:NH1	2.43	0.51
1:D:171:VAL:O	1:D:175:ILE:HG12	2.09	0.51
1:D:117:LEU:HA	1:D:120:MSE:HE2	1.93	0.50
1:C:61:ASN:O	1:C:62:ALA:CB	2.59	0.50
1:C:72:PHE:HZ	1:C:172:GLU:HG3	1.77	0.49
1:A:107:LEU:HD21	1:A:177:ILE:HD11	1.96	0.47
1:C:61:ASN:O	1:C:62:ALA:HB2	2.14	0.47
1:D:78:ASP:OD1	1:D:78:ASP:O	2.33	0.46
1:D:65:ILE:HD13	1:D:120:MSE:HE3	1.96	0.46
1:D:182:ALA:HB2	1:D:196:LEU:HD23	1.97	0.46
1:D:78:ASP:O	1:D:81:ARG:HB3	2.16	0.46
1:B:110:LYS:HD2	1:B:115:LYS:HD2	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:LYS:HG3	2:B:258:HOH:O	2.16	0.45
2:B:243:HOH:O	1:D:128:LYS:HE3	2.16	0.45
1:C:79:LEU:HD21	1:C:101:LYS:HD2	1.99	0.45
1:A:65:ILE:HG21	1:A:117:LEU:HD13	1.99	0.45
1:D:64:ILE:HG12	1:D:128:LYS:HB2	2.00	0.44
1:A:201:LYS:HE3	1:A:201:LYS:HB2	1.60	0.44
1:C:148:PHE:CZ	1:D:145:LEU:HD12	2.52	0.44
1:C:72:PHE:CZ	1:C:172:GLU:HG3	2.54	0.43
1:D:188:MSE:HE2	1:D:188:MSE:HB3	1.96	0.42
1:A:69:ARG:HB2	1:A:102:ASP:O	2.18	0.42
1:C:68:ALA:O	1:C:101:LYS:HA	2.19	0.42
1:B:174:LEU:HD23	1:B:204:VAL:HG11	2.02	0.42
1:B:63:LYS:HA	1:B:63:LYS:HD2	1.87	0.42
1:A:68:ALA:O	1:A:101:LYS:HA	2.21	0.41
1:A:118:LEU:HA	1:A:118:LEU:HD23	1.86	0.41
1:B:97:ILE:HD13	1:B:97:ILE:HG21	1.80	0.41
1:D:195:LYS:HA	1:D:198:LYS:HE3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/154 (94%)	142 (99%)	2 (1%)	0	100	100
1	B	143/154 (93%)	143 (100%)	0	0	100	100
1	C	143/154 (93%)	141 (99%)	0	2 (1%)	9	7
1	D	143/154 (93%)	140 (98%)	2 (1%)	1 (1%)	19	19
All	All	573/616 (93%)	566 (99%)	4 (1%)	3 (0%)	25	28

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	62	ALA
1	C	204	VAL
1	D	77	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/136 (96%)	129 (99%)	1 (1%)	79	88
1	B	129/136 (95%)	129 (100%)	0	100	100
1	C	125/136 (92%)	123 (98%)	2 (2%)	58	73
1	D	127/136 (93%)	125 (98%)	2 (2%)	58	73
All	All	511/544 (94%)	506 (99%)	5 (1%)	73	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	ARG
1	C	98	GLN
1	C	201	LYS
1	D	78	ASP
1	D	120	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	84	GLN
1	C	167	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Warning: The R factor obtained from EDS is 0.2538, which does not match the depositor's R factor of 0.20347. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	143/154 (92%)	-0.12	2 (1%) 73 70	10, 17, 31, 43	0
1	B	142/154 (92%)	-0.15	0 100 100	9, 17, 27, 35	0
1	C	142/154 (92%)	0.37	15 (10%) 13 10	8, 18, 28, 47	0
1	D	142/154 (92%)	0.53	8 (5%) 31 28	12, 20, 30, 38	0
All	All	569/616 (92%)	0.15	25 (4%) 39 36	8, 18, 30, 47	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	61	ASN	6.8
1	C	62	ALA	5.7
1	C	204	VAL	4.8
1	A	61	ASN	4.7
1	C	166	PRO	4.3
1	C	165	THR	4.2
1	C	205	ARG	3.8
1	D	77	ASP	3.2
1	C	169	GLU	2.9
1	C	78	ASP	2.8
1	D	189	HIS	2.7
1	C	81	ARG	2.7
1	C	164	LYS	2.6
1	C	174	LEU	2.5
1	C	167	GLN	2.5
1	D	177	ILE	2.4
1	C	173	ASP	2.3
1	C	189	HIS	2.3
1	D	96	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	170	LEU	2.2
1	A	191	HIS	2.2
1	D	78	ASP	2.2
1	D	62	ALA	2.1
1	D	84	GLN	2.1
1	D	181	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.