



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 3, 2025 – 08:17 PM JST

PDB ID : 9LSC / pdb_00009lsc
Title : Crystal structure of mRFP1 with a grafted calcium-binding sequence and two bound calcium ions in a calcium-free solution
Authors : Uehara, R.; Kamiya, Y.; Maeda, S.; Okamoto, K.; Toya, S.; Chiba, R.; Ame-saka, H.; Takano, K.; Matsumura, H.; Tanaka, S.-i.
Deposited on : 2025-02-04
Resolution : 1.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3869 (1.40-1.36)
Clashscore	180529	4183 (1.40-1.36)
Ramachandran outliers	177936	4116 (1.40-1.36)
Sidechain outliers	177891	4115 (1.40-1.36)
RSRZ outliers	164620	3867 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	<div> <div>4%</div> <div> <div></div> <div>75%</div> <div>•</div> <div>21%</div> </div> </div>
1	B	278	<div> <div>4%</div> <div> <div></div> <div>74%</div> <div>5%</div> <div>21%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4014 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Red fluorescent protein,grafted calcium-binding sequence.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	0	0
			1767	1122	296	341	8			
1	B	220	Total	C	N	O	S	0	0	0
			1767	1122	296	341	8			

There are 106 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-36	MET	-	initiating methionine	UNP A0A4V4ND72
A	-35	GLY	-	expression tag	UNP A0A4V4ND72
A	-34	GLY	-	expression tag	UNP A0A4V4ND72
A	-33	SER	-	expression tag	UNP A0A4V4ND72
A	-32	HIS	-	expression tag	UNP A0A4V4ND72
A	-31	HIS	-	expression tag	UNP A0A4V4ND72
A	-30	HIS	-	expression tag	UNP A0A4V4ND72
A	-29	HIS	-	expression tag	UNP A0A4V4ND72
A	-28	HIS	-	expression tag	UNP A0A4V4ND72
A	-27	HIS	-	expression tag	UNP A0A4V4ND72
A	-26	GLY	-	expression tag	UNP A0A4V4ND72
A	-25	MET	-	expression tag	UNP A0A4V4ND72
A	-24	ALA	-	expression tag	UNP A0A4V4ND72
A	-23	SER	-	expression tag	UNP A0A4V4ND72
A	-22	MET	-	expression tag	UNP A0A4V4ND72
A	-21	THR	-	expression tag	UNP A0A4V4ND72
A	-20	GLY	-	expression tag	UNP A0A4V4ND72
A	-19	GLY	-	expression tag	UNP A0A4V4ND72
A	-18	GLN	-	expression tag	UNP A0A4V4ND72
A	-17	GLN	-	expression tag	UNP A0A4V4ND72
A	-16	MET	-	expression tag	UNP A0A4V4ND72
A	-15	GLY	-	expression tag	UNP A0A4V4ND72
A	-14	ARG	-	expression tag	UNP A0A4V4ND72
A	-13	ASP	-	expression tag	UNP A0A4V4ND72
A	-12	LEU	-	expression tag	UNP A0A4V4ND72

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	TYR	-	expression tag	UNP A0A4V4ND72
A	-10	ASP	-	expression tag	UNP A0A4V4ND72
A	-9	ASP	-	expression tag	UNP A0A4V4ND72
A	-8	ASP	-	expression tag	UNP A0A4V4ND72
A	-7	ASP	-	expression tag	UNP A0A4V4ND72
A	-6	LYS	-	expression tag	UNP A0A4V4ND72
A	-5	ASP	-	expression tag	UNP A0A4V4ND72
A	-4	ARG	-	expression tag	UNP A0A4V4ND72
A	-3	TRP	-	expression tag	UNP A0A4V4ND72
A	-2	GLY	-	expression tag	UNP A0A4V4ND72
A	-1	SER	-	expression tag	UNP A0A4V4ND72
A	0	GLU	-	expression tag	UNP A0A4V4ND72
A	1	LEU	-	expression tag	UNP A0A4V4ND72
A	66	CRQ	GLN	chromophore	UNP A0A4V4ND72
A	66	CRQ	TYR	chromophore	UNP A0A4V4ND72
A	66	CRQ	GLY	chromophore	UNP A0A4V4ND72
A	232	GLY	-	expression tag	UNP A0A4V4ND72
A	233	GLY	-	expression tag	UNP A0A4V4ND72
A	234	SER	-	expression tag	UNP A0A4V4ND72
A	235	GLY	-	expression tag	UNP A0A4V4ND72
A	236	GLY	-	expression tag	UNP A0A4V4ND72
A	237	SER	-	expression tag	UNP A0A4V4ND72
A	238	GLU	-	expression tag	UNP A0A4V4ND72
A	239	ASN	-	expression tag	UNP A0A4V4ND72
A	240	LEU	-	expression tag	UNP A0A4V4ND72
A	241	TYR	-	expression tag	UNP A0A4V4ND72
A	242	PHE	-	expression tag	UNP A0A4V4ND72
A	243	GLN	-	expression tag	UNP A0A4V4ND72
B	-36	MET	-	initiating methionine	UNP A0A4V4ND72
B	-35	GLY	-	expression tag	UNP A0A4V4ND72
B	-34	GLY	-	expression tag	UNP A0A4V4ND72
B	-33	SER	-	expression tag	UNP A0A4V4ND72
B	-32	HIS	-	expression tag	UNP A0A4V4ND72
B	-31	HIS	-	expression tag	UNP A0A4V4ND72
B	-30	HIS	-	expression tag	UNP A0A4V4ND72
B	-29	HIS	-	expression tag	UNP A0A4V4ND72
B	-28	HIS	-	expression tag	UNP A0A4V4ND72
B	-27	HIS	-	expression tag	UNP A0A4V4ND72
B	-26	GLY	-	expression tag	UNP A0A4V4ND72
B	-25	MET	-	expression tag	UNP A0A4V4ND72
B	-24	ALA	-	expression tag	UNP A0A4V4ND72
B	-23	SER	-	expression tag	UNP A0A4V4ND72

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	MET	-	expression tag	UNP A0A4V4ND72
B	-21	THR	-	expression tag	UNP A0A4V4ND72
B	-20	GLY	-	expression tag	UNP A0A4V4ND72
B	-19	GLY	-	expression tag	UNP A0A4V4ND72
B	-18	GLN	-	expression tag	UNP A0A4V4ND72
B	-17	GLN	-	expression tag	UNP A0A4V4ND72
B	-16	MET	-	expression tag	UNP A0A4V4ND72
B	-15	GLY	-	expression tag	UNP A0A4V4ND72
B	-14	ARG	-	expression tag	UNP A0A4V4ND72
B	-13	ASP	-	expression tag	UNP A0A4V4ND72
B	-12	LEU	-	expression tag	UNP A0A4V4ND72
B	-11	TYR	-	expression tag	UNP A0A4V4ND72
B	-10	ASP	-	expression tag	UNP A0A4V4ND72
B	-9	ASP	-	expression tag	UNP A0A4V4ND72
B	-8	ASP	-	expression tag	UNP A0A4V4ND72
B	-7	ASP	-	expression tag	UNP A0A4V4ND72
B	-6	LYS	-	expression tag	UNP A0A4V4ND72
B	-5	ASP	-	expression tag	UNP A0A4V4ND72
B	-4	ARG	-	expression tag	UNP A0A4V4ND72
B	-3	TRP	-	expression tag	UNP A0A4V4ND72
B	-2	GLY	-	expression tag	UNP A0A4V4ND72
B	-1	SER	-	expression tag	UNP A0A4V4ND72
B	0	GLU	-	expression tag	UNP A0A4V4ND72
B	1	LEU	-	expression tag	UNP A0A4V4ND72
B	66	CRQ	GLN	chromophore	UNP A0A4V4ND72
B	66	CRQ	TYR	chromophore	UNP A0A4V4ND72
B	66	CRQ	GLY	chromophore	UNP A0A4V4ND72
B	232	GLY	-	expression tag	UNP A0A4V4ND72
B	233	GLY	-	expression tag	UNP A0A4V4ND72
B	234	SER	-	expression tag	UNP A0A4V4ND72
B	235	GLY	-	expression tag	UNP A0A4V4ND72
B	236	GLY	-	expression tag	UNP A0A4V4ND72
B	237	SER	-	expression tag	UNP A0A4V4ND72
B	238	GLU	-	expression tag	UNP A0A4V4ND72
B	239	ASN	-	expression tag	UNP A0A4V4ND72
B	240	LEU	-	expression tag	UNP A0A4V4ND72
B	241	TYR	-	expression tag	UNP A0A4V4ND72
B	242	PHE	-	expression tag	UNP A0A4V4ND72
B	243	GLN	-	expression tag	UNP A0A4V4ND72

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Ca 2	0	0
2	B	2	Total 2	Ca 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	249	Total 249	O 249	0	0
3	B	227	Total 227	O 227	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Red fluorescent protein,grafted calcium-binding sequence



- Molecule 1: Red fluorescent protein,grafted calcium-binding sequence



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.69Å 79.07Å 65.68Å 90.00° 91.07° 90.00°	Depositor
Resolution (Å)	37.56 – 1.38 37.56 – 1.38	Depositor EDS
% Data completeness (in resolution range)	98.6 (37.56-1.38) 98.6 (37.56-1.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.38Å)	Xtriage
Refinement program	PHENIX (1.17.1_3660: ???)	Depositor
R, R_{free}	0.193 , 0.208 0.193 , 0.208	Depositor DCC
R_{free} test set	87494 reflections (2.25%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 28.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4014	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 81.35 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6250e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CRQ, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/1784	0.53	0/2398
1	B	0.24	0/1784	0.51	0/2398
All	All	0.24	0/3568	0.52	0/4796

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1767	0	1705	9	0
1	B	1767	0	1705	9	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	249	0	0	0	0
3	B	227	0	0	0	0
All	All	4014	0	3410	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:GLU:HG2	1:A:188:MET:HG2	1.79	0.64
1:A:17:ARG:HH11	1:A:17:ARG:HG2	1.65	0.60
1:B:162:GLU:HG2	1:B:188:MET:HG2	1.85	0.58
1:A:149:ARG:HH11	1:A:200:LYS:HD2	1.69	0.57
1:B:17:ARG:HH12	1:B:30:GLU:CD	2.17	0.51

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/278 (77%)	210 (98%)	5 (2%)	0	100	100
1	B	215/278 (77%)	210 (98%)	5 (2%)	0	100	100
All	All	430/556 (77%)	420 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/228 (81%)	185 (100%)	0	100	100
1	B	185/228 (81%)	185 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	370/456 (81%)	370 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	178	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRQ	B	66	1	24,25,26	1.26	1 (4%)	27,34,36	3.02	5 (18%)
1	CRQ	A	66	1	24,25,26	1.29	2 (8%)	27,34,36	3.02	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRQ	B	66	1	-	3/10/32/33	0/2/2/2
1	CRQ	A	66	1	-	4/10/32/33	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRQ	CB2-CA2	4.73	1.39	1.35
1	B	66	CRQ	CB2-CA2	4.66	1.39	1.35
1	A	66	CRQ	CA2-C2	-2.02	1.46	1.48

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRQ	O2-C2-CA2	-11.18	124.68	130.96
1	B	66	CRQ	O2-C2-CA2	-10.32	125.17	130.96
1	B	66	CRQ	CA2-C2-N3	8.98	107.62	103.37
1	A	66	CRQ	CA2-C2-N3	8.46	107.37	103.37
1	B	66	CRQ	C2-CA2-N2	-4.99	105.44	108.93

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66	CRQ	C1-CA1-CB1-CG1
1	A	66	CRQ	N2-CA2-CB2-CG2
1	A	66	CRQ	C2-CA2-CB2-CG2
1	B	66	CRQ	C1-CA1-CB1-CG1
1	B	66	CRQ	N2-CA2-CB2-CG2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	CRQ	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/278 (78%)	0.71	11 (5%) 35 39	14, 18, 26, 30	0
1	B	219/278 (78%)	0.86	12 (5%) 32 35	15, 19, 27, 31	0
All	All	438/556 (78%)	0.79	23 (5%) 33 36	14, 19, 27, 31	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	17	ARG	3.5
1	B	176	GLY	3.4
1	B	174	LYS	3.3
1	B	77	ALA	3.3
1	B	132	ASP	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRQ	A	66	24/25	0.87	0.14	18,22,26,28	0
1	CRQ	B	66	24/25	0.91	0.11	19,22,27,28	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	B	301	1/1	0.97	0.15	20,20,20,20	0
2	CA	A	302	1/1	0.98	0.15	20,20,20,20	0
2	CA	A	301	1/1	0.99	0.19	18,18,18,18	0
2	CA	B	302	1/1	0.99	0.18	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CA A 302:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.