



wwPDB NMR Structure Validation Summary Report ⓘ

Dec 24, 2024 – 06:00 PM EST

PDB ID : 2LUP
BMRB ID : 18534
Title : RDC refined solution structure of double-stranded RNA binding domain of
S. cerevisiae RNase III (rnt1p) in complex with the terminal RNA hairpin of
snr47 precursor
Authors : Wang, Z.; Feigon, J.
Deposited on : 2012-06-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 71%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463
RNA backbone	6643	756

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	32	
2	B	90	

2 Ensemble composition and analysis

This entry contains 16 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	B:366-B:447 (82)	0.86	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 9, 14
2	4, 7
3	6, 13
4	3, 8
5	10, 11
Single-model clusters	12; 15; 16

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2437 atoms, of which 1067 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (32-MER).

Mol	Chain	Residues	Atoms						Trace
1	A	32	Total	C	H	N	O	P	0
			1028	306	346	124	221	31	

- Molecule 2 is a protein called Ribonuclease 3.

Mol	Chain	Residues	Atoms						Trace
2	B	90	Total	C	H	N	O	S	0
			1409	429	721	129	127	3	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	364	GLY	-	expression tag	UNP Q02555
B	365	SER	-	expression tag	UNP Q02555

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: RNA (32-MER)

Chain A: 



- Molecule 2: Ribonuclease 3

Chain B: 



4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 5. Colouring as in section 4.1 above.


- Molecule 1: RNA (32-MER)

Chain A: 



- Molecule 2: Ribonuclease 3

Chain B: 



A428	E429	N430	A431	L432	R433	D434	K435	K436	N437	L438	D439	F440	Y441	A442	K443	Q444	R445	A446	A447	I448	P449	R450	S451	F452	S453
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 16 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1317
Number of shifts mapped to atoms	1317
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	71%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.13±0.01	0±0/763 (0.0± 0.0%)	1.99±0.01	43±1/1188 (3.7± 0.1%)
2	B	0.27±0.02	0±0/638 (0.0± 0.0%)	0.45±0.02	0±0/859 (0.0± 0.0%)
All	All	0.85	0/22416 (0.0%)	1.55	695/32752 (2.1%)

There are no bond-length outliers.

5 of 45 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	25	G	N7-C8-N9	9.54	117.87	113.10	10	16
1	A	11	G	N7-C8-N9	9.40	117.80	113.10	16	16
1	A	16	G	N7-C8-N9	9.33	117.77	113.10	6	16
1	A	19	G	N7-C8-N9	9.30	117.75	113.10	10	16
1	A	1	G	N7-C8-N9	9.29	117.74	113.10	16	16

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	682	346	346	94±10
2	B	630	664	663	126±9
All	All	20992	16160	16144	3434

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 92.

5 of 1029 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:370:ALA:HB3	2:B:428:ALA:HB1	1.03	1.25	15	2
2:B:374:LEU:HD11	2:B:428:ALA:HB2	1.01	1.32	16	4
2:B:370:ALA:HB1	2:B:428:ALA:HB1	1.00	1.28	14	14
2:B:380:TYR:CB	2:B:383:LEU:HD22	0.98	1.88	13	1
2:B:374:LEU:HD11	2:B:428:ALA:CB	0.97	1.88	16	5

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	82/90 (91%)	56±2 (68±3%)	18±3 (22±3%)	8±2 (10±2%)	1	9
All	All	1312/1440 (91%)	889 (68%)	289 (22%)	134 (10%)	1	9

5 of 23 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	397	ASP	16
2	B	420	ILE	16
2	B	392	LYS	15
2	B	399	ASN	13
2	B	391	LYS	10

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	65/72 (90%)	34±3 (53±4%)	31±3 (47±4%)	0 1
All	All	1040/1152 (90%)	548 (53%)	492 (47%)	0 1

5 of 58 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	376	SER	16
2	B	388	VAL	16
2	B	412	LEU	16
2	B	421	LYS	16
2	B	422	ILE	16

6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	31/32 (97%)	4±2 (14±5%)	2±1 (5±3%)	0.13±0.02
All	All	497/512 (97%)	67 (13%)	24 (5%)	0.13

The overall RNA backbone suiteness is 0.13.

5 of 13 unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	17	A	16
1	A	18	A	15
1	A	15	A	13
1	A	13	U	6
1	A	16	G	5

5 of 7 unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	17	A	13
1	A	16	G	5
1	A	3	G	2
1	A	1	G	1
1	A	28	A	1

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 71% for the well-defined parts and 71% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1317
Number of shifts mapped to atoms	1317
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	7

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	89	2.18 ± 0.24	Should be applied
$^{13}\text{C}_\beta$	82	2.66 ± 0.21	Should be applied
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	84	0.51 ± 0.38	None needed (imprecise)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 71%, i.e. 1241 atoms were assigned a chemical shift out of a possible 1738. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	327/413 (79%)	166/169 (98%)	82/164 (50%)	79/80 (99%)
Sidechain	444/670 (66%)	270/437 (62%)	168/199 (84%)	6/34 (18%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	8/54 (15%)	8/25 (32%)	0/27 (0%)	0/2 (0%)
Sugar	352/352 (100%)	192/192 (100%)	160/160 (100%)	0/0 (—%)
Base	110/249 (44%)	63/153 (41%)	47/55 (85%)	0/41 (0%)
Overall	1241/1738 (71%)	699/976 (72%)	457/605 (76%)	85/157 (54%)

7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	B	391	LYS	CG	33.40	19.35 – 30.45	7.7
1	B	444	GLN	CG	26.40	28.36 – 39.21	-6.8
1	B	421	LYS	CD	22.42	23.50 – 34.42	-6.0
1	B	392	LYS	CD	22.50	23.50 – 34.42	-5.9
1	B	420	ILE	CD1	22.46	5.18 – 21.60	5.5
1	B	451	SER	CB	55.96	56.28 – 71.32	-5.2
1	B	401	ILE	CG2	10.80	10.93 – 24.12	-5.1

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:



8 NMR restraints analysis [i](#)

8.1 Conformationally restricting restraints [i](#)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	2360
Intra-residue ($ i-j =0$)	832
Sequential ($ i-j =1$)	756
Medium range ($ i-j >1$ and $ i-j <5$)	329
Long range ($ i-j \geq 5$)	397
Inter-chain	46
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	227
Number of unmapped restraints	0
Number of restraints per residue	21.2
Number of long range restraints per residue ¹	3.3

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations [i](#)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model [i](#)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	25.7	0.2
0.2-0.5 (Medium)	8.8	0.5
>0.5 (Large)	5.2	3.2

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	0.9	2.1
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	832	35.3	24	2.9	1.0	0	0.0	0.0
Backbone-Backbone	77	3.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	599	25.4	22	3.7	0.9	0	0.0	0.0
Sidechain-Sidechain	156	6.6	2	1.3	0.1	0	0.0	0.0
Sequential (i-j =1)	756	32.0	46	6.1	1.9	3	0.4	0.1
Backbone-Backbone	191	8.1	4	2.1	0.2	1	0.5	0.0
Backbone-Sidechain	414	17.5	36	8.7	1.5	2	0.5	0.1
Sidechain-Sidechain	151	6.4	6	4.0	0.3	0	0.0	0.0
Medium range (i-j >1 & i-j <5)	329	13.9	38	11.6	1.6	2	0.6	0.1
Backbone-Backbone	83	3.5	7	8.4	0.3	0	0.0	0.0
Backbone-Sidechain	126	5.3	19	15.1	0.8	1	0.8	0.0
Sidechain-Sidechain	120	5.1	12	10.0	0.5	1	0.8	0.0
Long range (i-j ≥5)	397	16.8	32	8.1	1.4	0	0.0	0.0
Backbone-Backbone	53	2.2	5	9.4	0.2	0	0.0	0.0
Backbone-Sidechain	159	6.7	20	12.6	0.8	0	0.0	0.0
Sidechain-Sidechain	185	7.8	7	3.8	0.3	0	0.0	0.0
Inter-chain	46	1.9	2	4.3	0.1	0	0.0	0.0
Backbone-Backbone	1	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	29	1.2	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	16	0.7	2	12.5	0.1	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	2360	100.0	142	6.0	6.0	5	0.2	0.2
Backbone-Backbone	405	17.2	16	4.0	0.7	1	0.2	0.0
Backbone-Sidechain	1327	56.2	97	7.3	4.1	3	0.2	0.1
Sidechain-Sidechain	628	26.6	29	4.6	1.2	1	0.2	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	4	14	12	10	1	41	0.28	1.31	0.28	0.16
2	3	16	14	8	1	42	0.28	1.39	0.3	0.16
3	7	16	14	8	0	45	0.32	2.27	0.44	0.15
4	4	17	9	9	1	40	0.32	1.66	0.4	0.14
5	5	14	11	11	0	41	0.25	1.33	0.29	0.14
6	6	17	11	9	0	43	0.28	1.81	0.37	0.15
7	6	18	13	9	0	46	0.34	2.06	0.45	0.16
8	8	15	11	9	0	43	0.29	1.57	0.35	0.15
9	6	15	11	11	1	44	0.36	1.78	0.4	0.16
10	4	14	19	8	1	46	0.27	1.52	0.31	0.14

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	5	16	11	12	1	45	0.37	2.17	0.47	0.18
12	3	15	14	13	1	46	0.36	3.2	0.58	0.18
13	4	14	12	7	2	39	0.34	1.58	0.37	0.19
14	6	12	10	8	1	37	0.29	1.58	0.33	0.16
15	6	14	8	10	1	39	0.27	1.37	0.27	0.16
16	8	12	14	8	0	42	0.26	1.51	0.3	0.15

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble ⓘ


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2218(IR:808, SQ:710, MR:291, LR:365, IC:44) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
9	16	12	10	1	48	1	6.2
5	7	5	3	0	20	2	12.5
1	1	2	3	0	7	3	18.8
3	3	5	1	0	12	4	25.0
1	3	1	5	0	10	5	31.2
2	1	2	1	0	6	6	37.5
0	3	4	2	0	9	7	43.8
0	1	1	0	0	2	8	50.0
1	2	1	0	0	4	9	56.2
1	1	0	1	1	4	10	62.5
0	3	0	0	0	3	11	68.8
0	0	0	1	0	1	12	75.0
0	1	1	1	0	3	13	81.2
0	0	1	2	0	3	14	87.5
1	1	1	2	0	5	15	93.8
0	3	2	0	0	5	16	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,614)	1:29:A:U:H4'	1:30:A:C:H5''	16	1.16	0.15	1.12
(1,1484)	2:408:B:ASP:H	2:406:B:VAL:HG11	16	1.15	0.52	1.5
(1,1484)	2:408:B:ASP:H	2:406:B:VAL:HG12	16	1.15	0.52	1.5
(1,1484)	2:408:B:ASP:H	2:406:B:VAL:HG13	16	1.15	0.52	1.5
(1,2354)	2:426:B:ARG:HD3	2:422:B:ILE:HD11	16	1.06	0.21	1.07
(1,2354)	2:426:B:ARG:HD3	2:422:B:ILE:HD12	16	1.06	0.21	1.07
(1,2354)	2:426:B:ARG:HD3	2:422:B:ILE:HD13	16	1.06	0.21	1.07
(1,2354)	2:426:B:ARG:HD2	2:422:B:ILE:HD11	16	1.06	0.21	1.07
(1,2354)	2:426:B:ARG:HD2	2:422:B:ILE:HD12	16	1.06	0.21	1.07
(1,2354)	2:426:B:ARG:HD2	2:422:B:ILE:HD13	16	1.06	0.21	1.07
(1,2347)	2:406:B:VAL:HG12	2:407:B:GLY:HA2	16	0.99	0.43	1.2
(1,2347)	2:406:B:VAL:HG12	2:407:B:GLY:HA3	16	0.99	0.43	1.2
(1,2347)	2:406:B:VAL:HG11	2:407:B:GLY:HA2	16	0.99	0.43	1.2
(1,2347)	2:406:B:VAL:HG11	2:407:B:GLY:HA3	16	0.99	0.43	1.2
(1,2347)	2:406:B:VAL:HG13	2:407:B:GLY:HA2	16	0.99	0.43	1.2
(1,2347)	2:406:B:VAL:HG13	2:407:B:GLY:HA3	16	0.99	0.43	1.2

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,2347)	2:406:B:VAL:HG13	2:410:B:THR:HA	16	0.99	0.43	1.2
(1,2086)	2:440:B:PHE:H	2:441:B:TYR:HE1	16	0.15	0.03	0.15
(1,2086)	2:440:B:PHE:H	2:441:B:TYR:HE2	16	0.15	0.03	0.15
(1,2338)	2:389:B:THR:HB	2:388:B:VAL:HG11	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:388:B:VAL:HG12	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:388:B:VAL:HG13	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:388:B:VAL:HG21	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:388:B:VAL:HG22	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:388:B:VAL:HG23	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:390:B:VAL:HG11	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:390:B:VAL:HG12	15	0.31	0.05	0.34
(1,2338)	2:389:B:THR:HB	2:390:B:VAL:HG13	15	0.31	0.05	0.34
(1,2345)	2:403:B:GLU:H	2:388:B:VAL:HG11	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:388:B:VAL:HG12	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:388:B:VAL:HG13	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:388:B:VAL:HG21	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:388:B:VAL:HG22	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:388:B:VAL:HG23	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:390:B:VAL:HG11	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:390:B:VAL:HG12	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:390:B:VAL:HG13	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:390:B:VAL:HG21	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:390:B:VAL:HG22	15	0.26	0.11	0.22
(1,2345)	2:403:B:GLU:H	2:390:B:VAL:HG23	15	0.26	0.11	0.22
(1,1136)	2:387:B:TYR:H	2:387:B:TYR:HE1	15	0.23	0.05	0.22
(1,1136)	2:387:B:TYR:H	2:387:B:TYR:HE2	15	0.23	0.05	0.22
(1,2358)	2:433:B:ARG:H	2:412:B:LEU:HB2	15	0.23	0.09	0.21
(1,2358)	2:433:B:ARG:H	2:412:B:LEU:HB3	15	0.23	0.09	0.21
(1,2114)	2:441:B:TYR:HE1	2:444:B:GLN:HB2	15	0.16	0.03	0.16

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1129)	2:387:B:TYR:HE1	2:404:B:CYS:HB3	12	3.2
(1,1130)	2:387:B:TYR:HE1	2:404:B:CYS:HB2	12	2.51
(1,1129)	2:387:B:TYR:HE1	2:404:B:CYS:HB3	3	2.27
(1,1129)	2:387:B:TYR:HE1	2:404:B:CYS:HB3	11	2.17
(1,1129)	2:387:B:TYR:HE1	2:404:B:CYS:HB3	7	2.06
(1,1130)	2:387:B:TYR:HE1	2:404:B:CYS:HB2	6	1.81
(1,1129)	2:387:B:TYR:HE1	2:404:B:CYS:HB3	9	1.78
(1,1130)	2:387:B:TYR:HE1	2:404:B:CYS:HB2	11	1.7
(1,1130)	2:387:B:TYR:HE1	2:404:B:CYS:HB2	4	1.66
(1,1130)	2:387:B:TYR:HE1	2:404:B:CYS:HB2	3	1.62

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
DELTA	32	14.1	1	3.1	0.4	0	0.0	0.0
GAMMA	28	12.3	6	21.4	2.6	0	0.0	0.0
ALPHA	26	11.5	0	0.0	0.0	0	0.0	0.0
EPSILON	27	11.9	0	0.0	0.0	0	0.0	0.0
ZETA	27	11.9	0	0.0	0.0	0	0.0	0.0
BETA	27	11.9	0	0.0	0.0	0	0.0	0.0
CHI	32	14.1	0	0.0	0.0	0	0.0	0.0
NU2	28	12.3	2	7.1	0.9	0	0.0	0.0
Total	227	100.0	9	4.0	4.0	0	0.0	0.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model ⓘ

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations									Mean (°)	Max (°)	SD (°)	Median (°)
	DELTA	GAMMA	ALPHA	EPSILON	ZETA	BETA	CHI	NU2	Total				
1	0	1	0	0	0	0	0	0	1	1.09	1.09	0.0	1.09
2	0	1	0	0	0	0	0	0	1	1.1	1.1	0.0	1.1
3	0	1	0	0	0	0	0	0	1	1.14	1.14	0.0	1.14
4	0	2	0	0	0	0	0	0	2	1.13	1.14	0.01	1.13
5	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0	0	0	0	1	1	1.57	1.57	0.0	1.57
8	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	1	0	0	0	0	0	0	1	1.56	1.56	0.0	1.56
10	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
11	1	2	0	0	0	0	0	0	3	1.48	2.1	0.44	1.26
12	0	2	0	0	0	0	0	0	2	1.09	1.16	0.07	1.09
13	0	0	0	0	0	0	0	0	0	0.0	0.0	0.0	0.0
14	0	1	0	0	0	0	0	0	1	1.13	1.13	0.0	1.13
15	0	1	0	0	0	0	0	0	1	1.11	1.11	0.0	1.11
16	0	0	0	0	0	0	0	1	1	1.07	1.07	0.0	1.07

10.2.1 Bar graph : Dihedral violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right


10.3 Dihedral-angle violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

DELTA	GAMMA	Number of violated restraints							Fraction of the ensemble	
		ALPHA	EPSILON	ZETA	BETA	CHI	NU2	Total	Count ¹	%
1	4	0	0	0	0	0	2	7	1	6.2
0	0	0	0	0	0	0	0	0	2	12.5
0	0	0	0	0	0	0	0	0	3	18.8
0	2	0	0	0	0	0	0	2	4	25.0
0	0	0	0	0	0	0	0	0	5	31.2
0	0	0	0	0	0	0	0	0	6	37.5
0	0	0	0	0	0	0	0	0	7	43.8
0	0	0	0	0	0	0	0	0	8	50.0
0	0	0	0	0	0	0	0	0	9	56.2
0	0	0	0	0	0	0	0	0	10	62.5
0	0	0	0	0	0	0	0	0	11	68.8
0	0	0	0	0	0	0	0	0	12	75.0
0	0	0	0	0	0	0	0	0	13	81.2
0	0	0	0	0	0	0	0	0	14	87.5
0	0	0	0	0	0	0	0	0	15	93.8
0	0	0	0	0	0	0	0	0	16	100.0

¹ Number of models with violations


10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble [i](#)



10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

10.4.1 Histogram : Distribution of mean dihedral-angle violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints [i](#)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.


Key	Atom-1	Atom-2	Atom-3	Atom-4	Models ¹	Mean	SD ²	Median
(1,131)	1:23:A:G:O5'	1:23:A:G:C5'	1:23:A:G:C4'	1:23:A:G:C3'	4	1.23	0.19	1.13
(1,123)	1:11:A:G:O5'	1:11:A:G:C5'	1:11:A:G:C4'	1:11:A:G:C3'	4	1.13	0.09	1.12

¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,212)	1:17:A:A:C5'	1:17:A:A:C4'	1:17:A:A:C3'	1:17:A:A:O3'	11	2.1
(1,170)	1:3:A:G:C1'	1:3:A:G:C2'	1:3:A:G:C3'	1:3:A:G:C4'	7	1.57
(1,131)	1:23:A:G:O5'	1:23:A:G:C5'	1:23:A:G:C4'	1:23:A:G:C3'	9	1.56
(1,123)	1:11:A:G:O5'	1:11:A:G:C5'	1:11:A:G:C4'	1:11:A:G:C3'	11	1.26
(1,124)	1:12:A:U:O5'	1:12:A:U:C5'	1:12:A:U:C4'	1:12:A:U:C3'	12	1.16
(1,131)	1:23:A:G:O5'	1:23:A:G:C5'	1:23:A:G:C4'	1:23:A:G:C3'	3	1.14
(1,123)	1:11:A:G:O5'	1:11:A:G:C5'	1:11:A:G:C4'	1:11:A:G:C3'	4	1.14
(1,131)	1:23:A:G:O5'	1:23:A:G:C5'	1:23:A:G:C4'	1:23:A:G:C3'	14	1.13
(1,121)	1:9:A:A:O5'	1:9:A:A:C5'	1:9:A:A:C4'	1:9:A:A:C3'	4	1.12
(1,123)	1:11:A:G:O5'	1:11:A:G:C5'	1:11:A:G:C4'	1:11:A:G:C3'	15	1.11