



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 01:27 PM EDT

PDB ID : 3LWV
Title : Structure of H/ACA RNP bound to a substrate RNA containing 2'-deoxyuridine
Authors : Zhou, J.; Liang, B.; Lv, C.; Yang, W.; Li, H.
Deposited on : 2010-02-24
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)
RNA backbone	3690	1181 (2.80-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	<div> <div> <div></div> <div>67%</div> <div>22%</div> <div>• 7%</div> </div> </div>
2	B	60	<div> <div> <div></div> <div>70%</div> <div>17%</div> <div>• 12%</div> </div> </div>
3	C	123	<div> <div> <div></div> <div>67%</div> <div>29%</div> <div>• •</div> </div> </div>
4	D	58	<div> <div> <div></div> <div>52%</div> <div>31%</div> <div>9%</div> <div>9%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	13	<div> <div>15%</div> <div>69%</div> <div>15%</div> <div>15%</div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable tRNA pseudouridine synthase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	0	0
			2521	1629	432	450	10			

- Molecule 2 is a protein called Ribosome biogenesis protein Nop10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	53	Total	C	N	O	S	0	0	0
			444	283	86	71	4			

- Molecule 3 is a protein called Large ribosomal subunit protein eL8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	120	Total	C	N	O	S	0	0	0
			916	585	151	177	3			

- Molecule 4 is a RNA chain called H/ACA RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	58	Total	C	N	O	P	0	0	0
			1236	552	227	400	57			

- Molecule 5 is a RNA chain called 5'-R(*GP*AP*GP*CP*GP*(du)P*GP*CP*GP*GP*UP*UP*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	11	Total	C	N	O	P	0	0	0
			234	105	42	77	10			

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	Zn 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	5	Total 5	O 5	0	0
7	B	1	Total 1	O 1	0	0
7	D	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable tRNA pseudouridine synthase B




- Molecule 2: Ribosome biogenesis protein Nop10



- Molecule 3: Large ribosomal subunit protein eL8




- Molecule 4: H/ACA RNA



- Molecule 5: 5'-R(*GP*AP*GP*CP*GP*(du)P*GP*CP*GP*GP*UP*UP*U)-3'

Chain E: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	186.84Å 64.33Å 83.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.64 – 2.50 34.64 – 2.50	Depositor EDS
% Data completeness (in resolution range)	51.2 (34.64-2.50) 96.5 (34.64-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.12 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.202 , 0.245 0.198 , 0.243	Depositor DCC
R_{free} test set	1725 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	66.3	Xtriage
Anisotropy	0.139	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5361	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2578	0.60	0/3488
2	B	0.51	0/457	0.69	2/613 (0.3%)
3	C	0.36	0/928	0.53	0/1253
4	D	0.55	0/1381	1.29	13/2152 (0.6%)
5	E	0.57	0/260	1.08	0/402
All	All	0.46	0/5604	0.87	15/7908 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	24	G	P-O3'-C3'	10.32	132.09	119.70
4	D	40	U	O4'-C1'-N1	9.72	115.98	108.20
4	D	56	C	P-O3'-C3'	9.54	131.15	119.70
4	D	40	U	P-O3'-C3'	8.12	129.45	119.70
4	D	56	C	C6-N1-C2	7.54	123.31	120.30
4	D	26	U	O4'-C1'-N1	-7.47	102.22	108.20
4	D	56	C	O4'-C1'-N1	6.92	113.73	108.20
2	B	8	CYS	CA-CB-SG	6.32	125.38	114.00
4	D	41	C	O4'-C1'-N1	5.69	112.75	108.20
4	D	26	U	N1-C1'-C2'	-5.42	106.04	112.00
4	D	45	C	O4'-C1'-N1	-5.39	103.88	108.20
2	B	20	CYS	CA-CB-SG	5.22	123.40	114.00
4	D	32	A	C3'-C2'-C1'	5.20	105.66	101.50
4	D	30	U	P-O3'-C3'	5.15	125.88	119.70
4	D	54	C	P-O3'-C3'	5.00	125.70	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2521	0	2588	75	0
2	B	444	0	451	6	0
3	C	916	0	964	23	0
4	D	1236	0	633	20	0
5	E	234	0	121	1	0
6	B	1	0	0	0	0
7	A	5	0	0	0	0
7	B	1	0	0	0	0
7	D	3	0	0	1	0
All	All	5361	0	4757	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (118) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:LYS:H	1:A:87:LYS:CE	1.80	0.94
1:A:20:GLU:HG2	1:A:282:ALA:HB3	1.51	0.92
1:A:141:GLN:HE22	1:A:182:TYR:H	1.14	0.88
1:A:11:ARG:N	1:A:11:ARG:HE	1.76	0.84
4:D:32:A:H2'	4:D:33:G:C8	2.17	0.80
1:A:87:LYS:H	1:A:87:LYS:HE3	1.47	0.78
1:A:87:LYS:H	1:A:87:LYS:HE2	1.50	0.76
2:B:10:LYS:HE3	2:B:10:LYS:HA	1.71	0.72
1:A:99:ALA:O	1:A:102:VAL:HG22	1.91	0.70
1:A:85:ASP:HB3	1:A:87:LYS:HZ1	1.58	0.68
3:C:119:VAL:O	3:C:123:MET:HG2	1.94	0.68
3:C:79:ILE:HG12	3:C:80:TYR:H	1.61	0.66
3:C:63:ILE:HG22	3:C:64:VAL:HG13	1.78	0.65
3:C:111:LEU:O	3:C:115:ILE:HG13	1.98	0.63
1:A:107:LEU:N	1:A:108:PRO:HD2	2.13	0.63
1:A:86:PRO:HD2	1:A:87:LYS:HZ1	1.61	0.63
3:C:79:ILE:HG12	3:C:80:TYR:N	2.14	0.62
3:C:16:LEU:HD13	3:C:118:LYS:HD3	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:LYS:NZ	1:A:304:LYS:HB3	2.15	0.61
1:A:20:GLU:HG2	1:A:282:ALA:CB	2.29	0.60
1:A:85:ASP:HB3	1:A:87:LYS:NZ	2.15	0.60
3:C:44:VAL:HG13	3:C:76:ILE:HD12	1.83	0.60
3:C:7:TYR:HD2	3:C:61:GLU:HG2	1.67	0.59
1:A:220:LEU:O	1:A:224:VAL:HG13	2.02	0.59
1:A:87:LYS:HE2	1:A:87:LYS:N	2.18	0.58
1:A:31:ASN:HD22	1:A:32:PRO:N	2.02	0.57
1:A:20:GLU:CG	1:A:282:ALA:HB3	2.29	0.57
4:D:32:A:H2'	4:D:33:G:H8	1.68	0.57
1:A:264:ALA:O	1:A:268:HIS:HD2	1.87	0.57
3:C:7:TYR:CD2	3:C:61:GLU:HG2	2.40	0.57
1:A:31:ASN:HD22	1:A:31:ASN:C	2.08	0.57
1:A:210:PRO:HG3	1:A:252:HIS:ND1	2.20	0.57
1:A:68:TRP:CE2	2:B:37:PRO:HG3	2.40	0.56
1:A:112:GLU:HB3	1:A:207:ARG:HB3	1.86	0.56
3:C:68:PRO:HB2	3:C:69:PRO:HD3	1.86	0.56
3:C:60:PRO:HD2	3:C:63:ILE:HD13	1.87	0.56
1:A:275:PRO:HB2	4:D:57:A:H5''	1.90	0.54
1:A:87:LYS:CE	1:A:87:LYS:N	2.62	0.53
1:A:233:ASP:OD2	1:A:233:ASP:N	2.42	0.53
2:B:10:LYS:HE3	2:B:10:LYS:CA	2.36	0.53
3:C:53:ILE:HD12	3:C:53:ILE:N	2.25	0.52
4:D:28:A:O2'	4:D:31:G:H4'	2.10	0.52
1:A:61:THR:HG22	1:A:86:PRO:HG3	1.90	0.52
1:A:31:ASN:ND2	1:A:33:ASP:H	2.08	0.51
5:E:13:G:H2'	5:E:14:G:H5'	1.91	0.51
1:A:154:ARG:NH1	1:A:156:ARG:HH11	2.08	0.51
1:A:207:ARG:NH1	1:A:212:LYS:HE3	2.25	0.51
1:A:71:LYS:HG3	1:A:72:ILE:N	2.25	0.51
4:D:56:C:H4'	4:D:57:A:O5'	2.10	0.51
1:A:31:ASN:HD22	1:A:32:PRO:CD	2.24	0.51
1:A:86:PRO:HD2	1:A:87:LYS:NZ	2.26	0.51
1:A:205:ARG:HD3	1:A:208:SER:HB2	1.93	0.51
1:A:213:GLU:HG2	1:A:217:LEU:HD22	1.92	0.51
3:C:18:GLU:O	3:C:22:GLN:HG2	2.10	0.50
1:A:154:ARG:NH2	1:A:180:GLY:O	2.44	0.50
1:A:11:ARG:N	1:A:11:ARG:NE	2.53	0.50
1:A:114:VAL:HG22	1:A:174:ARG:HG2	1.93	0.50
3:C:60:PRO:HD2	3:C:63:ILE:CD1	2.42	0.50
3:C:39:GLU:HG3	4:D:31:G:H1	1.76	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:GLU:CD	1:A:26:GLU:H	2.16	0.49
1:A:23:ILE:HD13	1:A:279:LYS:HG3	1.95	0.48
4:D:11:A:H2'	4:D:12:A:C8	2.49	0.47
1:A:112:GLU:CD	1:A:160:TYR:OH	2.53	0.47
1:A:107:LEU:N	1:A:108:PRO:CD	2.78	0.47
4:D:30:U:H4'	4:D:31:G:OP2	2.15	0.47
1:A:14:PRO:O	1:A:17:ILE:HG23	2.14	0.46
1:A:17:ILE:O	1:A:17:ILE:HG13	2.13	0.46
1:A:130:GLN:O	1:A:133:LYS:HE2	2.16	0.46
1:A:63:HIS:CE1	4:D:41:C:C5	3.04	0.46
1:A:232:GLU:C	1:A:233:ASP:OD2	2.55	0.46
4:D:28:A:C5	4:D:32:A:H4'	2.51	0.46
1:A:310:GLN:O	1:A:314:GLU:HG2	2.17	0.44
1:A:312:MET:HG2	1:A:320:ALA:HB1	1.99	0.44
3:C:34:ARG:HG3	3:C:103:ILE:CG1	2.47	0.44
1:A:230:TRP:HA	1:A:235:ILE:O	2.17	0.44
1:A:213:GLU:HG2	1:A:217:LEU:CD2	2.47	0.44
1:A:287:GLY:HA2	1:A:304:LYS:HE2	1.99	0.44
1:A:316:THR:O	1:A:317:LYS:HG3	2.17	0.44
3:C:115:ILE:O	3:C:119:VAL:HG13	2.17	0.44
3:C:116:ALA:O	3:C:119:VAL:HG22	2.18	0.44
4:D:26:U:H6	4:D:26:U:H2'	1.39	0.44
4:D:45:C:O2	4:D:45:C:H2'	2.17	0.44
1:A:187:ILE:HD12	1:A:200:MET:HB2	2.00	0.43
4:D:58:A:N3	4:D:58:A:H2'	2.32	0.43
1:A:264:ALA:O	1:A:268:HIS:CD2	2.70	0.43
3:C:32:LYS:NZ	3:C:103:ILE:HG21	2.34	0.43
4:D:11:A:H2'	4:D:12:A:H8	1.83	0.43
4:D:41:C:H2'	4:D:42:G:O4'	2.19	0.43
1:A:34:TRP:CE2	1:A:334:PRO:HG2	2.54	0.42
1:A:133:LYS:HE2	1:A:133:LYS:HB3	1.61	0.42
1:A:170:ASP:OD1	2:B:4:ARG:NH1	2.52	0.42
4:D:8:G:O2'	4:D:9:G:H5'	2.19	0.42
3:C:51:LEU:HD12	3:C:77:PRO:O	2.19	0.42
1:A:18:LYS:HE3	1:A:18:LYS:HB2	1.54	0.42
1:A:274:VAL:N	1:A:275:PRO:CD	2.83	0.42
3:C:21:LEU:O	3:C:25:GLU:HB2	2.19	0.42
1:A:268:HIS:CE1	1:A:330:ARG:HD3	2.54	0.42
1:A:285:LYS:NZ	1:A:285:LYS:HB2	2.33	0.42
1:A:31:ASN:HA	1:A:32:PRO:HD3	1.85	0.42
1:A:127:LYS:HA	1:A:127:LYS:HD3	1.75	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:333:TYR:HA	1:A:334:PRO:HD3	1.86	0.42
4:D:17:G:H2'	4:D:18:C:O4'	2.20	0.42
2:B:48:TRP:CD2	3:C:65:ALA:HB1	2.55	0.41
1:A:221:HIS:CD2	1:A:221:HIS:H	2.36	0.41
1:A:270:ALA:HB2	4:D:55:A:C5	2.55	0.41
4:D:1:G:H5'	7:D:59:HOH:O	2.20	0.41
4:D:23:U:C2'	4:D:24:G:H5'	2.50	0.41
1:A:261:SER:HA	1:A:336:LEU:HD23	2.02	0.41
1:A:328:MET:HA	1:A:329:PRO:HD3	1.96	0.41
2:B:39:ASP:N	2:B:40:PRO:HD3	2.35	0.41
1:A:31:ASN:C	1:A:31:ASN:ND2	2.74	0.41
1:A:44:GLU:O	1:A:48:GLN:HG2	2.21	0.41
1:A:302:LEU:HD22	1:A:327:PHE:HE1	1.85	0.41
1:A:51:VAL:HG11	1:A:211:PHE:CZ	2.56	0.41
1:A:154:ARG:NH1	1:A:156:ARG:NH1	2.69	0.41
1:A:250:VAL:O	1:A:250:VAL:CG1	2.69	0.40
3:C:37:THR:HA	3:C:99:SER:OG	2.21	0.40
1:A:247:GLU:HG2	1:A:297:ASP:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/340 (92%)	300 (96%)	12 (4%)	1 (0%)	37	56
2	B	51/60 (85%)	50 (98%)	1 (2%)	0	100	100
3	C	118/123 (96%)	113 (96%)	5 (4%)	0	100	100
All	All	482/523 (92%)	463 (96%)	18 (4%)	1 (0%)	44	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	279	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/289 (93%)	250 (93%)	18 (7%)	13	28
2	B	48/54 (89%)	44 (92%)	4 (8%)	9	19
3	C	97/99 (98%)	89 (92%)	8 (8%)	9	19
All	All	413/442 (93%)	383 (93%)	30 (7%)	11	24

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ARG
1	A	26	GLU
1	A	31	ASN
1	A	53	ASN
1	A	71	LYS
1	A	76	GLU
1	A	87	LYS
1	A	106	LEU
1	A	127	LYS
1	A	133	LYS
1	A	140	ILE
1	A	156	ARG
1	A	224	VAL
1	A	233	ASP
1	A	250	VAL
1	A	302	LEU
1	A	304	LYS
1	A	336	LEU
2	B	7	LYS
2	B	10	LYS
2	B	11	CYS
2	B	36	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	15	GLU
3	C	25	GLU
3	C	35	LYS
3	C	59	ASP
3	C	70	LEU
3	C	79	ILE
3	C	83	SER
3	C	120	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	46	HIS
1	A	53	ASN
1	A	63	HIS
1	A	141	GLN
1	A	221	HIS
1	A	268	HIS
3	C	48	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	D	57/58 (98%)	11 (19%)	3 (5%)
5	E	9/13 (69%)	2 (22%)	0
All	All	66/71 (92%)	13 (19%)	3 (4%)

All (13) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	D	10	A
4	D	25	A
4	D	26	U
4	D	27	C
4	D	28	A
4	D	29	A
4	D	32	A
4	D	41	C
4	D	56	C

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	57	A
4	D	58	A
5	E	13	G
5	E	14	G

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	D	24	G
4	D	32	A
4	D	56	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/340 (93%)	-0.16	2 (0%) 85 83	52, 68, 98, 130	0
2	B	53/60 (88%)	0.04	1 (1%) 66 63	54, 73, 97, 101	0
3	C	120/123 (97%)	0.10	2 (1%) 69 65	65, 91, 125, 141	0
4	D	58/58 (100%)	-0.26	0 100 100	68, 87, 152, 172	0
5	E	11/13 (84%)	0.79	2 (18%) 4 4	115, 156, 190, 196	0
All	All	559/594 (94%)	-0.08	7 (1%) 74 71	52, 77, 123, 196	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	20	ALA	2.8
5	E	9	G	2.7
5	E	12	C	2.6
2	B	55	ILE	2.5
1	A	107	LEU	2.4
1	A	76	GLU	2.1
3	C	104	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	ZN	B	61	1/1	0.95	0.16	583,583,583,583	0

6.5 Other polymers [i](#)

There are no such residues in this entry.