



wwPDB EM Validation Summary Report ⓘ

Oct 8, 2025 – 04:17 PM EDT

PDB ID : 9N11 / pdb_00009n11
EMDB ID : EMD-48802
Title : Mfd-bound E.coli RNA polymerase elongation complex + ADP-BeF3 - L1.5 state
Authors : Brewer, J.J.; Darst, S.A.; Campbell, E.A.
Deposited on : 2025-01-24
Resolution : 4.29 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev129
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
RNA backbone	6643	2191	-
Q-score	-	25397	4494 (3.79 - 4.79)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1166	
2	G	234	
2	H	234	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	1340	
4	J	1358	
5	K	91	
6	R	20	
7	N	64	
8	T	64	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 70698 atoms, of which 34981 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription-repair-coupling factor.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	1089	Total	C	H	N	O	S	0	0
			17246	5463	8626	1537	1584	36		

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	HIS	-	expression tag	UNP A0A024L3Y3
A	-16	HIS	-	expression tag	UNP A0A024L3Y3
A	-15	HIS	-	expression tag	UNP A0A024L3Y3
A	-14	HIS	-	expression tag	UNP A0A024L3Y3
A	-13	HIS	-	expression tag	UNP A0A024L3Y3
A	-12	HIS	-	expression tag	UNP A0A024L3Y3
A	-11	SER	-	expression tag	UNP A0A024L3Y3
A	-10	SER	-	expression tag	UNP A0A024L3Y3
A	-9	GLY	-	expression tag	UNP A0A024L3Y3
A	-8	LEU	-	expression tag	UNP A0A024L3Y3
A	-7	GLU	-	expression tag	UNP A0A024L3Y3
A	-6	VAL	-	expression tag	UNP A0A024L3Y3
A	-5	LEU	-	expression tag	UNP A0A024L3Y3
A	-4	PHE	-	expression tag	UNP A0A024L3Y3
A	-3	GLN	-	expression tag	UNP A0A024L3Y3
A	-2	GLY	-	expression tag	UNP A0A024L3Y3
A	-1	PRO	-	expression tag	UNP A0A024L3Y3
A	0	HIS	-	expression tag	UNP A0A024L3Y3

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	G	224	Total	C	H	N	O	S	0	0
			3488	1080	1763	303	336	6		
2	H	217	Total	C	H	N	O	S	0	0
			3368	1043	1701	293	325	6		

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	I	1316	Total	C	H	N	O	S	0	0
			20754	6507	10386	1808	2010	43		

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	J	1336	Total	C	H	N	O	S	0	0
			20985	6524	10602	1852	1957	50		

- Molecule 5 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	K	79	Total	C	H	N	O	S	0	0
			1261	382	634	118	126	1		

- Molecule 6 is a RNA chain called RNA (5'-R(P*GP*GP*AP*GP*AP*GP*GP*UP*A)-3').

Mol	Chain	Residues	Atoms						AltConf	Trace
6	R	9	Total	C	H	N	O	P	0	0
			300	89	99	42	61	9		

- Molecule 7 is a DNA chain called DNA (49-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
7	N	49	Total	C	H	N	O	P	0	0
			1560	478	549	194	290	49		

- Molecule 8 is a DNA chain called DNA (55-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
8	T	55	Total	C	H	N	O	P	0	0
			1733	531	621	192	335	54		

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
9	J	1	Total	Mg	0
			1	1	


- Molecule 10 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
10	J	2	Total	Zn	0
			2	2	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcription-repair-coupling factor






• Molecule 2: DNA-directed RNA polymerase subunit alpha

Chain G: 87% 9%




• Molecule 2: DNA-directed RNA polymerase subunit alpha


Chain H: 83% 9% 7%




• Molecule 3: DNA-directed RNA polymerase subunit beta





• Molecule 4: DNA-directed RNA polymerase subunit beta'







- Molecule 5: DNA-directed RNA polymerase subunit omega




- Molecule 6: RNA (5'-R(P*GP*GP*AP*GP*AP*GP*GP*UP*A)-3')



- Molecule 7: DNA (49-MER)



- Molecule 8: DNA (55-MER)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	13808	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	51.2	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.787	Depositor
Minimum map value	-0.342	Depositor
Average map value	0.005	Depositor
Map value standard deviation	0.034	Depositor
Recommended contour level	0.168	Depositor
Map size (Å)	324.9, 324.9, 324.9	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.083, 1.083, 1.083	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/8792	0.33	0/11907
2	G	0.22	0/1745	0.36	0/2366
2	H	0.19	0/1686	0.39	0/2286
3	I	0.21	0/10533	0.39	0/14214
4	J	0.21	0/10540	0.39	0/14231
5	K	0.16	0/629	0.34	0/847
6	R	0.27	0/226	0.32	0/352
7	N	0.26	0/1135	0.46	0/1748
8	T	0.29	0/1242	0.50	0/1913
All	All	0.20	0/36528	0.38	0/49864

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	779	PRO	Peptide
1	A	784	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8620	8626	8611	65	0
2	G	1725	1763	1763	12	0
2	H	1667	1701	1699	16	0
3	I	10368	10386	10379	77	0
4	J	10383	10602	10597	132	0
5	K	627	634	634	2	0
6	R	201	99	99	0	0
7	N	1011	549	550	2	0
8	T	1112	621	622	6	0
9	J	1	0	0	0	0
10	J	2	0	0	0	0
All	All	35717	34981	34954	298	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 298 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:770:MET:CE	1:A:975:TYR:HE2	1.72	1.01
1:A:770:MET:CE	1:A:975:TYR:CE2	2.44	0.99
1:A:770:MET:HE2	1:A:975:TYR:CE2	1.97	0.99
4:J:848:VAL:HG12	4:J:858:VAL:HG22	1.50	0.91
4:J:1060:VAL:HG22	4:J:1106:ILE:HG22	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1081/1166 (93%)	1004 (93%)	74 (7%)	3 (0%)	37	72
2	G	220/234 (94%)	200 (91%)	20 (9%)	0	100	100
2	H	213/234 (91%)	193 (91%)	20 (9%)	0	100	100
3	I	1312/1340 (98%)	1220 (93%)	91 (7%)	1 (0%)	48	83
4	J	1330/1358 (98%)	1251 (94%)	79 (6%)	0	100	100
5	K	77/91 (85%)	71 (92%)	6 (8%)	0	100	100
All	All	4233/4423 (96%)	3939 (93%)	290 (7%)	4 (0%)	50	83

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	GLN
3	I	31	GLN
1	A	127	ALA
1	A	345	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	909/989 (92%)	906 (100%)	3 (0%)	91	92
2	G	191/202 (95%)	190 (100%)	1 (0%)	86	90
2	H	184/202 (91%)	183 (100%)	1 (0%)	86	90
3	I	1132/1155 (98%)	1119 (99%)	13 (1%)	70	80
4	J	1118/1134 (99%)	1099 (98%)	19 (2%)	56	73
5	K	67/75 (89%)	67 (100%)	0	100	100
All	All	3601/3757 (96%)	3564 (99%)	37 (1%)	71	82

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	J	727	ASP
4	J	1115	ILE
4	J	809	VAL
4	J	963	VAL
3	I	1154	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
3	I	513	GLN
4	J	805	GLN
3	I	761	GLN
4	J	1195	GLN
4	J	266	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
6	R	8/20 (40%)	1 (12%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
6	R	14	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-48802. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 151




Y Index: 149




Z Index: 149


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.168. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 344 nm³; this corresponds to an approximate mass of 311 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.233 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.233 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.29	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	7.75	10.66	7.94

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.75 differs from the reported value 4.29 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-48802 and PDB model 9N11. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.168 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.168).

9.4 Atom inclusion [i](#)



At the recommended contour level, 63% of all backbone atoms, 63% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.168) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.6330	<div></div> 0.2060
A	<div></div> 0.1980	<div></div> 0.0840
G	<div></div> 0.7970	<div></div> 0.2560
H	<div></div> 0.7980	<div></div> 0.2400
I	<div></div> 0.7780	<div></div> 0.2530
J	<div></div> 0.7750	<div></div> 0.2390
K	<div></div> 0.6640	<div></div> 0.2030
N	<div></div> 0.8040	<div></div> 0.2360
R	<div></div> 1.0000	<div></div> 0.3460
T	<div></div> 0.8050	<div></div> 0.2360

1.0

0.0

<0.0