



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 03:58 PM EDT

PDB ID : 3N6R  
Title : CRYSTAL STRUCTURE OF the holoenzyme of PROPIONYL-COA CARBOXYLASE (PCC)  
Authors : Huang, C.S.; Sadre-Bazzaz, K.; Tong, L.  
Deposited on : 2010-05-26  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	681	<div> <div>33%</div> <div>44%</div> <div>9%</div> <div>13%</div> </div>
1	C	681	<div> <div>31%</div> <div>45%</div> <div>10%</div> <div>13%</div> </div>
1	E	681	<div> <div>32%</div> <div>44%</div> <div>10%</div> <div>13%</div> </div>
1	G	681	<div> <div>35%</div> <div>50%</div> <div>9%</div> <div>5%</div> </div>
1	I	681	<div> <div>35%</div> <div>50%</div> <div>10%</div> <div>5%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	681	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>2%35%50%9%5%</div></div>
2	B	531	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>%49%43%• 5%</div></div>
2	D	531	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>49%42%• • 5%</div></div>
2	F	531	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>51%40%• 5%</div></div>
2	H	531	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>51%40%• • 5%</div></div>
2	J	531	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>51%39%• • 5%</div></div>
2	L	531	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>51%40%• 5%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 51921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Propionyl-CoA carboxylase, alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	591	Total	C	N	O	S	0	0	0
			4507	2828	794	857	28			
1	C	591	Total	C	N	O	S	0	0	0
			4507	2828	794	857	28			
1	E	591	Total	C	N	O	S	0	0	0
			4507	2828	794	857	28			
1	G	646	Total	C	N	O	S	0	0	0
			4950	3108	869	942	31			
1	I	646	Total	C	N	O	S	0	0	0
			4950	3108	869	942	31			
1	K	646	Total	C	N	O	S	0	0	0
			4950	3108	869	942	31			

- Molecule 2 is a protein called Propionyl-CoA carboxylase, beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	506	Total	C	N	O	S	0	0	0
			3910	2462	683	744	21			
2	D	506	Total	C	N	O	S	0	0	0
			3910	2462	683	744	21			
2	F	506	Total	C	N	O	S	0	0	0
			3910	2462	683	744	21			
2	H	506	Total	C	N	O	S	0	0	0
			3910	2462	683	744	21			
2	J	506	Total	C	N	O	S	0	0	0
			3910	2462	683	744	21			
2	L	506	Total	C	N	O	S	0	0	0
			3910	2462	683	744	21			

There are 126 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	11	MET	-	EXPRESSION TAG	UNP Q168G2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	12	GLY	-	EXPRESSION TAG	UNP Q168G2
B	13	SER	-	EXPRESSION TAG	UNP Q168G2
B	14	SER	-	EXPRESSION TAG	UNP Q168G2
B	15	HIS	-	EXPRESSION TAG	UNP Q168G2
B	16	HIS	-	EXPRESSION TAG	UNP Q168G2
B	17	HIS	-	EXPRESSION TAG	UNP Q168G2
B	18	HIS	-	EXPRESSION TAG	UNP Q168G2
B	19	HIS	-	EXPRESSION TAG	UNP Q168G2
B	20	HIS	-	EXPRESSION TAG	UNP Q168G2
B	21	SER	-	EXPRESSION TAG	UNP Q168G2
B	22	SER	-	EXPRESSION TAG	UNP Q168G2
B	23	GLY	-	EXPRESSION TAG	UNP Q168G2
B	24	LEU	-	EXPRESSION TAG	UNP Q168G2
B	25	VAL	-	EXPRESSION TAG	UNP Q168G2
B	26	PRO	-	EXPRESSION TAG	UNP Q168G2
B	27	ARG	-	EXPRESSION TAG	UNP Q168G2
B	28	GLY	-	EXPRESSION TAG	UNP Q168G2
B	29	SER	-	EXPRESSION TAG	UNP Q168G2
B	30	HIS	-	EXPRESSION TAG	UNP Q168G2
B	31	MET	-	EXPRESSION TAG	UNP Q168G2
D	11	MET	-	EXPRESSION TAG	UNP Q168G2
D	12	GLY	-	EXPRESSION TAG	UNP Q168G2
D	13	SER	-	EXPRESSION TAG	UNP Q168G2
D	14	SER	-	EXPRESSION TAG	UNP Q168G2
D	15	HIS	-	EXPRESSION TAG	UNP Q168G2
D	16	HIS	-	EXPRESSION TAG	UNP Q168G2
D	17	HIS	-	EXPRESSION TAG	UNP Q168G2
D	18	HIS	-	EXPRESSION TAG	UNP Q168G2
D	19	HIS	-	EXPRESSION TAG	UNP Q168G2
D	20	HIS	-	EXPRESSION TAG	UNP Q168G2
D	21	SER	-	EXPRESSION TAG	UNP Q168G2
D	22	SER	-	EXPRESSION TAG	UNP Q168G2
D	23	GLY	-	EXPRESSION TAG	UNP Q168G2
D	24	LEU	-	EXPRESSION TAG	UNP Q168G2
D	25	VAL	-	EXPRESSION TAG	UNP Q168G2
D	26	PRO	-	EXPRESSION TAG	UNP Q168G2
D	27	ARG	-	EXPRESSION TAG	UNP Q168G2
D	28	GLY	-	EXPRESSION TAG	UNP Q168G2
D	29	SER	-	EXPRESSION TAG	UNP Q168G2
D	30	HIS	-	EXPRESSION TAG	UNP Q168G2
D	31	MET	-	EXPRESSION TAG	UNP Q168G2
F	11	MET	-	EXPRESSION TAG	UNP Q168G2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	12	GLY	-	EXPRESSION TAG	UNP Q168G2
F	13	SER	-	EXPRESSION TAG	UNP Q168G2
F	14	SER	-	EXPRESSION TAG	UNP Q168G2
F	15	HIS	-	EXPRESSION TAG	UNP Q168G2
F	16	HIS	-	EXPRESSION TAG	UNP Q168G2
F	17	HIS	-	EXPRESSION TAG	UNP Q168G2
F	18	HIS	-	EXPRESSION TAG	UNP Q168G2
F	19	HIS	-	EXPRESSION TAG	UNP Q168G2
F	20	HIS	-	EXPRESSION TAG	UNP Q168G2
F	21	SER	-	EXPRESSION TAG	UNP Q168G2
F	22	SER	-	EXPRESSION TAG	UNP Q168G2
F	23	GLY	-	EXPRESSION TAG	UNP Q168G2
F	24	LEU	-	EXPRESSION TAG	UNP Q168G2
F	25	VAL	-	EXPRESSION TAG	UNP Q168G2
F	26	PRO	-	EXPRESSION TAG	UNP Q168G2
F	27	ARG	-	EXPRESSION TAG	UNP Q168G2
F	28	GLY	-	EXPRESSION TAG	UNP Q168G2
F	29	SER	-	EXPRESSION TAG	UNP Q168G2
F	30	HIS	-	EXPRESSION TAG	UNP Q168G2
F	31	MET	-	EXPRESSION TAG	UNP Q168G2
H	11	MET	-	EXPRESSION TAG	UNP Q168G2
H	12	GLY	-	EXPRESSION TAG	UNP Q168G2
H	13	SER	-	EXPRESSION TAG	UNP Q168G2
H	14	SER	-	EXPRESSION TAG	UNP Q168G2
H	15	HIS	-	EXPRESSION TAG	UNP Q168G2
H	16	HIS	-	EXPRESSION TAG	UNP Q168G2
H	17	HIS	-	EXPRESSION TAG	UNP Q168G2
H	18	HIS	-	EXPRESSION TAG	UNP Q168G2
H	19	HIS	-	EXPRESSION TAG	UNP Q168G2
H	20	HIS	-	EXPRESSION TAG	UNP Q168G2
H	21	SER	-	EXPRESSION TAG	UNP Q168G2
H	22	SER	-	EXPRESSION TAG	UNP Q168G2
H	23	GLY	-	EXPRESSION TAG	UNP Q168G2
H	24	LEU	-	EXPRESSION TAG	UNP Q168G2
H	25	VAL	-	EXPRESSION TAG	UNP Q168G2
H	26	PRO	-	EXPRESSION TAG	UNP Q168G2
H	27	ARG	-	EXPRESSION TAG	UNP Q168G2
H	28	GLY	-	EXPRESSION TAG	UNP Q168G2
H	29	SER	-	EXPRESSION TAG	UNP Q168G2
H	30	HIS	-	EXPRESSION TAG	UNP Q168G2
H	31	MET	-	EXPRESSION TAG	UNP Q168G2
J	11	MET	-	EXPRESSION TAG	UNP Q168G2


*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
J	12	GLY	-	EXPRESSION TAG	UNP Q168G2
J	13	SER	-	EXPRESSION TAG	UNP Q168G2
J	14	SER	-	EXPRESSION TAG	UNP Q168G2
J	15	HIS	-	EXPRESSION TAG	UNP Q168G2
J	16	HIS	-	EXPRESSION TAG	UNP Q168G2
J	17	HIS	-	EXPRESSION TAG	UNP Q168G2
J	18	HIS	-	EXPRESSION TAG	UNP Q168G2
J	19	HIS	-	EXPRESSION TAG	UNP Q168G2
J	20	HIS	-	EXPRESSION TAG	UNP Q168G2
J	21	SER	-	EXPRESSION TAG	UNP Q168G2
J	22	SER	-	EXPRESSION TAG	UNP Q168G2
J	23	GLY	-	EXPRESSION TAG	UNP Q168G2
J	24	LEU	-	EXPRESSION TAG	UNP Q168G2
J	25	VAL	-	EXPRESSION TAG	UNP Q168G2
J	26	PRO	-	EXPRESSION TAG	UNP Q168G2
J	27	ARG	-	EXPRESSION TAG	UNP Q168G2
J	28	GLY	-	EXPRESSION TAG	UNP Q168G2
J	29	SER	-	EXPRESSION TAG	UNP Q168G2
J	30	HIS	-	EXPRESSION TAG	UNP Q168G2
J	31	MET	-	EXPRESSION TAG	UNP Q168G2
L	11	MET	-	EXPRESSION TAG	UNP Q168G2
L	12	GLY	-	EXPRESSION TAG	UNP Q168G2
L	13	SER	-	EXPRESSION TAG	UNP Q168G2
L	14	SER	-	EXPRESSION TAG	UNP Q168G2
L	15	HIS	-	EXPRESSION TAG	UNP Q168G2
L	16	HIS	-	EXPRESSION TAG	UNP Q168G2
L	17	HIS	-	EXPRESSION TAG	UNP Q168G2
L	18	HIS	-	EXPRESSION TAG	UNP Q168G2
L	19	HIS	-	EXPRESSION TAG	UNP Q168G2
L	20	HIS	-	EXPRESSION TAG	UNP Q168G2
L	21	SER	-	EXPRESSION TAG	UNP Q168G2
L	22	SER	-	EXPRESSION TAG	UNP Q168G2
L	23	GLY	-	EXPRESSION TAG	UNP Q168G2
L	24	LEU	-	EXPRESSION TAG	UNP Q168G2
L	25	VAL	-	EXPRESSION TAG	UNP Q168G2
L	26	PRO	-	EXPRESSION TAG	UNP Q168G2
L	27	ARG	-	EXPRESSION TAG	UNP Q168G2
L	28	GLY	-	EXPRESSION TAG	UNP Q168G2
L	29	SER	-	EXPRESSION TAG	UNP Q168G2
L	30	HIS	-	EXPRESSION TAG	UNP Q168G2
L	31	MET	-	EXPRESSION TAG	UNP Q168G2

- Molecule 3 is 5-(HEXAHYDRO-2-OXO-1H-THIENO[3,4-D]IMIDAZOL-6-YL)PENTANAL

(three-letter code: BTI) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	10	2	2	1		
3	C	1	Total	C	N	O	S	0	0
			15	10	2	2	1		
3	E	1	Total	C	N	O	S	0	0
			15	10	2	2	1		
3	G	1	Total	C	N	O	S	0	0
			15	10	2	2	1		
3	I	1	Total	C	N	O	S	0	0
			15	10	2	2	1		
3	K	1	Total	C	N	O	S	0	0
			15	10	2	2	1		




### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Propionyl-CoA carboxylase, alpha subunit





- Molecule 1: Propionyl-CoA carboxylase, alpha subunit



● Molecule 1: Propionyl-CoA carboxylase, alpha subunit








• Molecule 1: Propionyl-CoA carboxylase, alpha subunit









• Molecule 2: Propionyl-CoA carboxylase, beta subunit


Chain D: 49% 42% 5%



• Molecule 2: Propionyl-CoA carboxylase, beta subunit


Chain F: 51% 40% 5%






• Molecule 2: Propionyl-CoA carboxylase, beta subunit


Chain H: 51% 40% 5%



• Molecule 2: Propionyl-CoA carboxylase, beta subunit


Chain J: 51% 39% 5%





• Molecule 2: Propionyl-CoA carboxylase, beta subunit

Chain L: 51% 40% 5%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	133.89Å 159.17Å 153.74Å 113.87° 101.03° 108.99°	Depositor
Resolution (Å)	29.36 – 3.20 29.36 – 3.14	Depositor EDS
% Data completeness (in resolution range)	92.4 (29.36-3.20) 90.6 (29.36-3.14)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 3.11Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.212 , 0.245 0.206 , 0.240	Depositor DCC
$R_{free}$ test set	12641 reflections (7.45%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.8	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 51.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for -h,h+k+l,-l	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	51921	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BTI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/4586	0.73	1/6209 (0.0%)
1	C	0.51	0/4586	0.74	1/6209 (0.0%)
1	E	0.52	0/4586	0.74	1/6209 (0.0%)
1	G	0.52	0/5036	0.74	1/6811 (0.0%)
1	I	0.51	0/5036	0.74	1/6811 (0.0%)
1	K	0.51	0/5036	0.73	1/6811 (0.0%)
2	B	0.62	1/3990 (0.0%)	0.81	0/5399
2	D	0.61	1/3990 (0.0%)	0.82	0/5399
2	F	0.61	1/3990 (0.0%)	0.82	0/5399
2	H	0.61	0/3990	0.80	0/5399
2	J	0.61	0/3990	0.82	0/5399
2	L	0.61	0/3990	0.81	0/5399
All	All	0.56	3/52806 (0.0%)	0.77	6/71454 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	E	0	1
1	G	0	1
1	I	0	1
1	K	0	1
All	All	0	5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	365	CYS	CB-SG	-6.58	1.71	1.82

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	365	CYS	CB-SG	-5.84	1.72	1.81
2	F	365	CYS	CB-SG	-5.36	1.73	1.81

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	422(B)	ALA	N-CA-C	5.95	127.07	111.00
1	K	422(B)	ALA	N-CA-C	5.46	125.75	111.00
1	G	422(B)	ALA	N-CA-C	5.44	125.69	111.00
1	I	422(B)	ALA	N-CA-C	5.36	125.48	111.00
1	E	422(B)	ALA	N-CA-C	5.19	125.01	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	420	TYR	Sidechain
1	E	420	TYR	Sidechain
1	G	420	TYR	Sidechain
1	I	420	TYR	Sidechain
1	K	420	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4507	0	4511	424	0
1	C	4507	0	4512	449	0
1	E	4507	0	4512	449	0
1	G	4950	0	4944	491	0
1	I	4950	0	4943	486	0
1	K	4950	0	4943	474	0
2	B	3910	0	3851	260	0
2	D	3910	0	3851	251	0
2	F	3910	0	3851	229	0
2	H	3910	0	3851	239	0
2	J	3910	0	3851	233	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	L	3910	0	3851	230	0
3	A	15	0	15	1	0
3	C	15	0	15	0	0
3	E	15	0	15	1	0
3	G	15	0	15	0	0
3	I	15	0	15	2	0
3	K	15	0	15	0	0
All	All	51921	0	51561	3984	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 3984 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:338:ILE:HD11	1:E:348:LEU:HB2	1.25	1.17
1:A:338:ILE:HD11	1:A:348:LEU:HB2	1.26	1.16
1:C:115:PRO:HG2	1:C:116:PRO:HD3	1.29	1.14
1:A:433:THR:HG22	1:A:435:VAL:H	1.06	1.11
1:G:400:LEU:HD13	1:G:449:ILE:HD11	1.33	1.11

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	585/681 (86%)	451 (77%)	99 (17%)	35 (6%)	1	12
1	C	585/681 (86%)	456 (78%)	94 (16%)	35 (6%)	1	12
1	E	585/681 (86%)	449 (77%)	98 (17%)	38 (6%)	1	10
1	G	638/681 (94%)	502 (79%)	100 (16%)	36 (6%)	2	14

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	638/681 (94%)	504 (79%)	94 (15%)	40 (6%)	1	10
1	K	638/681 (94%)	505 (79%)	92 (14%)	41 (6%)	1	10
2	B	504/531 (95%)	444 (88%)	49 (10%)	11 (2%)	6	35
2	D	504/531 (95%)	445 (88%)	47 (9%)	12 (2%)	6	34
2	F	504/531 (95%)	451 (90%)	43 (8%)	10 (2%)	7	38
2	H	504/531 (95%)	451 (90%)	43 (8%)	10 (2%)	7	38
2	J	504/531 (95%)	452 (90%)	43 (8%)	9 (2%)	8	41
2	L	504/531 (95%)	455 (90%)	39 (8%)	10 (2%)	7	38
All	All	6693/7272 (92%)	5565 (83%)	841 (13%)	287 (4%)	2	20

5 of 287 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100	ALA
1	A	137	GLN
1	A	138	ALA
1	A	277	ASP
1	A	331	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	474/540 (88%)	417 (88%)	57 (12%)	5	22
1	C	474/540 (88%)	420 (89%)	54 (11%)	5	24
1	E	474/540 (88%)	418 (88%)	56 (12%)	5	23
1	G	520/540 (96%)	461 (89%)	59 (11%)	6	25
1	I	520/540 (96%)	461 (89%)	59 (11%)	6	25
1	K	520/540 (96%)	460 (88%)	60 (12%)	5	24
2	B	415/437 (95%)	392 (94%)	23 (6%)	21	57
2	D	415/437 (95%)	392 (94%)	23 (6%)	21	57

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	415/437 (95%)	394 (95%)	21 (5%)	24	60
2	H	415/437 (95%)	390 (94%)	25 (6%)	19	54
2	J	415/437 (95%)	385 (93%)	30 (7%)	14	47
2	L	415/437 (95%)	392 (94%)	23 (6%)	21	57
All	All	5472/5862 (93%)	4982 (91%)	490 (9%)	9	34

5 of 490 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	499	ARG
1	K	548	ASP
1	G	666	MET
1	K	473	PHE
2	L	238	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 124 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	167	GLN
1	K	342	GLN
1	G	564	GLN
1	K	188	ASN
2	L	151	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	BTI	C	801	1	15,16,16	1.72	1 (6%)	20,21,21	2.10	6 (30%)
3	BTI	E	801	1	15,16,16	1.76	1 (6%)	20,21,21	2.22	7 (35%)
3	BTI	G	801	1	15,16,16	1.86	1 (6%)	20,21,21	2.33	7 (35%)
3	BTI	I	801	1	15,16,16	1.77	1 (6%)	20,21,21	2.11	6 (30%)
3	BTI	K	801	1	15,16,16	1.76	1 (6%)	20,21,21	2.08	5 (25%)
3	BTI	A	801	1	15,16,16	1.72	1 (6%)	20,21,21	2.05	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BTI	C	801	1	-	0/6/27/27	0/2/2/2
3	BTI	E	801	1	-	1/6/27/27	0/2/2/2
3	BTI	G	801	1	-	5/6/27/27	0/2/2/2
3	BTI	I	801	1	-	3/6/27/27	0/2/2/2
3	BTI	K	801	1	-	3/6/27/27	0/2/2/2
3	BTI	A	801	1	-	2/6/27/27	0/2/2/2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	801	BTI	O3-C3	6.63	1.37	1.23
3	I	801	BTI	O3-C3	6.37	1.36	1.23
3	E	801	BTI	O3-C3	6.36	1.36	1.23
3	K	801	BTI	O3-C3	6.31	1.36	1.23
3	C	801	BTI	O3-C3	6.22	1.36	1.23

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	801	BTI	C6-C5-N3	-4.53	107.34	113.18
3	E	801	BTI	C6-C5-N3	-4.24	107.72	113.18
3	G	801	BTI	C4-N2-C3	-4.09	107.49	112.56
3	I	801	BTI	C6-S1-C2	4.03	98.37	89.98
3	E	801	BTI	C4-N2-C3	-3.98	107.62	112.56

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	801	BTI	S1-C2-C7-C8
3	G	801	BTI	C4-C2-C7-C8
3	I	801	BTI	C9-C10-C11-O11
3	K	801	BTI	C2-C7-C8-C9
3	A	801	BTI	C2-C7-C8-C9

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	801	BTI	1	0
3	I	801	BTI	2	0
3	A	801	BTI	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	591/681 (86%)	-0.31	5 (0%) 86 78	17, 65, 113, 128	0
1	C	591/681 (86%)	-0.26	4 (0%) 87 81	17, 65, 113, 128	0
1	E	591/681 (86%)	-0.22	11 (1%) 66 53	17, 66, 113, 128	0
1	G	646/681 (94%)	-0.28	4 (0%) 89 83	15, 67, 112, 128	0
1	I	646/681 (94%)	-0.22	9 (1%) 75 63	16, 67, 112, 127	0
1	K	646/681 (94%)	-0.24	11 (1%) 70 57	16, 68, 112, 128	0
2	B	506/531 (95%)	-0.87	4 (0%) 86 78	10, 26, 57, 101	0
2	D	506/531 (95%)	-0.86	2 (0%) 92 89	11, 26, 58, 101	0
2	F	506/531 (95%)	-0.87	2 (0%) 92 89	12, 27, 58, 100	0
2	H	506/531 (95%)	-0.89	1 (0%) 95 94	10, 26, 58, 101	0
2	J	506/531 (95%)	-0.87	1 (0%) 95 94	11, 26, 58, 100	0
2	L	506/531 (95%)	-0.88	1 (0%) 95 94	11, 26, 57, 100	0
All	All	6747/7272 (92%)	-0.53	55 (0%) 86 78	10, 43, 108, 128	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	329	TYR	3.9
1	E	95	ASP	3.6
2	F	476(A)	LEU	3.5
1	K	159	GLU	3.5
2	D	476(A)	LEU	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BTI	I	801	15/15	0.91	0.23	57,65,68,69	0
3	BTI	K	801	15/15	0.93	0.16	39,50,57,62	0
3	BTI	G	801	15/15	0.94	0.20	43,49,56,60	0
3	BTI	A	801	15/15	0.94	0.16	30,42,56,62	0
3	BTI	E	801	15/15	0.94	0.18	31,39,49,50	0
3	BTI	C	801	15/15	0.95	0.13	30,38,46,47	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.