



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 05:05 AM EDT

PDB ID : 5N9F
Title : Crystal Structure of Drosophila DHX36 helicase in complex with ssDNA CpG_A
Authors : Chen, W.-F.; Rety, S.; Hai-Lei Guo, H.-L.; Wu, W.-Q.; Liu, N.-N.; Liu, Q.-W.; Dai, Y.-X.; Xi, X.-G.
Deposited on : 2017-02-24
Resolution : 2.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	944	<div> <div></div> <div>66%</div> <div>24%</div> <div>• 9%</div> </div>
1	B	944	<div> <div></div> <div>65%</div> <div>25%</div> <div>• 9%</div> </div>
2	C	10	<div> <div>30%</div> <div>50%</div> <div>50%</div> </div>
2	D	10	<div> <div>40%</div> <div>60%</div> <div>40%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CG9323, isoform A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	859	Total	C	N	O	S	0	0	0
			6900	4357	1219	1279	45			
1	B	855	Total	C	N	O	S	0	0	0
			6844	4322	1206	1271	45			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	943	VAL	-	expression tag	UNP Q8SWT2
A	944	ASP	-	expression tag	UNP Q8SWT2
B	943	VAL	-	expression tag	UNP Q8SWT2
B	944	ASP	-	expression tag	UNP Q8SWT2

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*GP*GP*AP*CP*GP*AP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	P	0	0	0
			192	89	40	54	9			
2	D	10	Total	C	N	O	P	0	0	0
			195	89	40	56	10			


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		


- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Chain A: 

Residue	State	Percentage
MET	66%	
GLN	66%	
ARG	66%	
ASP	66%	
GLN	66%	
ASP	66%	
SER	66%	
ASP	66%	
GLY	66%	
SER	66%	
ASN	66%	
ALA	66%	
ARG	66%	
LYS	66%	
GLY	66%	
ASN	66%	
ARG	66%	
PRO	66%	
PRO	66%	
GLY	66%	
LEU	66%	
ARG	66%	
GLY	66%	
LYS	66%	
ASP	66%	
ILE	66%	
GLY	66%	
LEU	66%	
TYR	66%	
TYR	66%	
ASN	66%	
LEU	66%	
ALA	66%	
ARG	66%	
GLN	66%	
GLN	66%	
LYS	66%	
LYS	66%	
ASP	66%	
ARG	66%	
GLY	66%	
GLU	66%	
ASN	66%	
ALA	66%	
GLU	66%	
SER	66%	
LYS	66%	
GLU	66%	
PRO	66%	
GLN	66%	
152	24%	
157	24%	
160	24%	
165	24%	
175	24%	
179	24%	
180	24%	
181	24%	
182	24%	
183	24%	
184	24%	
185	24%	
186	24%	
187	24%	
188	24%	
189	24%	
190	24%	
191	24%	
192	24%	
193	24%	
194	24%	
195	24%	
196	24%	
197	24%	
198	24%	
199	24%	
200	24%	
201	24%	
202	24%	
203	24%	
204	24%	
205	24%	
206	24%	
207	24%	
208	24%	
209	24%	
210	24%	
211	24%	
212	24%	
213	24%	
214	24%	
215	24%	
216	24%	
217	24%	
218	24%	
219	24%	
220	24%	
221	24%	
222	24%	
223	24%	
224	24%	
225	24%	
226	24%	
227	24%	
228	24%	
229	24%	
230	24%	
231	24%	
232	24%	
233	24%	
234	24%	
235	24%	
236	24%	
237	24%	
238	24%	
239	24%	
240	24%	
241	24%	
242	24%	
243	24%	
244	24%	
245	24%	
246	24%	
247	24%	
248	24%	
249	24%	
250	24%	
251	24%	
252	24%	
253	24%	
254	24%	
255	24%	
256	24%	
257	24%	
258	24%	
259	24%	
260	24%	
261	24%	
262	24%	
263	24%	
264	24%	
265	24%	
266	24%	
267	24%	
268	24%	
269	24%	
270	24%	
271	24%	
272	24%	
273	24%	
274	24%	
275	24%	
276	24%	
277	24%	
278	24%	
279	24%	
280	24%	
281	24%	
282	24%	
283	24%	
284	24%	
285	24%	
286	24%	
287	24%	
288	24%	
289	24%	
290	24%	



- Chain B: 



- Molecule 2: DNA (5'-D(*GP*GP*GP*GP*AP*CP*GP*AP*TP*C)-3')



- Molecule 2: DNA (5'-D(*GP*GP*GP*GP*AP*CP*GP*AP*TP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	306.68Å 51.31Å 164.86Å 90.00° 115.55° 90.00°	Depositor
Resolution (Å)	67.10 – 2.97 138.35 – 2.97	Depositor EDS
% Data completeness (in resolution range)	95.4 (67.10-2.97) 95.4 (138.35-2.97)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.96Å)	Xtriage
Refinement program	PHENIX (1.11_2563: ???)	Depositor
R, R_{free}	0.214 , 0.270 0.215 , 0.269	Depositor DCC
R_{free} test set	2301 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	68.1	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14133	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/7022	0.47	0/9474
1	B	0.27	0/6964	0.46	0/9398
2	C	0.56	0/216	0.81	0/334
2	D	0.48	0/219	0.82	0/338
All	All	0.28	0/14421	0.48	0/19544

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	142	ASN	Peptide
1	B	142	ASN	Peptide
1	B	862	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6900	0	7008	178	0
1	B	6844	0	6926	181	0
2	C	192	0	101	6	0
2	D	195	0	100	4	0
3	B	1	0	0	0	0
4	B	1	0	0	1	0
All	All	14133	0	14135	359	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 359 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:175:VAL:HG11	1:B:315:GLU:HG2	1.42	0.98
1:A:147:LEU:HD21	1:A:151:LYS:HE3	1.55	0.89
1:B:461:PRO:HG3	1:B:673:LEU:HD11	1.57	0.86
1:A:529:VAL:HG13	1:A:533:ASN:HB2	1.60	0.84
1:A:891:ASN:HA	1:A:894:LYS:HG2	1.61	0.82

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	853/944 (90%)	820 (96%)	30 (4%)	3 (0%)	34 69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	849/944 (90%)	815 (96%)	30 (4%)	4 (0%)	29 64
All	All	1702/1888 (90%)	1635 (96%)	60 (4%)	7 (0%)	34 69

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	144	LYS
1	B	143	ALA
1	B	862	GLN
1	A	791	ARG
1	B	792	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	767/842 (91%)	747 (97%)	20 (3%)	46 75
1	B	757/842 (90%)	740 (98%)	17 (2%)	52 79
All	All	1524/1684 (90%)	1487 (98%)	37 (2%)	49 77

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	614	ARG
1	B	829	GLN
1	B	640	MET
1	B	783	ARG
1	A	509	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	520	ASN
1	B	467	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	94	GLN
1	B	463	HIS
1	A	781	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	859/944 (90%)	0.24	14 (1%) 72 55	33, 67, 119, 185	0
1	B	855/944 (90%)	0.14	9 (1%) 80 65	36, 69, 129, 191	0
2	C	10/10 (100%)	1.20	3 (30%) 0 0	81, 126, 156, 182	0
2	D	10/10 (100%)	1.99	4 (40%) 0 0	86, 128, 172, 194	0
All	All	1734/1908 (90%)	0.20	30 (1%) 70 53	33, 68, 128, 194	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	791	ARG	8.7
1	A	787	GLN	7.9
1	A	788	ILE	6.1
2	D	10	DT	4.5
2	D	11	DC	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CL	B	1002	1/1	0.48	0.27	94,94,94,94	0
3	NA	B	1001	1/1	0.90	0.31	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.