



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 17, 2024 – 04:50 AM EST

PDB ID : 4NE9
Title : PCSK9 in complex with LDLR peptide
Authors : Liu, S.
Deposited on : 2013-10-28
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	<div> <div>7%</div> <div> <div></div> <div>77%</div> <div>12%</div> <div>•</div> <div>9%</div> </div> </div>
1	B	540	<div> <div>9%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>•</div> <div>9%</div> </div> </div>
2	C	152	<div> <div></div> <div> <div>52%</div> <div>7%</div> <div>•</div> <div>39%</div> </div> </div>
2	P	152	<div> <div>%</div> <div> <div></div> <div>54%</div> <div>7%</div> <div>39%</div> </div> </div>
3	D	26	<div> <div></div> <div> <div>92%</div> <div>8%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	E	2	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8999 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	491	Total	C	N	O	S	0	1	0
			3646	2246	676	692	32			
1	B	490	Total	C	N	O	S	0	1	0
			3623	2230	669	692	32			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	474	ILE	VAL	engineered mutation	UNP Q8NBP7
A	670	GLU	GLY	engineered mutation	UNP Q8NBP7
B	474	ILE	VAL	engineered mutation	UNP Q8NBP7
B	670	GLU	GLY	engineered mutation	UNP Q8NBP7

- Molecule 2 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	92	Total	C	N	O	S	0	0	0
			740	474	133	131	2			
2	P	92	Total	C	N	O	S	0	0	0
			740	474	133	131	2			

- Molecule 3 is a protein called Low-density lipoprotein receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	26	Total	C	N	O	S	0	0	0
			189	111	33	41	4			

- Molecule 4 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	1	Total	Ca	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	9	Total	O	0	0
			9	9		
7	B	5	Total	O	0	0
			5	5		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	4	Total	O	0	0
			4	4		
7	P	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proprotein convertase subtilisin/kexin type 9



- Molecule 1: Proprotein convertase subtilisin/kexin type 9



- Molecule 2: Proprotein convertase subtilisin/kexin type 9

Topic	Count
T61	10
H65	10
R66	10
D70	10
V79	10
R96	10
R97	10
A100	10
Q101	10
R104	10
L108	10
L118	10
L123	10
L133	10
Q152	10

-

-
- A diagram showing a sequence of four colored blocks: G1 (green), T2 (yellow), V15 (yellow), and L26 (green). The blocks are arranged horizontally and connected by thin lines.

- | | NAG1 | FUC2 |
|-----|------|------|
| 1 | 0.00 | 0.00 |
| 2 | 0.00 | 0.00 |
| 3 | 0.00 | 0.00 |
| 4 | 0.00 | 0.00 |
| 5 | 0.00 | 0.00 |
| 6 | 0.00 | 0.00 |
| 7 | 0.00 | 0.00 |
| 8 | 0.00 | 0.00 |
| 9 | 0.00 | 0.00 |
| 10 | 0.00 | 0.00 |
| 11 | 0.00 | 0.00 |
| 12 | 0.00 | 0.00 |
| 13 | 0.00 | 0.00 |
| 14 | 0.00 | 0.00 |
| 15 | 0.00 | 0.00 |
| 16 | 0.00 | 0.00 |
| 17 | 0.00 | 0.00 |
| 18 | 0.00 | 0.00 |
| 19 | 0.00 | 0.00 |
| 20 | 0.00 | 0.00 |
| 21 | 0.00 | 0.00 |
| 22 | 0.00 | 0.00 |
| 23 | 0.00 | 0.00 |
| 24 | 0.00 | 0.00 |
| 25 | 0.00 | 0.00 |
| 26 | 0.00 | 0.00 |
| 27 | 0.00 | 0.00 |
| 28 | 0.00 | 0.00 |
| 29 | 0.00 | 0.00 |
| 30 | 0.00 | 0.00 |
| 31 | 0.00 | 0.00 |
| 32 | 0.00 | 0.00 |
| 33 | 0.00 | 0.00 |
| 34 | 0.00 | 0.00 |
| 35 | 0.00 | 0.00 |
| 36 | 0.00 | 0.00 |
| 37 | 0.00 | 0.00 |
| 38 | 0.00 | 0.00 |
| 39 | 0.00 | 0.00 |
| 40 | 0.00 | 0.00 |
| 41 | 0.00 | 0.00 |
| 42 | 0.00 | 0.00 |
| 43 | 0.00 | 0.00 |
| 44 | 0.00 | 0.00 |
| 45 | 0.00 | 0.00 |
| 46 | 0.00 | 0.00 |
| 47 | 0.00 | 0.00 |
| 48 | 0.00 | 0.00 |
| 49 | 0.00 | 0.00 |
| 50 | 0.00 | 0.00 |
| 51 | 0.00 | 0.00 |
| 52 | 0.00 | 0.00 |
| 53 | 0.00 | 0.00 |
| 54 | 0.00 | 0.00 |
| 55 | 0.00 | 0.00 |
| 56 | 0.00 | 0.00 |
| 57 | 0.00 | 0.00 |
| 58 | 0.00 | 0.00 |
| 59 | 0.00 | 0.00 |
| 60 | 0.00 | 0.00 |
| 61 | 0.00 | 0.00 |
| 62 | 0.00 | 0.00 |
| 63 | 0.00 | 0.00 |
| 64 | 0.00 | 0.00 |
| 65 | 0.00 | 0.00 |
| 66 | 0.00 | 0.00 |
| 67 | 0.00 | 0.00 |
| 68 | 0.00 | 0.00 |
| 69 | 0.00 | 0.00 |
| 70 | 0.00 | 0.00 |
| 71 | 0.00 | 0.00 |
| 72 | 0.00 | 0.00 |
| 73 | 0.00 | 0.00 |
| 74 | 0.00 | 0.00 |
| 75 | 0.00 | 0.00 |
| 76 | 0.00 | 0.00 |
| 77 | 0.00 | 0.00 |
| 78 | 0.00 | 0.00 |
| 79 | 0.00 | 0.00 |
| 80 | 0.00 | 0.00 |
| 81 | 0.00 | 0.00 |
| 82 | 0.00 | 0.00 |
| 83 | 0.00 | 0.00 |
| 84 | 0.00 | 0.00 |
| 85 | 0.00 | 0.00 |
| 86 | 0.00 | 0.00 |
| 87 | 0.00 | 0.00 |
| 88 | 0.00 | 0.00 |
| 89 | 0.00 | 0.00 |
| 90 | 0.00 | 0.00 |
| 91 | 0.00 | 0.00 |
| 92 | 0.00 | 0.00 |
| 93 | 0.00 | 0.00 |
| 94 | 0.00 | 0.00 |
| 95 | 0.00 | 0.00 |
| 96 | 0.00 | 0.00 |
| 97 | 0.00 | 0.00 |
| 98 | 0.00 | 0.00 |
| 99 | 0.00 | 0.00 |
| 100 | 0.00 | 0.00 |

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	126.40Å 131.39Å 134.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.02 – 2.60 30.02 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.2 (30.02-2.60) 95.1 (30.02-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.61Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.190 , 0.211 0.198 , 0.220	Depositor DCC
R_{free} test set	3329 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	57.7	Xtriage
Anisotropy	0.335	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.016 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8999	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/3718	0.69	0/5048
1	B	0.51	0/3693	0.70	0/5014
2	C	0.51	0/757	0.73	0/1023
2	P	0.54	0/757	0.71	0/1023
3	D	0.56	0/190	0.70	0/255
All	All	0.51	0/9115	0.70	0/12363

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3646	0	3559	34	0
1	B	3623	0	3531	36	0
2	C	740	0	750	4	0
2	P	740	0	750	6	0
3	D	189	0	168	1	0
4	E	24	0	22	0	0
5	B	14	0	13	0	0
6	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	9	0	0	0	0
7	B	5	0	0	0	0
7	C	4	0	0	0	0
7	P	4	0	0	0	0
All	All	8999	0	8793	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:558:VAL:HG21	1:A:606:LEU:HD13	1.38	1.04
1:A:203:THR:O	1:A:204:ASP:HB2	1.57	1.02
1:B:203:THR:O	1:B:204:ASP:HB2	1.57	1.02
1:A:558:VAL:CG2	1:A:606:LEU:HD13	1.91	1.00
1:B:344:GLN:HE22	1:B:425:ASN:H	1.12	0.88
1:A:344:GLN:HE22	1:A:425:ASN:H	1.16	0.87
1:A:186:ASP:OD1	1:A:288:PRO:HG2	1.93	0.68
1:B:278:GLN:HE22	2:P:88:LEU:H	1.41	0.67
1:A:548:THR:HG22	1:A:596:ILE:HG22	1.77	0.67
1:A:641:THR:HG22	1:A:643:HIS:CE1	2.30	0.67
1:B:186:ASP:OD1	1:B:288:PRO:HG2	1.95	0.66
1:B:548:THR:HG22	1:B:596:ILE:HG22	1.79	0.65
1:B:641:THR:HG22	1:B:643:HIS:CE1	2.31	0.64
1:A:518:GLU:H	1:A:518:GLU:CD	2.02	0.63
1:A:212:ASP:H	1:A:256:GLN:HE22	1.46	0.62
1:B:212:ASP:H	1:B:256:GLN:HE22	1.46	0.62
1:B:518:GLU:H	1:B:518:GLU:CD	2.03	0.62
1:B:539:ALA:HB2	1:B:550:VAL:HG13	1.83	0.60
1:A:203:THR:O	1:A:204:ASP:CB	2.38	0.59
1:A:539:ALA:HB2	1:A:550:VAL:HG13	1.85	0.58
1:B:277:VAL:HG13	2:P:117:GLY:HA2	1.85	0.58
2:C:101:GLN:HE21	2:C:133:LEU:HD11	1.72	0.55
1:A:632:LEU:H	1:A:652:ASN:ND2	2.03	0.55
1:A:185:LEU:HD11	1:A:271:ILE:HD11	1.88	0.54
1:B:311:LEU:HB2	1:B:333:VAL:HG12	1.89	0.54
1:B:549:ARG:HG2	1:B:589:VAL:HG22	1.89	0.54
1:A:529:LEU:HG	1:A:532:ALA:HB2	1.89	0.54
1:B:185:LEU:HD11	1:B:271:ILE:HD11	1.89	0.54
1:B:632:LEU:H	1:B:652:ASN:ND2	2.06	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:549:ARG:HG2	1:A:589:VAL:HG22	1.90	0.54
1:A:476:ARG:HH21	1:A:503:GLN:NE2	2.07	0.53
1:A:237:ARG:O	1:A:243:LYS:HD2	2.09	0.52
1:A:200:VAL:HG22	1:A:247:MET:HB2	1.92	0.52
1:A:295:ARG:HH11	2:P:65:HIS:CE1	2.28	0.52
1:B:212:ASP:H	1:B:256:GLN:NE2	2.07	0.51
1:B:638:LEU:HB2	1:B:673:THR:HB	1.93	0.51
1:B:476:ARG:HH21	1:B:503:GLN:NE2	2.09	0.51
1:B:203:THR:O	1:B:204:ASP:CB	2.38	0.50
1:B:550:VAL:HG22	1:B:596:ILE:HG23	1.94	0.49
1:B:278:GLN:NE2	2:P:88:LEU:H	2.09	0.49
1:A:638:LEU:HB2	1:A:673:THR:HB	1.94	0.49
1:A:212:ASP:H	1:A:256:GLN:NE2	2.09	0.48
1:B:374:ASP:OD2	3:D:15:VAL:HG21	2.12	0.48
1:B:490:SER:HB2	1:B:520:VAL:HG12	1.94	0.48
1:A:557:HIS:CE1	1:A:602:HIS:HB2	2.49	0.48
1:B:615:ILE:HG13	1:B:616:PRO:HD2	1.95	0.48
1:B:361:LEU:HD22	1:B:442:ALA:HB2	1.95	0.47
1:B:200:VAL:HG22	1:B:247:MET:HB2	1.97	0.47
1:A:490:SER:HB2	1:A:520:VAL:HG12	1.94	0.47
1:A:361:LEU:HD22	1:A:442:ALA:HB2	1.96	0.47
1:A:550:VAL:HG22	1:A:596:ILE:HG23	1.96	0.47
2:C:79:VAL:HG22	2:C:123:LEU:HD13	1.97	0.46
1:B:551:HIS:HB3	1:B:587:GLN:HB2	1.97	0.46
1:A:551:HIS:HB3	1:A:587:GLN:HB2	1.97	0.46
1:A:303:ARG:HE	1:A:303:ARG:HB2	1.48	0.44
1:A:539:ALA:CB	1:A:550:VAL:HG13	2.48	0.44
1:A:277:VAL:HG12	1:A:278:GLN:HG2	1.99	0.43
2:C:100:ALA:O	2:C:104:ARG:HG2	2.18	0.43
1:B:558:VAL:HG21	1:B:608:CYS:SG	2.58	0.43
1:B:529:LEU:HA	1:B:530:PRO:HD2	1.66	0.43
1:B:237:ARG:O	1:B:243:LYS:HD2	2.18	0.43
1:A:591:HIS:CD2	1:A:593:GLU:H	2.37	0.42
1:B:591:HIS:CD2	1:B:593:GLU:H	2.38	0.42
1:A:632:LEU:H	1:A:652:ASN:HD22	1.67	0.42
1:B:277:VAL:HG13	2:P:117:GLY:CA	2.49	0.42
1:B:465:SER:HB3	1:B:473:ALA:HB2	2.01	0.41
1:B:539:ALA:CB	1:B:550:VAL:HG13	2.48	0.41
2:P:69:LYS:HD3	2:P:72:TRP:CE2	2.56	0.41
1:B:591:HIS:HD2	1:B:593:GLU:H	1.70	0.40
1:A:345:PRO:HD3	1:A:424:ILE:HG23	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:591:HIS:HD2	1:A:593:GLU:H	1.69	0.40
1:B:180:VAL:HG22	1:B:282:PRO:HG2	2.02	0.40
1:B:295:ARG:HH11	2:C:65:HIS:CE1	2.38	0.40
1:A:304:LEU:HD12	1:A:304:LEU:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	482/540 (89%)	458 (95%)	18 (4%)	6 (1%)	11	24
1	B	479/540 (89%)	453 (95%)	19 (4%)	7 (2%)	8	18
2	C	90/152 (59%)	86 (96%)	4 (4%)	0	100	100
2	P	90/152 (59%)	87 (97%)	3 (3%)	0	100	100
3	D	24/26 (92%)	23 (96%)	1 (4%)	0	100	100
All	All	1165/1410 (83%)	1107 (95%)	45 (4%)	13 (1%)	12	26

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	245	ALA
1	B	530	PRO
1	A	543	GLU
1	A	642	SER
1	B	204	ASP
1	B	447	SER
1	B	543	GLU
1	A	204	ASP
1	B	642	SER
1	A	618	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	618	PRO
1	A	280	VAL
1	B	617	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/431 (91%)	364 (93%)	27 (7%)	13	28
1	B	389/431 (90%)	361 (93%)	28 (7%)	12	26
2	C	79/127 (62%)	71 (90%)	8 (10%)	6	12
2	P	79/127 (62%)	74 (94%)	5 (6%)	15	32
3	D	22/22 (100%)	21 (96%)	1 (4%)	23	47
All	All	960/1138 (84%)	891 (93%)	69 (7%)	12	26

All (69) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	SER
1	A	179	LEU
1	A	190	GLN
1	A	197	GLU
1	A	215	ARG
1	A	222	LYS
1	A	303	ARG
1	A	361	LEU
1	A	375	CYS
1	A	411	LEU
1	A	413	GLN
1	A	422	ASP
1	A	454	GLN
1	A	507	LEU
1	A	518	GLU
1	A	550	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	555	GLN
1	A	567	GLU
1	A	570	ASP
1	A	587	GLN
1	A	592	ARG
1	A	606	LEU
1	A	641	THR
1	A	642	SER
1	A	651[A]	ASP
1	A	651[B]	ASP
1	A	659	ARG
1	B	178	SER
1	B	179	LEU
1	B	190	GLN
1	B	194	ARG
1	B	219	GLN
1	B	222	LYS
1	B	283	LEU
1	B	304	LEU
1	B	361	LEU
1	B	375	CYS
1	B	411	LEU
1	B	413	GLN
1	B	422	ASP
1	B	454	GLN
1	B	518	GLU
1	B	531	GLN
1	B	550	VAL
1	B	555	GLN
1	B	567	GLU
1	B	570	ASP
1	B	587	GLN
1	B	592	ARG
1	B	606	LEU
1	B	641	THR
1	B	642	SER
1	B	651[A]	ASP
1	B	651[B]	ASP
1	B	659	ARG
2	C	66	ARG
2	C	70	ASP
2	C	96	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	97	ARG
2	C	104	ARG
2	C	108	LEU
2	C	118	LEU
2	C	133	LEU
3	D	2	THR
2	P	66	ARG
2	P	70	ASP
2	P	104	ARG
2	P	108	LEU
2	P	118	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	256	GLN
1	A	278	GLN
1	A	298	ASN
1	A	344	GLN
1	A	413	GLN
1	A	464	HIS
1	A	513	ASN
1	A	587	GLN
1	A	591	HIS
1	A	602	HIS
1	A	643	HIS
1	A	652	ASN
1	B	256	GLN
1	B	278	GLN
1	B	298	ASN
1	B	344	GLN
1	B	413	GLN
1	B	464	HIS
1	B	513	ASN
1	B	537	HIS
1	B	554	GLN
1	B	587	GLN
1	B	591	HIS
1	B	643	HIS
1	B	652	ASN
2	C	65	HIS
2	C	101	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	P	65	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	E	1	1,4	14,14,15	0.34	0	17,19,21	0.55	0
4	FUC	E	2	4	10,10,11	0.46	0	14,14,16	0.80	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	FUC	E	2	4	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	2	FUC	C1-O5-C5	2.47	118.80	112.97


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	901	1	14,14,15	0.34	0	17,19,21	0.75	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	901	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	901	NAG	C1-O5-C5	2.72	115.83	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	491/540 (90%)	0.15	38 (7%) 21 17	34, 61, 127, 158	1 (0%)
1	B	490/540 (90%)	0.28	50 (10%) 13 11	36, 65, 139, 189	1 (0%)
2	C	92/152 (60%)	-0.31	0 100 100	41, 60, 87, 105	0
2	P	92/152 (60%)	-0.54	2 (2%) 62 57	35, 47, 70, 87	0
3	D	26/26 (100%)	-0.11	0 100 100	50, 61, 79, 92	0
All	All	1191/1410 (84%)	0.11	90 (7%) 21 17	34, 61, 127, 189	2 (0%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	178	SER	8.3
1	A	178	SER	7.8
1	B	585	PRO	7.3
1	B	168	ALA	7.0
1	A	168	ALA	6.8
1	B	176	GLY	6.7
1	B	177	GLY	6.1
1	B	220	ALA	5.9
1	A	585	PRO	5.8
1	B	572	GLY	5.6
1	B	682	ARG	5.4
1	A	682	ARG	5.2
1	A	572	GLY	5.1
1	B	449	HIS	5.0
1	B	618	PRO	5.0
1	A	179	LEU	4.7
1	B	448	THR	4.6
1	B	617	ALA	4.6
1	B	221	SER	4.5
1	A	515	PHE	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	659	ARG	4.1
1	A	219	GLN	4.1
1	B	212	ASP	4.0
1	A	544	ALA	3.9
1	B	544	ALA	3.9
1	B	537	HIS	3.9
1	B	515	PHE	3.8
1	A	568	VAL	3.8
1	B	640	GLY	3.7
1	B	166	TYR	3.6
1	B	153	SER	3.5
1	A	615	ILE	3.3
1	B	219	GLN	3.3
1	A	617	ALA	3.3
1	B	671	ALA	3.3
1	A	616	PRO	3.2
1	B	588	CYS	3.2
1	B	167	ARG	3.2
1	B	546	MET	3.2
1	B	179	LEU	3.1
1	B	447	SER	3.1
1	B	164	PRO	3.1
1	A	537	HIS	3.1
1	A	516	GLY	3.0
1	A	659	ARG	3.0
1	A	280	VAL	3.0
1	B	643	HIS	2.9
1	B	592	ARG	2.9
1	B	222	LYS	2.9
1	A	201	MET	2.8
1	B	496	ARG	2.8
2	P	116	HIS	2.7
1	A	640	GLY	2.7
1	B	470	MET	2.6
1	B	551	HIS	2.6
1	A	470	MET	2.6
1	A	554	GLN	2.6
1	A	542	ALA	2.6
1	A	546	MET	2.6
1	A	618	PRO	2.6
1	B	571	LEU	2.6
1	A	215	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	643	HIS	2.6
1	A	642	SER	2.5
1	B	223	CYS	2.5
1	B	589	VAL	2.4
1	A	454	GLN	2.4
1	B	454	GLN	2.4
1	B	507	LEU	2.4
1	A	589	VAL	2.4
1	A	445	PRO	2.4
1	A	571	LEU	2.3
1	B	620	GLU	2.3
1	B	499	ARG	2.3
1	B	531	GLN	2.3
1	A	592	ARG	2.3
1	B	446	PRO	2.2
2	P	96	ARG	2.2
1	B	639	PRO	2.2
1	B	591	HIS	2.2
1	A	222	LYS	2.1
1	A	163	PRO	2.1
1	B	159	GLU	2.1
1	B	568	VAL	2.1
1	B	619	GLN	2.0
1	A	166	TYR	2.0
1	A	549	ARG	2.0
1	B	642	SER	2.0
1	A	473	ALA	2.0
1	A	671	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	B	901	14/15	0.59	0.19	123,134,138,141	0
6	CA	D	101	1/1	0.99	0.03	74,74,74,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.