



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 23, 2025 – 12:11 PM EDT

PDB ID : 9OB6 / pdb\_00009ob6  
Title : CDK2/CyclinE bound to compound 21 with P-loop in the EE and CC conformations  
Authors : Baird, J.; Holliday, M.H.  
Deposited on : 2025-04-22  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	<div> <div>9%</div> <div>89%</div> <div>10%</div> </div>
2	B	285	<div> <div>17%</div> <div>86%</div> <div>9%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4778 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	P	S	0	7	0
			2448	1588	415	436	1	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP P24941
A	-1	HIS	-	expression tag	UNP P24941
A	0	MET	-	expression tag	UNP P24941


- Molecule 2 is a protein called G1/S-specific cyclin-E1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	274	Total	C	N	O	S	0	0	0
			2243	1458	366	402	17			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	94	GLY	-	expression tag	UNP P24864
B	95	SER	-	expression tag	UNP P24864

- Molecule 3 is (1R,3R)-3-{3-[2-(4-methoxyphenyl)acetamido]-1H-pyrazol-5-yl}cyclopentyl propan-2-ylcarbamate (CCD ID: A1CAI) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>4</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			29	21	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	46	Total	O	0	0
			46	46		
4	B	12	Total	O	0	0
			12	12		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cyclin-dependent kinase 2



#### • Molecule 2: G1/S-specific cyclin-E1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.10Å 101.10Å 151.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.01 – 2.00 84.01 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (84.01-2.00) 100.0 (84.01-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.227 , 0.256 0.230 , 0.260	Depositor DCC
$R_{free}$ test set	51545 reflections (3.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.3	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4778	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1CAI, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/2499	0.46	0/3390
2	B	0.23	0/2299	0.40	0/3120
All	All	0.23	0/4798	0.43	0/6510

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2448	0	2488	14	0
2	B	2243	0	2245	13	0
3	A	29	0	0	1	0
4	A	46	0	0	0	0
4	B	12	0	0	1	0
All	All	4778	0	4733	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12[A]:GLU:HG3	1:A:17[A]:VAL:HG22	1.80	0.64
1:A:34:LYS:HE3	1:A:75:LYS:HD2	1.85	0.58
1:A:22:ARG:HH12	1:A:24:LYS:HD2	1.68	0.58
2:B:145:ARG:HD3	2:B:289:VAL:HG22	1.86	0.57
2:B:260:ASN:HD22	2:B:261:ASP:H	1.53	0.57
2:B:123:LYS:HD2	2:B:244:LEU:HG	1.88	0.55
1:A:57:GLU:OE1	2:B:123:LYS:NZ	2.40	0.55
1:A:72:THR:HG22	1:A:73:GLU:HG3	1.89	0.54
2:B:272:GLN:HB3	2:B:358:HIS:HB3	1.91	0.53
2:B:277:GLN:HG2	2:B:343:SER:HB2	1.94	0.50
1:A:255:LEU:HG	1:A:259:GLY:HA3	1.94	0.49
1:A:81:GLU:O	3:A:301:A1CAI:N12	2.46	0.49
2:B:167:TYR:HB3	2:B:245:THR:HB	1.95	0.48
1:A:49:ILE:HG22	2:B:202:LEU:HD21	1.96	0.47
1:A:155:PRO:HD3	2:B:255:GLN:HB2	1.95	0.47
1:A:175:LEU:HA	1:A:235:ASP:HB2	1.95	0.47
1:A:38:ASP:OD1	1:A:38:ASP:N	2.37	0.45
2:B:229:MET:HA	2:B:232:MET:HE2	2.00	0.43
1:A:223:ASP:H	1:A:226:VAL:HG22	1.83	0.43
2:B:110:TRP:O	4:B:401:HOH:O	2.22	0.43
2:B:271:PRO:HB2	2:B:274:ILE:HG23	2.02	0.42
1:A:128:LEU:HD13	1:A:189:LEU:HD13	2.01	0.42
1:A:4:PHE:HD1	1:A:4:PHE:HA	1.82	0.41
2:B:227:LEU:HA	2:B:230:GLU:HG2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	302/301 (100%)	291 (96%)	10 (3%)	1 (0%)	37	35
2	B	272/285 (95%)	266 (98%)	4 (2%)	2 (1%)	19	14

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	574/586 (98%)	557 (97%)	14 (2%)	3 (0%)	25	21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	VAL
2	B	365	ASP
2	B	182	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/265 (100%)	260 (98%)	6 (2%)	45	49
2	B	247/255 (97%)	239 (97%)	8 (3%)	34	35
All	All	513/520 (99%)	499 (97%)	14 (3%)	38	42

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	THR
1	A	59	ASN
1	A	161	HIS
1	A	199	ARG
1	A	252	VAL
1	A	293	VAL
2	B	260	ASN
2	B	262	LEU
2	B	274	ILE
2	B	304	LEU
2	B	355	GLU
2	B	364	ARG
2	B	369	LEU
2	B	371	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
2	B	136	GLN
2	B	260	ASN
2	B	273	GLN
2	B	320	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	TPO	A	160	1	8,10,11	2.59	1 (12%)	10,14,16	1.30	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	160	1	-	0/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	160	TPO	P-OG1	7.15	1.72	1.59

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	TPO	OG1-P-O1P	-2.48	100.50	109.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	A1CAI	A	301	-	29,31,31	2.33	9 (31%)	39,42,42	2.39	17 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1CAI	A	301	-	-	7/22/31/31	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	A1CAI	C08-N09	5.04	1.46	1.35
3	A	301	A1CAI	C07-C08	5.03	1.62	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	A1CAI	C13-N12	4.37	1.40	1.34
3	A	301	A1CAI	C07-C06	4.07	1.58	1.51
3	A	301	A1CAI	O19-C20	3.76	1.41	1.35
3	A	301	A1CAI	C10-N09	3.67	1.48	1.40
3	A	301	A1CAI	C20-N21	3.07	1.42	1.34
3	A	301	A1CAI	C18-C17	2.20	1.57	1.52
3	A	301	A1CAI	C22-N21	2.17	1.51	1.47

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	A1CAI	C07-C08-N09	6.40	126.69	114.73
3	A	301	A1CAI	C26-C13-N12	-4.89	103.73	110.28
3	A	301	A1CAI	C14-C13-N12	4.24	126.49	120.23
3	A	301	A1CAI	C13-C26-C10	4.03	109.53	105.91
3	A	301	A1CAI	O19-C20-O25	-3.67	119.15	124.55
3	A	301	A1CAI	O19-C20-N21	3.67	114.94	110.26
3	A	301	A1CAI	O27-C08-C07	-3.38	114.67	121.99
3	A	301	A1CAI	C26-C10-N11	-2.98	106.21	110.47
3	A	301	A1CAI	C05-C06-C28	-2.98	113.81	118.23
3	A	301	A1CAI	O27-C08-N09	-2.79	118.64	123.64
3	A	301	A1CAI	C26-C10-N09	-2.70	127.39	131.11
3	A	301	A1CAI	C14-C18-C17	2.46	108.74	103.01
3	A	301	A1CAI	C16-C15-C14	2.44	107.76	103.39
3	A	301	A1CAI	C18-C14-C15	2.43	104.64	100.78
3	A	301	A1CAI	C07-C06-C05	2.40	124.42	120.89
3	A	301	A1CAI	C29-C28-C06	2.36	124.11	121.00
3	A	301	A1CAI	C04-C03-C29	-2.24	116.90	120.16

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	301	A1CAI	C04-C03-O02-C01
3	A	301	A1CAI	C29-C03-O02-C01
3	A	301	A1CAI	N21-C20-O19-C17
3	A	301	A1CAI	O25-C20-O19-C17
3	A	301	A1CAI	O19-C20-N21-C22
3	A	301	A1CAI	O25-C20-N21-C22
3	A	301	A1CAI	C16-C17-O19-C20

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	A1CAI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	297/301 (98%)	0.59	26 (8%) 17 16	31, 59, 106, 168	7 (2%)
2	B	274/285 (96%)	1.20	48 (17%) 5 4	55, 83, 125, 194	0
All	All	571/586 (97%)	0.88	74 (12%) 9 7	31, 71, 120, 194	7 (1%)

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	15[A]	TYR	14.1
1	A	14[A]	THR	12.9
1	A	154	VAL	6.8
1	A	16[A]	GLY	5.0
1	A	13[A]	GLY	4.2
1	A	155	PRO	4.2
2	B	267	LEU	4.2
2	B	102	GLY	4.2
2	B	265	VAL	4.0
2	B	256	VAL	3.8
1	A	158	THR	3.8
2	B	126	THR	3.6
2	B	367	LEU	3.5
2	B	375	ALA	3.5
2	B	270	TYR	3.4
2	B	262	LEU	3.4
1	A	298	LEU	3.4
2	B	117	TRP	3.4
2	B	103	SER	2.9
2	B	319	TYR	2.9
2	B	258	TYR	2.8
1	A	157	ARG	2.8
1	A	163	VAL	2.7
1	A	293	VAL	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	110	TRP	2.7
1	A	4	PHE	2.7
2	B	105	LEU	2.7
2	B	114	GLU	2.6
1	A	153	GLY	2.6
1	A	296	LEU	2.6
1	A	179	TYR	2.5
1	A	41	THR	2.5
1	A	294	PRO	2.5
2	B	373	ALA	2.5
2	B	268	PRO	2.5
2	B	187	LEU	2.5
2	B	266	LEU	2.5
2	B	263	HIS	2.4
2	B	259	LEU	2.4
1	A	162	GLU	2.4
1	A	1	MET	2.4
2	B	188	LEU	2.4
2	B	257	ALA	2.4
1	A	289	VAL	2.4
2	B	356	ASP	2.4
2	B	330	TRP	2.3
1	A	292	PRO	2.3
2	B	107	VAL	2.3
2	B	337	VAL	2.3
2	B	251	ASN	2.3
2	B	369	LEU	2.3
2	B	203	GLU	2.2
2	B	139	LEU	2.2
2	B	285	CYS	2.2
2	B	318	GLY	2.2
2	B	124	GLU	2.2
2	B	215	TYR	2.2
2	B	349	PHE	2.2
2	B	140	LEU	2.2
1	A	151	ALA	2.1
2	B	138	PRO	2.1
2	B	211	HIS	2.1
1	A	17[A]	VAL	2.1
2	B	274	ILE	2.1
2	B	242	SER	2.1
2	B	123	LYS	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	218	ASP	2.1
1	A	25	LEU	2.1
2	B	284	LEU	2.1
1	A	72	THR	2.1
2	B	371	ASP	2.0
1	A	295	HIS	2.0
2	B	312	LEU	2.0
2	B	252	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	A	160	11/12	0.95	0.13	52,61,67,67	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	A1CAI	A	301	29/29	0.92	0.12	41,52,71,73	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.