



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 9, 2024 – 09:10 AM EST

PDB ID : 1OCC  
Title : STRUCTURE OF BOVINE HEART CYTOCHROME C OXIDASE AT THE FULLY OXIDIZED STATE  
Authors : Tsukihara, T.; Aoyama, H.; Yamashita, E.; Tomizaki, T.; Yamaguchi, H.; Shinzawa-Itoh, K.; Nakashima, R.; Yaono, R.; Yoshikawa, S.  
Deposited on : 1996-04-18  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)



















The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	514	74% 23% .
1	N	514	75% 22% .
2	B	227	62% 34% .
2	O	227	62% 34% .
3	C	261	72% 24% 5%
3	P	261	72% 23% 5%
4	D	147	73% 22% . .
4	Q	147	71% 24% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	109	 77% 20% •
5	R	109	 78% 20% •
6	F	98	 61% 33% 6%
6	S	98	 64% 30% 6%
7	G	84	 61% 30% 8% •
7	T	84	 60% 31% 8% •
8	H	85	 58% 22% 8% 12%
8	U	85	 58% 21% 9% 12%
9	I	73	 77% 19% •
9	V	73	 75% 21% •
10	J	59	 66% 20% 5% • 5%
10	W	59	 66% 20% 5% • 5%
11	K	56	 64% 23% 12%
11	X	56	 64% 23% 12%
12	L	47	 81% 17% •
12	Y	47	 81% 17% •
13	M	46	 67% 24% • 7%
13	Z	46	 70% 22% • 7%

## 2 Entry composition [i](#)

There are 17 unique types of molecules in this entry. The entry contains 28722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	514	Total	C	N	O	S	0	0	0
			4025	2690	623	677	35			
1	N	514	Total	C	N	O	S	0	0	0
			4025	2690	623	677	35			

- Molecule 2 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	227	Total	C	N	O	S	0	0	0
			1822	1184	281	339	18			
2	O	227	Total	C	N	O	S	0	0	0
			1822	1184	281	339	18			

- Molecule 3 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	261	Total	C	N	O	S	0	0	0
			2124	1420	338	353	13			
3	P	261	Total	C	N	O	S	0	0	0
			2124	1420	338	353	13			

- Molecule 4 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	144	Total	C	N	O	S	0	0	0
			1195	777	196	218	4			
4	Q	144	Total	C	N	O	S	0	0	0
			1195	777	196	218	4			

- Molecule 5 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	109	Total	C	N	O	S	0	0	0
			878	558	150	168	2			
5	R	109	Total	C	N	O	S	0	0	0
			878	558	150	168	2			

- Molecule 6 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	98	Total	C	N	O	S	0	0	0
			748	464	134	145	5			
6	S	98	Total	C	N	O	S	0	0	0
			748	464	134	145	5			

- Molecule 7 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	84	Total	C	N	O	S	0	0	0
			672	431	129	111	1			
7	T	84	Total	C	N	O	S	0	0	0
			672	431	129	111	1			

- Molecule 8 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	75	Total	C	N	O	S	0	0	0
			628	395	114	114	5			
8	U	75	Total	C	N	O	S	0	0	0
			628	395	114	114	5			

- Molecule 9 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	73	Total	C	N	O	S	0	0	0
			598	388	107	99	4			
9	V	73	Total	C	N	O	S	0	0	0
			598	388	107	99	4			

- Molecule 10 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	56	Total	C	N	O	S	0	0	0
			441	285	73	80	3			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	W	56	Total	C	N	O	S	0	0	0
			441	285	73	80	3			

- Molecule 11 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	49	Total	C	N	O	S	0	0	0
			384	250	65	67	2			
11	X	49	Total	C	N	O	S	0	0	0
			384	250	65	67	2			

- Molecule 12 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	47	Total	C	N	O	S	0	0	0
			386	257	65	62	2			
12	Y	47	Total	C	N	O	S	0	0	0
			386	257	65	62	2			

- Molecule 13 is a protein called CYTOCHROME C OXIDASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
13	M	43	Total	C	N	O	0	0	0
			335	223	53	59			
13	Z	43	Total	C	N	O	0	0	0
			335	223	53	59			


- Molecule 14 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	A	1	Total	Cu	0	0
			1	1		
14	B	2	Total	Cu	0	0
			2	2		
14	N	1	Total	Cu	0	0
			1	1		
14	O	2	Total	Cu	0	0
			2	2		

- Molecule 15 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
15	A	1	Total Mg 1 1	0	0
15	N	1	Total Mg 1 1	0	0

- Molecule 16 is HEME-A (three-letter code: HEA) (formula:  $C_{49}H_{56}FeN_4O_6$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	A	1	Total C Fe N O 60 49 1 4 6	0	0
16	A	1	Total C Fe N O 60 49 1 4 6	0	0
16	N	1	Total C Fe N O 60 49 1 4 6	0	0
16	N	1	Total C Fe N O 60 49 1 4 6	0	0

- Molecule 17 is ZINC ION (three-letter code: ZN) (formula:  $Zn$ ).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
17	F	1	Total Zn 1 1	0	0
17	S	1	Total Zn 1 1	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: CYTOCHROME C OXIDASE



#### • Molecule 1: CYTOCHROME C OXIDASE




#### • Molecule 2: CYTOCHROME C OXIDASE









• Molecule 2: CYTOCHROME C OXIDASE



• Molecule 3: CYTOCHROME C OXIDASE




• Molecule 3: CYTOCHROME C OXIDASE




• Molecule 4: CYTOCHROME C OXIDASE

Chain D:  73% 22% • •



• Molecule 4: CYTOCHROME C OXIDASE

Chain Q:  71% 24% • •




• Molecule 5: CYTOCHROME C OXIDASE

Chain E:  77% 20% •




• Molecule 5: CYTOCHROME C OXIDASE

Chain R:  78% 20% •



• Molecule 6: CYTOCHROME C OXIDASE

Chain F:  61% 33% 6%




• Molecule 6: CYTOCHROME C OXIDASE

Chain S:  64% 30% 6%



• Molecule 7: CYTOCHROME C OXIDASE

Chain G:  61% 30% 8% •



• Molecule 7: CYTOCHROME C OXIDASE

Chain T:  60% 31% 8% •




• Molecule 8: CYTOCHROME C OXIDASE

Chain H:  58% 22% 8% 12%




• Molecule 8: CYTOCHROME C OXIDASE

Chain U:  58% 21% 9% 12%



• Molecule 9: CYTOCHROME C OXIDASE

Chain I:  77% 19% •




• Molecule 9: CYTOCHROME C OXIDASE

Chain V:  75% 21% •




• Molecule 10: CYTOCHROME C OXIDASE

Chain J:  66% 20% 5% 5% •



• Molecule 10: CYTOCHROME C OXIDASE

Chain W:  66% 20% 5% 5% •





• Molecule 11: CYTOCHROME C OXIDASE





• Molecule 11: CYTOCHROME C OXIDASE



• Molecule 12: CYTOCHROME C OXIDASE




• Molecule 12: CYTOCHROME C OXIDASE



• Molecule 13: CYTOCHROME C OXIDASE



• Molecule 13: CYTOCHROME C OXIDASE



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	189.10Å 210.50Å 178.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.80	Depositor
% Data completeness (in resolution range)	88.7 (10.00-2.80)	Depositor
$R_{merge}$	0.78	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.201 , 0.252	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	28722	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, HEA, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/4164	0.76	1/5688 (0.0%)
1	N	0.60	0/4164	0.76	1/5688 (0.0%)
2	B	0.57	0/1868	0.79	0/2544
2	O	0.57	0/1868	0.79	0/2544
3	C	0.56	0/2211	0.68	0/3023
3	P	0.56	0/2211	0.68	0/3023
4	D	0.56	0/1229	0.64	1/1658 (0.1%)
4	Q	0.56	0/1229	0.64	1/1658 (0.1%)
5	E	0.50	0/898	0.66	0/1218
5	R	0.50	0/898	0.66	0/1218
6	F	0.56	0/765	0.81	0/1038
6	S	0.56	0/765	0.81	0/1038
7	G	0.54	0/699	0.73	1/950 (0.1%)
7	T	0.55	0/699	0.73	1/950 (0.1%)
8	H	0.55	0/648	0.73	0/877
8	U	0.55	0/648	0.73	0/877
9	I	0.60	0/611	0.64	0/810
9	V	0.60	0/611	0.64	0/810
10	J	0.61	0/451	0.72	0/610
10	W	0.61	0/451	0.72	0/610
11	K	0.57	0/398	0.66	0/546
11	X	0.57	0/398	0.66	0/546
12	L	0.63	0/399	0.62	0/534
12	Y	0.63	0/399	0.62	0/534
13	M	0.51	0/345	0.65	0/470
13	Z	0.51	0/345	0.65	0/470
All	All	0.57	0/29372	0.72	6/39932 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	Q	133	GLY	N-CA-C	6.44	129.21	113.10
4	D	133	GLY	N-CA-C	6.44	129.21	113.10
7	G	5	LYS	N-CA-C	5.22	125.10	111.00
7	T	5	LYS	N-CA-C	5.22	125.10	111.00
1	A	435	GLY	N-CA-C	5.18	126.04	113.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4025	0	4003	91	0
1	N	4025	0	4003	90	0
2	B	1822	0	1834	72	0
2	O	1822	0	1834	73	0
3	C	2124	0	2044	50	0
3	P	2124	0	2044	51	0
4	D	1195	0	1183	32	0
4	Q	1195	0	1183	33	0
5	E	878	0	868	22	0
5	R	878	0	868	20	0
6	F	748	0	728	27	0
6	S	748	0	728	24	0
7	G	672	0	645	40	0
7	T	672	0	645	42	0
8	H	628	0	582	29	0
8	U	628	0	582	30	0
9	I	598	0	612	15	0
9	V	598	0	612	14	0
10	J	441	0	439	14	0
10	W	441	0	439	14	0
11	K	384	0	366	9	0
11	X	384	0	366	10	0
12	L	386	0	388	8	0
12	Y	386	0	388	8	0
13	M	335	0	352	12	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
13	Z	335	0	352	10	0
14	A	1	0	0	0	0
14	B	2	0	0	0	0
14	N	1	0	0	0	0
14	O	2	0	0	0	0
15	A	1	0	0	0	0
15	N	1	0	0	0	0
16	A	120	0	108	8	0
16	N	120	0	108	6	0
17	F	1	0	0	0	0
17	S	1	0	0	0	0
All	All	28722	0	28304	697	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 697 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:T:5:LYS:HZ2	7:T:6:GLY:H	1.13	0.97
1:A:365:ILE:HD12	2:B:87:MET:HE1	1.52	0.90
1:N:365:ILE:HD12	2:O:87:MET:HE1	1.51	0.90
7:G:5:LYS:HZ2	7:G:6:GLY:H	1.14	0.89
3:P:112:LEU:HG	3:P:118:PRO:HB3	1.55	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	512/514 (100%)	479 (94%)	29 (6%)	4 (1%)	16 44

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	N	512/514 (100%)	479 (94%)	29 (6%)	4 (1%)	16	44
2	B	225/227 (99%)	203 (90%)	19 (8%)	3 (1%)	10	32
2	O	225/227 (99%)	203 (90%)	19 (8%)	3 (1%)	10	32
3	C	259/261 (99%)	249 (96%)	10 (4%)	0	100	100
3	P	259/261 (99%)	249 (96%)	10 (4%)	0	100	100
4	D	142/147 (97%)	135 (95%)	7 (5%)	0	100	100
4	Q	142/147 (97%)	135 (95%)	7 (5%)	0	100	100
5	E	107/109 (98%)	104 (97%)	3 (3%)	0	100	100
5	R	107/109 (98%)	104 (97%)	3 (3%)	0	100	100
6	F	96/98 (98%)	86 (90%)	6 (6%)	4 (4%)	2	8
6	S	96/98 (98%)	86 (90%)	6 (6%)	4 (4%)	2	8
7	G	82/84 (98%)	67 (82%)	10 (12%)	5 (6%)	1	3
7	T	82/84 (98%)	67 (82%)	10 (12%)	5 (6%)	1	3
8	H	73/85 (86%)	64 (88%)	8 (11%)	1 (1%)	9	30
8	U	73/85 (86%)	64 (88%)	8 (11%)	1 (1%)	9	30
9	I	71/73 (97%)	65 (92%)	6 (8%)	0	100	100
9	V	71/73 (97%)	65 (92%)	6 (8%)	0	100	100
10	J	54/59 (92%)	48 (89%)	4 (7%)	2 (4%)	2	9
10	W	54/59 (92%)	48 (89%)	4 (7%)	2 (4%)	2	9
11	K	47/56 (84%)	41 (87%)	6 (13%)	0	100	100
11	X	47/56 (84%)	41 (87%)	6 (13%)	0	100	100
12	L	45/47 (96%)	42 (93%)	3 (7%)	0	100	100
12	Y	45/47 (96%)	42 (93%)	3 (7%)	0	100	100
13	M	41/46 (89%)	39 (95%)	2 (5%)	0	100	100
13	Z	41/46 (89%)	39 (95%)	2 (5%)	0	100	100
All	All	3508/3612 (97%)	3244 (92%)	226 (6%)	38 (1%)	12	37

5 of 38 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	328	HIS
1	A	508	PRO
6	F	2	SER

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
6	F	87	THR
6	F	95	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/427 (100%)	389 (91%)	38 (9%)	8	25
1	N	427/427 (100%)	389 (91%)	38 (9%)	8	25
2	B	211/211 (100%)	191 (90%)	20 (10%)	7	22
2	O	211/211 (100%)	191 (90%)	20 (10%)	7	22
3	C	226/226 (100%)	199 (88%)	27 (12%)	4	14
3	P	226/226 (100%)	199 (88%)	27 (12%)	4	14
4	D	128/129 (99%)	120 (94%)	8 (6%)	15	42
4	Q	128/129 (99%)	120 (94%)	8 (6%)	15	42
5	E	95/95 (100%)	89 (94%)	6 (6%)	15	42
5	R	95/95 (100%)	89 (94%)	6 (6%)	15	42
6	F	81/81 (100%)	76 (94%)	5 (6%)	15	43
6	S	81/81 (100%)	76 (94%)	5 (6%)	15	43
7	G	68/68 (100%)	50 (74%)	18 (26%)	0	1
7	T	68/68 (100%)	50 (74%)	18 (26%)	0	1
8	H	67/75 (89%)	58 (87%)	9 (13%)	3	10
8	U	67/75 (89%)	58 (87%)	9 (13%)	3	10
9	I	58/58 (100%)	53 (91%)	5 (9%)	8	27
9	V	58/58 (100%)	53 (91%)	5 (9%)	8	27
10	J	47/50 (94%)	40 (85%)	7 (15%)	2	8
10	W	47/50 (94%)	40 (85%)	7 (15%)	2	8
11	K	39/46 (85%)	37 (95%)	2 (5%)	20	51
11	X	39/46 (85%)	37 (95%)	2 (5%)	20	51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
12	L	40/40 (100%)	38 (95%)	2 (5%)	20	51
12	Y	40/40 (100%)	38 (95%)	2 (5%)	20	51
13	M	37/38 (97%)	34 (92%)	3 (8%)	9	29
13	Z	37/38 (97%)	34 (92%)	3 (8%)	9	29
All	All	3048/3088 (99%)	2748 (90%)	300 (10%)	6	21

5 of 300 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	P	199	VAL
10	W	1	PHE
4	Q	59	LEU
7	T	34	ASN
5	E	80	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 66 such sidechains are listed below:

Mol	Chain	Res	Type
8	U	28	ASN
9	V	8	GLN
13	Z	39	ASN
8	H	28	ASN
8	H	25	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
16	HEA	A	515	1	58,67,67	1.34	7 (12%)	63,103,103	1.44	10 (15%)
16	HEA	A	516	1	58,67,67	1.52	8 (13%)	63,103,103	1.46	13 (20%)
16	HEA	N	515	1	58,67,67	1.34	7 (12%)	63,103,103	1.44	11 (17%)
16	HEA	N	516	1	58,67,67	1.52	8 (13%)	63,103,103	1.46	13 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	HEA	A	515	1	-	5/32/76/76	-
16	HEA	A	516	1	-	5/32/76/76	-
16	HEA	N	515	1	-	5/32/76/76	-
16	HEA	N	516	1	-	5/32/76/76	-

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
16	N	516	HEA	C3A-C2A	-6.65	1.31	1.40
16	A	516	HEA	C3A-C2A	-6.64	1.31	1.40
16	N	515	HEA	C11-C3B	-4.09	1.46	1.51
16	A	515	HEA	C11-C3B	-4.06	1.46	1.51
16	A	516	HEA	C3A-CMA	-3.81	1.37	1.46

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
16	A	515	HEA	C17-C18-C19	-4.17	118.09	127.62

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
16	N	515	HEA	C17-C18-C19	-4.15	118.12	127.62
16	A	516	HEA	C4A-CHB-C1B	3.63	127.35	122.56
16	N	516	HEA	C4A-CHB-C1B	3.63	127.35	122.56
16	A	515	HEA	C13-C14-C15	-3.33	120.00	127.62

There are no chirality outliers.

5 of 20 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
16	A	515	HEA	C15-C16-C17-C18
16	N	515	HEA	C15-C16-C17-C18
16	N	516	HEA	CAD-CBD-CGD-O1D
16	A	516	HEA	CAD-CBD-CGD-O1D
16	A	516	HEA	CAD-CBD-CGD-O2D


There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
16	A	515	HEA	4	0
16	A	516	HEA	4	0
16	N	515	HEA	2	0
16	N	516	HEA	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.